



Prépublications du Département de Mathématiques

Université de La Rochelle
Avenue Michel Crépeau
17042 La Rochelle Cedex 1
<http://www.univ-lr.fr/labo/lmca>

Nonexistence of Global Solutions to an Elliptic Equation with a Dynamical Boundary Condition

Mokhtar Kirane, Eric Nabana et Stanislav I. Pohozaev

Avril 2004

Classification 35J65, 35G30, 35J55.

Mots clés:

2004/06

Nonexistence of global solutions to an elliptic equation with a dynamical boundary condition

Mokhtar KIRANE, Eric NABANA and Stanislav I. POHOZAEV

ABSTRACT. We consider the equation $\Delta u = 0$ posed in $Q := (0, +\infty) \times \Omega$, $\Omega := \{x = (x', x_N)/x' \in R^{N-1}, x_N > 0\}$, with the dynamical boundary condition $B(t, x', 0)u_{tt} + A(t, x', 0)u_t - u_{x_N} \geq D(t, x', 0)|u|^q$ on $\Sigma := (0, \infty) \times R^{N-1} \times \{0\}$ and give conditions on the coefficient functions $A(t, x', 0)$, $B(t, x', 0)$ and $D(t, x', 0)$ for the nonexistence of global solutions.

Contents

1. Introduction
2. Preliminaries
3. The result

1. Introduction

In this paper, we consider the problem **(EI)**

$$\begin{cases} \Delta u = 0, & \text{on } Q, \\ B u_{tt} + A u_t - u_{x_N} \geq D |u|^q, & \text{on } \Sigma, \end{cases}$$

subject to the initial conditions

$$u(0, x', 0) = u_0(x', 0), \quad u_t(0, x', 0) = u_1(x', 0), \quad x' \in R^{N-1}, \quad (1)$$

where $\Delta = \partial_1^2 + \dots + \partial_N^2$ is the Laplace operator with respect to $x = (x', x_N) \in \Omega$, $u_t = \partial u / \partial t$, $u_{x_N} = \partial u / \partial x_N$, and $u_0, u_1 \in L^2_{loc}(R^{N-1} \times \{0\})$, $D \in L^\infty_{loc}(\Sigma)$, $A, A_t, B, B_t, B_{tt} \in L^2_{loc}(\Sigma)$. The functions A, B and D are assumed nonnegative; D positive for large x , A and B don't vanish simultaneously.

Before describing our result in detail, let us dwell on some literature related to equations with dynamical boundary conditions. These kind of problems have been studied for a long time; see [9], [8], [11], [5]. More information is contained in the book by Lions [12]; In chapter 11 of [12], the existence of weak solutions to the Laplace equation with various nonlinear dynamical boundary conditions of parabolic and hyperbolic type is studied. More recently, Kirane [10] considered blow up for three equations with dynamical boundary conditions of parabolic and hyperbolic types. Later, Escher [6] addressed the questions of local solvability and blow up for such problems. Andreucci and Gianni [3] discussed the global existence and blow up issues for a degenerate parabolic problem with nonlocal dynamical

boundary conditions, this on one hand. On the other hand, Apushkinskaya and Nazarov [1] present a survey on the recent results on boundary value problems with boundary conditions described by second-order Venttsel operators. They paid a special attention to nonlinear problems for elliptic and parabolic equations. They stated a priori estimates and existence results in Sobolev and Hölder spaces.

In [2], Amann and Fila derived a Fujita's type result for the Laplace equation with a parabolic dynamical boundary condition with constant coefficients posed in a half-space; they were followed by Fila and Quittner [7] who discussed the same problem as in [2] but in a bounded domain.

In this paper, we generalize the results of [2] concerning blow up to inequalities with non constant coefficients rather than equations with constant coefficients. Observe also that our technique of proof is different of that used by Amann and Fila which parallels Fujita's one; we rather follow an idea from the papers of Baras and Pierre [4], Mitidieri and Pohozaev [13] which is based on a judicious choice of the test function in the weak formulation of the problem, and a scaling of the variables.

2. Preliminaries

The coefficient functions $A(t, x', 0)$, $B(t, x', 0)$ and $D(t, x', 0)$ appearing in the boundary condition are assumed to verify the hypotheses

$$\text{H1. } \int_{R^{N-1}} u(0, x', 0) B(0, x', 0) dx' \geq 0;$$

$$\text{H2. } \int_{R^{N-1}} \left(B_t(0, x', 0) + A_t(0, x', 0) \right) u(0, x', 0) dx' \leq 0;$$

$$\text{H3. } B_{tt} - A_t \leq 0, \text{ in case we consider nonnegative solutions } u \geq 0,$$

or $B_{tt} - A_t = 0$, in case the considered solutions u are of indefinite sign;

$$\text{H4. } |2B_t - A|D^{-\frac{1}{q}} \leq C_1 t^{\sigma_1} |x'|^{\delta_1}, \quad |B|D^{-\frac{1}{q}} \leq C_2 t^{\sigma_2} |x'|^{\delta_2}$$

for $|x'| \gg 1, t > 0$, where $(\delta_1 - \delta_2)/(\sigma_1 - \sigma_2 + 1) > 0$.

H5. The function D is strictly positive for x' outside a large ball.

Definition. By a solution of (EI) on Q subject to the conditions (1), we mean a

function $u \in L^q_{loc}(\Sigma)$ such that

$$\begin{aligned}
\int_{\Sigma} D(t, x', 0) |u|^q \tilde{\varphi} &\leq - \int_{R^{N-1}} u_t(0, x', 0) B(0, x', 0) \tilde{\varphi}(0, x', 0) \\
&+ \int_{R^{N-1}} B_t(0, x', 0) u(0, x', 0) \tilde{\varphi}(0, x', 0) \\
&+ \int_{R^{N-1}} B(0, x', 0) \tilde{\varphi}_t(0, x', 0) u(0, x', 0) \\
&+ \int_{\Sigma} u \left(B_{tt} \tilde{\varphi} + 2B_t \tilde{\varphi}_t + B \tilde{\varphi}_{tt} \right) \\
&- \int_{R^{N-1}} u(0, x', 0) A(0, x', 0) \tilde{\varphi}(0, x', 0) \\
&- \int_{\Sigma} u \left(A_t \tilde{\varphi} + A \tilde{\varphi}_t + \Delta \tilde{\varphi} - \tilde{\varphi}_{x_N} \right) \tag{2}
\end{aligned}$$

for any test function $\tilde{\varphi} \in C_0^2(R_+ \times \Omega)$, $\tilde{\varphi}(T, x', x_N) = 0$ for T large enough.

For later use, we set

$$\tilde{\chi} := (\delta_1 - \delta_2)/(\sigma_1 - \sigma_2 + 1) = \chi/\mu > 0.$$

The Result

Now, we are in force to announce our main result.

Theorem. *Assume that conditions H1-H5 are satisfied, and*

- either (i) $\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi} < 0$, and $q \geq (\tilde{\chi} + N - 1)/(\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi})$;
- or (ii) $\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi} > 0$ and $1 \leq q \leq -(\tilde{\chi} + N - 1)/(\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi})$.

Then problem (EI)-(1) doesn't admit global non trivial solutions.

Proof. The proof is by contradiction. So, we assume that the solution is global. Let $\varphi_0 \in C_0^2(R)$, $\varphi_0 \geq 0$, φ_0 decreasing be such that

$$\varphi_0(\xi) = \begin{cases} 1 & \text{if } 0 \leq |\xi| \leq 1, \\ 0 & \text{if } |\xi| \geq 2. \end{cases}$$

Next, let's define $\psi = \psi(t, x)$ as the solution of

$$\begin{cases} \Delta \psi &= 0, & t \in R_+, x \in R^{N-1} \times R^+, \\ \psi(t, x', 0) &= \varphi(t, x', 0), & t \in R_+, x' \in R^{N-1}. \end{cases}$$

We choose

$$\varphi(t, x') = \varphi_0^\lambda(\xi),$$

where $\xi = \theta^{-1}/(t^\lambda + |x'|^\mu)$, θ a positive real and λ any real greater than p , such that the integrals

$$\int_{\text{supp}\varphi_t} |\varphi_t|^p \varphi^{1-p} \quad \text{and} \quad \int_{\text{supp}\varphi_{tt}} |\varphi_{tt}|^p \varphi^{1-p}$$

are finite.

The function ψ is given by the Poisson's formula

$$\psi(t, x', x_N) = \frac{2x_N}{\sigma_N} \int_{R^{N-1}} \frac{\varphi(t, y, 0)}{(|y - x'|^2 + x_N^2)^{N/2}} dy,$$

where σ_N/N is the volume of the unit ball; so, we have

$$\psi_{x_N}(t, x', 0) = \frac{2}{\sigma_N} \int_{R^{N-1}} \frac{\varphi(t, y, 0)}{|y - x'|^N} dy$$

so

$$-\varphi_{x_N}(t, x', 0) = -\psi_{x_N}(t, x', 0) < 0.$$

Multiplying Equation $(EI)_1$ by ψ and integrating, we obtain

$$\int_Q \Delta u \psi = 0,$$

which, in the light of the Green formula, yields

$$\int_\Sigma u_{x_N}(t, x', 0) \psi(t, x', 0) = \int_\Sigma u(t, x', 0) \psi_{x_N}(t, x', 0).$$

or

$$\int_\Sigma u_{x_N}(t, x', 0) \varphi(t, x', 0) = \int_\Sigma u(t, x', 0) \varphi_{x_N}(t, x', 0).$$

because on Σ , $\psi = \varphi$.

Now, using φ in (2) as a test function and splitting the expression into terms, we obtain

$$\begin{aligned} \int_\Sigma B(t, x', 0) u_{tt} \varphi &= \int_{R^{N-1}} [u_t B \varphi]_0^T - \int_\Sigma (B \varphi)_t u_t \\ &= \int_{R^{N-1}} [u_t B \varphi]_0^T + \int_{R^{N-1}} [u (B \varphi)_t]_0^T + \int_\Sigma u (B \varphi)_{tt} \\ &= - \int_{R^{N-1}} u_t(0, x', 0) B(0, x', 0) \varphi(0, x', 0) \\ &\quad + \int_{R^{N-1}} B_t(0, x', 0) u(0, x', 0) \varphi(0, x', 0) \\ &\quad + \int_{R^{N-1}} B(0, x', 0) \varphi_t(0, x', 0) u(0, x', 0) \\ &\quad + \int_\Sigma u (B_{tt} \varphi + 2B_t \varphi_t + B \varphi_{tt}). \end{aligned}$$

Also, we have

$$\begin{aligned} \int_{\Sigma} A(t, x', 0) u_t \varphi &= \int_{R^{N-1}} [uA\varphi]_0^T - \int_{\Sigma} u(A\varphi)_t \\ &= - \int_{R^{N-1}} u(0, x') A(0, x', 0) \varphi(0, x', 0) - \int_{\Sigma} uA_t \varphi - \int_{\Sigma} uA\varphi_t. \end{aligned}$$

So if we assume $u \geq 0$, we may then write

$$\begin{aligned} \int_{\Sigma} D(t, x', 0) |u|^q \varphi &\leq - \int_{R^{N-1}} u_t(0, x', 0) B(0, x', 0) \varphi(0, x', 0) \\ &\quad + \int_{R^{N-1}} B_t(0, x', 0) u(0, x', 0) \varphi(0, x', 0) \\ &\quad + \int_{R^{N-1}} B(0, x', 0) \varphi_t(0, x', 0) u(0, x', 0) \\ &\quad + \int_{\Sigma} u(B_{tt}\varphi + 2B_t\varphi_t + B\varphi_{tt}) \\ &\quad - \int_{\Sigma} u(0, x', 0) A(0, x', 0) \varphi(0, x', 0) \\ &\quad - \int_{\Sigma} uA_t \varphi - \int_{\Sigma} uA\varphi_t - \int_{R^{N-1}} u\psi_{\sigma}. \end{aligned} \quad (3)$$

In the case where $B_{tt} - A_t = 0$, the solution u may change sign.

By (H1) and (H2), (3) becomes

$$\int_{\Sigma} D(t, x', 0) |u|^q \varphi \leq \int_{\Sigma} u(2B_t - A)\varphi_t + \int_{\Sigma} uB\varphi_{tt}. \quad (4)$$

Using the ε -Young inequality to estimate the right hand side of (4), we obtain

$$\begin{aligned} \int_{\Sigma} D(t, x', 0) |u|^q \varphi &\leq \varepsilon \int_{\Sigma} D(t, x', 0) |u|^q \varphi + C(\varepsilon) \int_{\Sigma} |2B_t - A|^p |\varphi_t|^p D^{1-p} \varphi^{1-p} \\ &\quad + \varepsilon \int_{\Sigma} D(t, x', 0) |u|^q \varphi + C(\varepsilon) \int_{\Sigma} |B|^p |\varphi_{tt}|^p D^{1-p} \varphi^{1-p}, \end{aligned}$$

for some positive constants ε and $C(\varepsilon)$, and where $p + q = pq$.

For ε small enough, we obtain

$$\int_{\Sigma} D |u|^q \varphi \leq C \int_{\Sigma} |2B_t - A|^p |\varphi_t|^p D^{1-p} \varphi^{1-p} + C \int_{\Sigma} |B|^p |\varphi_{tt}|^p D^{1-p} \varphi^{1-p}. \quad (5)$$

The right hand side of (5) is finite thanks to our choice of the test function.

At this stage, we introduce the scaled variables

$$t = \tau \theta^x \quad \text{and} \quad x' = \eta \theta^\mu,$$

so

$$\xi := (t^{\frac{1}{\chi}} + |x'|^{\frac{1}{\mu}})/\theta = \tau^{\frac{1}{\chi}} + |\eta|^{\frac{1}{\mu}}.$$

Observe that

$$\text{supp}\varphi \subset \{(\tau, \eta) \in R^2 : 0 \leq \tau^{\frac{1}{\chi}} + |\eta|^{\frac{1}{\mu}} \leq 2\} =: C_{1,2},$$

$$\text{supp}\varphi_t \subset \{(\tau, \eta) \in R^2 : 1 \leq \tau^{\frac{1}{\chi}} + |\eta|^{\frac{1}{\mu}} \leq 2\}, \quad \text{supp}\varphi_{tt} \subset C_{1,2}.$$

Now, using the scaled variables, we obtain the estimates

$$\begin{aligned} |2B_t - A|^p D^{1-p} &\leq C\theta^{p(\chi\sigma_1 + \mu\delta_1)} \tau^{p\sigma_1} |\eta|^{p\delta_1}, \\ \varphi_t &= \theta^{-p\chi}, \quad \varphi_{tt} = \theta^{-2p\chi}, \quad dt dx' = \theta^{\chi + (N-1)\mu} d\tau d\eta. \end{aligned}$$

So,

$$\int_{\Sigma} |2B_t - A|^p D^{1-p} |\varphi_t|^p \varphi^{1-p} dt dx' \leq C_1 \theta^{\Lambda_1}, \quad (6)$$

where

$$\Lambda_1 = \chi\sigma_1 p + \mu\delta_1 p - \chi p + \chi + \mu(N-1).$$

Similarly, we obtain the estimate

$$\int_{\Sigma} |B|^p D^{1-p} |\varphi_{tt}|^p \varphi^{1-p} \leq C_2 \theta^{\Lambda_2}, \quad (7)$$

where

$$\Lambda_2 = \chi\sigma_2 p + \mu\delta_2 p - 2\chi p + \chi\mu(N-1).$$

Now, we choose $\Lambda_1 \leq 0$ and $\Lambda_2 \leq 0$, that is

$$\begin{aligned} \tilde{\chi}(\sigma_1 p - p + 1) + (\delta_1 p + N - 1) &\leq 0, \\ \tilde{\chi}(\sigma_2 p - 2p + 1) + (\delta_2 p + N - 1) &\leq 0, \end{aligned}$$

where we set $\tilde{\chi} = \chi/\mu$.

For an optimal choice of χ and μ , we take

$$\tilde{\chi} = (\delta_2 - \delta_1)/(\sigma_1 - \sigma_2 + 1).$$

As $\tilde{\chi}$ has to be positive, the inequality $(\delta_2 - \delta_1)(\sigma_1 - \sigma_2 + 1) > 0$ is required.

Finally, we have

$$(\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi})p + (\tilde{\chi} + N - 1) \leq 0, \quad \text{with } \tilde{\chi} = (\delta_2 - \delta_1)/(\sigma_1 - \sigma_2 + 1). \quad (8)$$

There are two possibilities

- either (i) $\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi} < 0 \implies p \geq (\tilde{\chi} + N - 1)/(\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi})$;

- or (ii) $\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi} > 0 \implies 1 \leq p \leq -(\tilde{\chi} + N - 1)/(\tilde{\chi}\sigma_1 + \delta_1 - \tilde{\chi})$.

In the case $\Lambda_1 < 0, \Lambda_2 < 0$, using (6), (7) and (8) we obtain

$$\int_{\Sigma} D|u|^q \leq 0 \implies u = 0.$$

If Λ_1 or $\Lambda_2 = 0$, we have

$$\int_{\Sigma} D|u|^q \leq \text{Const.} < \infty. \quad (9)$$

Now, estimating inequality (4) via the Hölder inequality and using (9), we may write

$$\begin{aligned} \int_{\Sigma} D|u|^q \varphi &\leq \left(\int_{\{\theta \leq t^\chi + |x'|^\mu \leq 2\theta\}} D(t, x') |u|^q \varphi \right)^{\frac{1}{q}} \\ &\cdot \left(\left(\int_{\text{supp} \varphi_t} |2B_t - A|^p \frac{|\varphi_t|^p}{(D\varphi)^{p-1}} \right)^{\frac{1}{p}} + \left(\int_{\text{supp} \varphi_{tt}} |B|^p \frac{|\varphi_{tt}|^p}{(D\varphi)^{p-1}} \right)^{\frac{1}{p}} \right). \end{aligned}$$

As

$$\lim_{\theta \rightarrow \infty} \left(\int_{\{\theta \leq t^\chi + |x'|^\mu \leq 2\theta\}} D(t, x', 0) |u|^q \varphi \right) = 0,$$

we deduce from the former estimate that

$$\int_{\Sigma} D|u|^q \leq 0 \implies u = 0.$$

The proof is complete.

Remark 1. In the case where $B \neq 0$, the natural choice is $\chi = \mu = 2$ because the temporal and spatial derivatives which appear are of the same order; in this case $\tilde{\chi} = 1$ and hence $\sigma_1 + \delta_1 - 1 < 0$; the condition on q is then $q \leq N/(N + 1 - \sigma_1 - \delta_1)$.

Remark 2. Amann and Fila showed that the problem

$$(\mathbf{AF}) \quad \begin{cases} \Delta u = 0, & (0, \infty) \times \Omega, \\ \partial_t u + \partial_n u = u^q, & (0, \infty) \times \partial\Omega, \\ u(0, x') = \phi, & \partial\Omega. \end{cases}$$

admits, for each ϕ bounded and uniformly continuous on R^{N-1} , a unique maximal solution u_ϕ . Moreover, they showed that

- If $T_\phi < \infty$ then $\lim_{t \rightarrow T_\phi} \|u_\phi\|_\infty = \infty$.
- If $q \leq N/(N - 1)$ then every nonzero maximal solution blows up in finite time.
- If $q > N/(N - 1)$ then there are solutions that exist globally as well as solutions that blow up in finite time.

In our notations, Problem (AF) corresponds to

$$B = 0, \quad A = D = 1, \quad \sigma_1 = \delta_1 = 0.$$

In this case, (8) reads

$$-p\tilde{\chi} + \tilde{\chi} + N - 1 \leq 0 \implies \frac{q}{q-1} = p \geq 1 + \frac{N-1}{\tilde{\chi}}.$$

So if we choose $\tilde{\chi} = 1$, then $q \leq N/(N-1)$ for $N \geq 2$; this corresponds to the result of Amann and Fila as expected.

References

- [1] Apushkinskaya, D. E. and A.I. Nazarov, A survey of results on nonlinear Venttsel problems. *Appl. Math.* 45 (2000), no. 1, 69–80.
- [2] Amann, H. and M. Fila, A Fujita-type theorem for the Laplace equation with a dynamical boundary condition. *Acta Math. Univ. Comenian. (N.S.)* 66 (1997), no. 2, 321–328.
- [3] Andreucci, D. and R. Gianni, Global existence and blow up in a parabolic problem with nonlocal dynamical boundary conditions. *Adv. Differential Equations* 1 (1996), no. 5, 729–752.
- [4] Baras, P. and M. Pierre, Problème paraboliques semi-linéaires avec données mesures. *Applicable Analysis* 18 (1984), 11-149.
- [5] Carslaw, H. S. and J. C. Jaeger, *Conduction of heat in solids*. Oxford University Press, 1959, xx+510 pp.
- [6] Escher, J., Nonlinear elliptic systems with dynamical boundary conditions. *Math. Z.* 210 (1992), no. 3, 413–439.
- [7] Fila, M. and P. Quittner, Large time behavior of solutions of a semilinear parabolic equation with a nonlinear dynamical boundary condition. *Topics in nonlinear analysis*, 251–272, *Progr. Nonlinear Differential Equations Appl.*, 35, Birkhäuser, Basel, 1999.
- [8] Friedman, A. and M. Shinbrot, The initial value problem for the linearized equations of water waves. *J. Math. Mech.* 17 (1967) 107–180.
- [9] Garipov, R. M., On the linear theory of gravity waves: the theorem of existence and uniqueness. *Arch. Rational Mech. Anal.* 24 (1967) 352–362.
- [10] Kirane, M., Blow-up for some equations with semilinear dynamical boundary conditions of parabolic and hyperbolic type. *Hokkaido Math. J.* 21 (1992), no. 2, 221–229.

- [11] Langer, R.E., A problem in diffusion or in the flow of heat for a solid in contact with a fluid, *Tohoku Math. J.*, 35 (1932), 260-275.
- [12] Lions, J.-L., Quelques méthodes de résolution des problèmes aux limites non linéaires. Dunod; Gauthier-Villars, Paris 1969 xx+554 pp.
- [13] Mitidieri, E. and S. I. Pohozaev, Absence of positive solutions for systems of quasilinear elliptic equations and inequalities in R^N . (Russian) *Dokl. Akad. Nauk* 366 (1999), no. 1, 13–17.

Mokhtar Kirane

Laboratoire de Mathématiques et Applications, Université de La Rochelle,
Pôle Sciences et Technologie,
Avenue M. Crepeau, 17000 La Rochelle, France
e-mail: mkirane@univ-lr.fr

Eric Nabana

LAMFA, CNRS UPRES A 6119, Université de Picardie Jules Verne,
Faculté de Mathématiques et d'Informatique,
33, rue Saint-Leu, 80039 Amiens Cedex 01, France

Stanislav I. Pohozaev

Steklov Mathematical Institute,
Russian Academy of Sciences ,
Gubkina 8, 117966 Moscow, Russia

Liste des prépublications

- 99-1 Monique Jeanblanc et Nicolas Privault. A complete market model with Poisson and Brownian components. A paraître dans *Proceedings of the Seminar on Stochastic Analysis, Random Fields and Applications*, Ascona, 1999.
- 99-2 Laurence Cherfils et Alain Miranville. Generalized Cahn-Hilliard equations with a logarithmic free energy. A paraître dans *Revista de la Real Academia de Ciencias*.
- 99-3 Jean-Jacques Prat et Nicolas Privault. Explicit stochastic analysis of Brownian motion and point measures on Riemannian manifolds. *Journal of Functional Analysis* **167** (1999) 201-242.
- 99-4 Changgui Zhang. Sur la fonction q -Gamma de Jackson. A paraître dans *Aequationes Math.*
- 99-5 Nicolas Privault. A characterization of grand canonical Gibbs measures by duality. A paraître dans *Potential Analysis*.
- 99-6 Guy Wallet. La variété des équations surstables. A paraître dans *Bulletin de la Société Mathématique de France*.
- 99-7 Nicolas Privault et Jiang-Lun Wu. Poisson stochastic integration in Hilbert spaces. *Annales Mathématiques Blaise Pascal*, **6** (1999) 41-61.
- 99-8 Augustin Fruchard et Reinhard Schäfke. Sursabilité et résonance.
- 99-9 Nicolas Privault. Connections and curvature in the Riemannian geometry of configuration spaces. *C. R. Acad. Sci. Paris, Série I* **330** (2000) 899-904.
- 99-10 Fabienne Marotte et Changgui Zhang. Multisommabilité des séries entières solutions formelles d'une équation aux q -différences linéaire analytique. A paraître dans *Annales de l'Institut Fourier*, 2000.
- 99-11 Knut Aase, Bernt Øksendal, Nicolas Privault et Jan Ubøe. White noise generalizations of the Clark-Haussmann-Ocone theorem with application to mathematical finance. *Finance and Stochastics*, **4** (2000) 465-496.
- 00-01 Eric Benoît. Canards en un point pseudo-singulier nœud. A paraître dans *Bulletin de la Société Mathématique de France*.
- 00-02 Nicolas Privault. Hypothesis testing and Skorokhod stochastic integration. *Journal of Applied Probability*, **37** (2000) 560-574.
- 00-03 Changgui Zhang. La fonction thêta de Jacobi et la sommabilité des séries entières q -Gevrey, I. *C. R. Acad. Sci. Paris, Série I* **331** (2000) 31-34.
- 00-04 Guy Wallet. Déformation topologique par changement d'échelle.
- 00-05 Nicolas Privault. Quantum stochastic calculus for the uniform measure and Boolean convolution. A paraître dans *Séminaire de Probabilités XXXV*.
- 00-06 Changgui Zhang. Sur les fonctions q -Bessel de Jackson.

- 00-07 Laure Coutin, David Nualart et Ciprian A. Tudor. Tanaka formula for the fractional Brownian motion. A paraître dans *Stochastic Processes and their Applications*.
- 00-08 Nicolas Privault. On logarithmic Sobolev inequalities for normal martingales. *Annales de la Faculté des Sciences de Toulouse* **9** (2000) 509-518.
- 01-01 Emanuelle Augeraud-Veron et Laurent Augier. Stabilizing endogenous fluctuations by fiscal policies; Global analysis on piecewise continuous dynamical systems. A paraître dans *Studies in Nonlinear Dynamics and Econometrics*
- 01-02 Delphine Boucher. About the polynomial solutions of homogeneous linear differential equations depending on parameters. A paraître dans *Proceedings of the 1999 International Symposium on Symbolic and Algebraic Computation: ISSAC 99, Sam Dooley Ed., ACM, New York 1999*.
- 01-03 Nicolas Privault. Quasi-invariance for Lévy processes under anticipating shifts.
- 01-04 Nicolas Privault. Distribution-valued iterated gradient and chaotic decompositions of Poisson jump times functionals.
- 01-05 Christian Houdré et Nicolas Privault. Deviation inequalities: an approach via covariance representations.
- 01-06 Abdallah El Hamidi. Remarques sur les sentinelles pour les systèmes distribués
- 02-01 Eric Benoît, Abdallah El Hamidi et Augustin Fruchard. On combined asymptotic expansions in singular perturbation.
- 02-02 Rachid Bebbouchi et Eric Benoît. Equations différentielles et familles bien nées de courbes planes.
- 02-03 Abdallah El Hamidi et Gennady G. Laptev. Nonexistence of solutions to systems of higher-order semilinear inequalities in cone-like domains.
- 02-04 Hassan Lakhel, Youssef Ouknine, et Ciprian A. Tudor. Besov regularity for the indefinite Skorohod integral with respect to the fractional Brownian motion: the singular case.
- 02-05 Nicolas Privault et Jean-Claude Zambrini. Markovian bridges and reversible diffusions with jumps.
- 02-06 Abdallah El Hamidi et Gennady G. Laptev. Existence and Nonexistence Results for Reaction-Diffusion Equations in Product of Cones.
- 02-07 Guy Wallet. Nonstandard generic points.
- 02-08 Gilles Bailly-Maitre. On the monodromy representation of polynomials.
- 02-09 Abdallah El Hamidi. Necessary conditions for local and global solvability of nondiagonal degenerate systems.

- 02-10 Abdallah El Hamidi et Amira Obeid. Systems of Semilinear higher order evolution inequalities on the Heisenberg group.
- 03-01 Abdallah El Hamidi et Gennady G. Laptev. Non existence de solutions d'inéquations semilinéaires dans des domaines coniques.
- 03-02 Eris Benoît et Marie-Joëlle Rochet. A continuous model of biomass size spectra governed by predation and the effects of fishing on them.
- 03-03 Catherine Stenger: On a conjecture of Wolfgang Wasow concerning the nature of turning points.
- 03-04 Christian Houdré et Nicolas Privault. Surface measures and related functional inequalities on configuration spaces.
- 03-05 Abdallah El Hamidi et Mokhtar Kirane. Nonexistence results of solutions to systems of semilinear differential inequalities on the Heisenberg group.
- 03-06 Uwe Franz, Nicolas Privault et René Schott. Non-Gaussian Malliavin calculus on real Lie algebras.
- 04-01 Abdallah El Hamidi. Multiple solutions to a nonlinear elliptic equation involving Paneitz type operators
- 04-02 Mohamed Amara, Amira Obeid et Guy Vallet. Relaxed formulation and existence result of the degenerated elliptic small disturbance model.
- 04-03 Hippolyte d'Albis et Emmanuelle Augeraud-Veron. Competitive Growth in a Life-cycle Model: Existence and Dynamics
- 04-04 Sadjia Aït-Mokhtar: Third order differential equations with fixed critical points.
- 04-05 Mokhtar Kirane et Nasser-eddine Tatar. Asymptotic Behavior for a Reaction Diffusion System with Unbounded Coefficients.
- 04-06 Mokhtar Kirane, Eric Nabana et Stanislav I. Pohozaev. Nonexistence of Global Solutions to an Elliptic Equation with a Dynamical Boundary Condition.