



Prépublications du Département de Mathématiques

Université de La Rochelle
Avenue Michel Crépear
17042 La Rochelle Cedex 1
<http://www.univ-lr.fr/lab0/lmca/>

Some families of polynomial automorphisms

Eric Edo et Jean-Philippe Furter
Mai 2004

Classification: 08A35, 13B25, 14E09, 14J50.

Mots clés:

Some Families of Polynomial Automorphisms.

Eric EDO,

Département de Mathématiques Pures,

Université Bordeaux I,

351 Cours de la Libération, 33 400 Talence, France,

email : Eric.Edo@math.u-bordeaux.fr

Jean-Philippe FURTER,

Département de Mathématiques,

Université de La Rochelle,

avenue M. Crépeau, 17 000 La Rochelle, France,

email : jpfurter@univ-lr.fr

Abstract. We give a family of polynomial automorphisms of the complex affine plane whose generic length is 3 and degenerating in an automorphism of length 1 with surprisingly high degree.

Math. Subj. Class : 08A35; 13B25; 14E09; 14J50

1. Introduction.

The group G of polynomial automorphisms of the complex plane can be endowed with the structure of an infinite-dimensional algebraic variety (see [10]). Furthermore, using the structure of amalgamated product of G , one can define the multidegree of an element σ of G (see below). It is a finite sequence $d = (d_1, \dots, d_l)$ of integers ≥ 2 . We then say that l is the length of σ . A natural question is: What are the relations between the (Zariski) topology of G and the multidegree? A first answer is given in [6]. It is shown that the length is a lower semicontinuous function on the group G (a family of automorphisms of length l can only degenerate in an automorphism of length $l' \leq l$).

Let us denote by G_d the set of automorphisms of G whose multidegree is $d = (d_1, \dots, d_l)$. We obtain a partition of G by the G_d when d describes the set D of finite sequences of integers ≥ 2 . It was conjectured in [6] that $\overline{G_d}$ (the Zariski closure of G_d) is saturated for the equivalence relation “to have the same multidegree”, i.e. there should exist a subset $E(d) \subset D$ of multidegrees $e = (e_1, \dots, e_m)$ such that $\overline{G_d} = \bigcup_{e \in E(d)} G_e$. This is equivalent to saying that if $G_e \cap \overline{G_d} \neq \emptyset$, then $G_e \subset \overline{G_d}$.

In this paper, we show that this conjecture is false. We make explicit a result of

[2] to obtain a family of automorphisms belonging generically to $G_{(11,3,3)}$ and degenerating in an automorphism of $G_{(19)}$. Therefore, we have $G_{(19)} \cap \overline{G_{(11,3,3)}} \neq \emptyset$. However, for grounds of dimension, one cannot have $G_{(19)} \subset \overline{G_{(11,3,3)}}$. Indeed, $G_{(d_1, \dots, d_l)}$ is a constructible subset of G of dimension $d_1 + \dots + d_l + 6$ (see [4] and [5]).

2. The multidegree of an automorphism.

This notion is introduced by Friedland and Milnor (see [4]).

We denote by G the group of polynomial automorphisms of the complex plane $\mathbb{A}_{\mathbb{C}}^2 = \text{Spec}(\mathbb{C}[X, Y])$. An element σ of G is identified with its sequence (f, g) of coordinate functions, where $f, g \in \mathbb{C}[X, Y]$. We define the degree of σ by $\deg \sigma = \max\{\deg(f), \deg(g)\}$. Let $\sigma = (f, g) \in G$ and $\sigma' = (f', g') \in G$. We denote by $\sigma\sigma'$ the composition, *i.e.* $\sigma\sigma' = (f(f', g'), g(f', g'))$.

Let

$$A := \{(aX + bY + c, a'X + b'Y + c'); a, b, c, a', b', c' \in \mathbb{C}, ab' - a'b \neq 0\}$$

be the subgroup of affine automorphisms and let

$$B := \{(aX + P(Y), bY + c); a, b, c \in \mathbb{C}, P \in \mathbb{C}[Y], ab \neq 0\}$$

be the subgroup of triangular automorphisms (B may be viewed as a Borel subgroup of G).

If $\sigma \in G$, by the Jung-van der Kulk theorem (see [7] and [8]), one can write $\sigma = \alpha_1\beta_1 \dots \alpha_k\beta_k\alpha_{k+1}$ where the α_j (resp. β_j) belong to A (resp. B). By contracting such an expression, one might as well suppose that it is reduced, *i.e.* $\forall j, \beta_j \notin A$ and $\forall j, 2 \leq j \leq k, \alpha_j \notin B$. It follows from the amalgamated structure of G that if $\sigma = \alpha'_1\beta'_1 \dots \alpha'_l\beta'_l\alpha'_{l+1}$ is another reduced expression of f , then $k = l$ and there exist $(\gamma_j)_{1 \leq j \leq k}, (\delta_j)_{1 \leq j \leq k}$ in $A \cap B$ such that $\alpha'_1 = \alpha_1\gamma_1^{-1}, \alpha'_j = \delta_{j-1}\alpha_j\gamma_j^{-1}$ (for $2 \leq j \leq k$), $\alpha'_{k+1} = \delta_k\alpha_{k+1}$ and $\beta'_j = \gamma_j\beta_j\delta_j^{-1}$ (for $1 \leq j \leq k$). Following [4], we define the multidegree of σ by $d(\sigma) := (\deg \beta_1, \dots, \deg \beta_k)$ which does not depend on the choice of the reduced expression. Therefore, we have a multidegree function $d : G \rightarrow D$, where D denotes the set of finite sequences of integers ≥ 2 (including the empty sequence).

3. The structure of infinite-dimensional variety of G .

This notion is introduced by Shafarevich (see [10]).

When $n \geq 1$ is an integer, we set $G_{\leq n} := \{\sigma \in G, \deg \sigma \leq n\}$. The subset $G_{\leq n}$ is naturally endowed with the structure of an algebraic variety (see [BCW]). The equality $G = \bigcup_n G_{\leq n}$ endows G with the structure of an infinite-dimensional algebraic

variety.

Let us recall that a set V with a fixed sequence of subsets V_n , each of which has a structure of finite-dimensional algebraic variety, is called an infinite-dimensional algebraic variety if the following conditions are satisfied:

- 1) $V = \bigcup_n V_n$;
- 2) V_n is a closed algebraic subvariety of V_{n+1} .

Each of the V_n will be considered with its Zariski topology and we endow V with the topology of the inductive limit, in which a set $W \subset V$ is closed if and only if $W \cap V_n$ is closed in V_n for each n .

4. Families of automorphisms of the affine plane.

4.1. The Nagata automorphism.

The first non trivial example of a family of automorphisms of the plane comes from the Nagata automorphism (see [9]). Let us set $R = \mathbb{C}[Z]$. The Nagata automorphism is the R -automorphism of $\mathbb{A}_R^2 = \text{Spec } R[X, Y]$ defined by $N = (X - 2Y(XZ + Y^2) - Z(XZ + Y^2)^2, Y + Z(XZ + Y^2))$. We thus obtain a family of automorphisms of $\mathbb{A}_{\mathbb{C}}^2$ parametrized by $\mathbb{A}_{\mathbb{C}}^1 = \text{Spec } \mathbb{C}[Z]$, *i.e.* a morphism from $\mathbb{A}_{\mathbb{C}}^1$ to G .

4.2. The group $G(R)$.

When R is a ring, let us denote by $G(R)$ the group of polynomial R -automorphisms of $\mathbb{A}_R^2 = \text{Spec } R[X, Y]$. We still identify an element σ of $G(R)$ with the couple (f, g) of its coordinate functions, where $f, g \in R[X, Y]$.

Let us notice that knowing an element of $G(R)$ is equivalent to knowing an (algebraic) family of automorphisms parametrized by the affine scheme $V = \text{Spec } R$. In particular, when $R = \mathbb{C}[Z]$, an element σ of $G(\mathbb{C}[Z])$ induces a morphism from $\mathbb{A}_{\mathbb{C}}^1$ to G . We denote by $\sigma_{Z \rightarrow z}$ the image of $z \in \mathbb{A}_{\mathbb{C}}^1 = \mathbb{C}$ by this induced morphism.

For example, if N denotes the Nagata automorphism, we have $N_{Z \rightarrow z} = (X - 2Y(zX + Y^2) - z(zX + Y^2)^2, Y + z(zX + Y^2))$ for all $z \in \mathbb{C}$ and in particular $N_{Z \rightarrow 0} = (X - 2Y^3, Y)$.

4.3. Degeneration.

We shall say that a family $\sigma \in G(\mathbb{C}[Z])$ degenerates at a point z_0 of \mathbb{C} if the multidegree of $\sigma_{Z \rightarrow z_0}$ is different from the generic multidegree of σ , *i.e.* the multidegree of σ as an element of $G(\mathbb{C}(Z))$. For example, one could easily check that the Nagata automorphism splits in $N = (X - Z^{-1}Y^2, Y)(X, Z^2X + Y)(X + Z^{-1}Y^2, Y)$ in $G(\mathbb{C}(Z))$. This last expression shows that the generic multidegree of σ (which is also in this example the multidegree of $N_{Z \rightarrow z}$ for all $z \in \mathbb{C}^*$) is $(2, 2)$. However, the multidegree of $N_{Z \rightarrow 0} = (X - 2Y^3, Y)$ is (3) , hence $G_{(3)} \cap \overline{G_{(2,2)}} \neq \emptyset$.

This family degenerates for $z = 0$ and we observe that the length has strictly

decreased. This illustrates the main result of [6]:

Theorem 1. *The length map $l : G \rightarrow \mathbb{Z}$ is lower semicontinuous.*

However, even if the length behaves well with respect to the topology, it seems that it is not the case of the multidegree (at least in length 3).

4.4. Constructing the elements of $G(\mathbb{C}[Z])$.

The next result (see Lemma 1.1.8. p. 5 of [3]) is very useful to construct the elements of $G(\mathbb{C}[Z])$.

Proposition. *Let $\sigma = (f, g) \in G(\mathbb{C}(Z))$, then the following assertions are equivalent:*

- (i) $\sigma \in G(\mathbb{C}[Z])$;
- (ii) $f, g \in \mathbb{C}[Z][X, Y]$ and $\text{Jac } \sigma \in \mathbb{C}^*$, where $\text{Jac } \sigma := \frac{\partial f}{\partial X} \frac{\partial g}{\partial Y} - \frac{\partial f}{\partial Y} \frac{\partial g}{\partial X}$.

Remark. We reformulate the assertion $f, g \in \mathbb{C}[Z][X, Y]$ by saying that the coefficients of f and g (or even the ones of σ) belong to $\mathbb{C}[Z]$ (the coefficients are of course the ones of $X^p Y^q$).

Furthermore, since $\mathbb{C}(Z)$ is a field, we know (by the Jung-van der Kulk theorem) that any element of $G(\mathbb{C}(Z))$ can be expressed as a composition of affine and triangular automorphisms.

Therefore, to construct any element of $G(\mathbb{C}[Z])$, it is sufficient to compose affine and/or triangular automorphisms $\sigma_1, \dots, \sigma_n$ with coefficients in $\mathbb{C}(Z)$, in such a way that:

- 1) $\sigma = \sigma_1 \cdots \sigma_n$ has its coefficients in $\mathbb{C}[Z]$;
- 2) $\text{Jac } \sigma = \prod_{j=1}^n \text{Jac } \sigma_j \in \mathbb{C}^*$.

For example, using the splitting of the Nagata automorphism in $G(\mathbb{C}(Z))$ (see 4.3.), one can verify assumptions 1) and 2) and the proposition shows that N belongs to $G(\mathbb{C}[Z])$.

5. Some families of automorphisms with generic length 3.

Our main outcome is the following:

Theorem 2. *Let $a, b \geq 2$ and $c \geq 1$ be integers, then*

$$G_{(a+c(ab-1))} \cap \overline{G_{(a+(c-1)(ab-1), b, a)}} \neq \emptyset.$$

Example. Fixing some values for (a, b, c) , we obtain:
 $G_{(5)} \cap \overline{G_{(2,2,2)}} \neq \emptyset$, $G_{(7)} \cap \overline{G_{(2,3,2)}} \neq \emptyset$ and $G_{(19)} \cap \overline{G_{(11,3,3)}} \neq \emptyset$.

>From Theorem 2, we can deduce the

Corollary. Let $a, b \geq 2$ and $c \geq 1$ be integers with $(a, b) \neq (2, 2)$, then the closure of $G_{(a+(c-1)(ab-1), b, a)}$ in G is not saturated for the equivalence relation “to have the same multidegree”.

Proof. Let us recall that $G_{(d_1, \dots, d_l)}$ is a constructible subset of G of dimension $d_1 + \dots + d_l + 6$. Let us set $d = (a + (c-1)(ab-1), b, a)$ and $e = (a + c(ab-1))$. We argue by contradiction. If $\overline{G_d}$ was saturated for the equivalence relation “to have the same multidegree”, then we would have $G_e \subset \overline{G_d} \setminus G_d$ by Theorem 2. Hence the dimension of G_e would be strictly less than that of G_d , whence $a + c(ab-1) + 6 < a + (c-1)(ab-1) + b + a + 6$, i.e. $(a-1)(b-1) < 2$, which is absurd. \square

Example. We have thus shown that $\overline{G_{(2,3,2)}}$ and $\overline{G_{(11,3,3)}}$ are not saturated for the equivalence relation “to have the same multidegree”.

Theorem 2 is a direct consequence of the following one:

Theorem 3. Let $a, b \geq 2$ and $c \geq 1$ be integers.

Let us define triangular automorphisms $\beta, \beta', \beta'' \in G(\mathbb{C}[Z])$ by

$$\beta := (X + Y^a, Y),$$

$$\beta' := (X + ZY^b, Y) \text{ and}$$

$$\beta'' := (X - Y^a \sum_{n=0}^{c-1} \frac{1}{bn+1} \binom{a(bn+1)}{n} (-ZY^{ab-1})^n, Y)$$

and affine automorphisms $\alpha, \alpha' \in G(\mathbb{C}(Z))$ by

$$\alpha := (Z^c X, Y) \text{ and}$$

$$\alpha' := (Y, X).$$

If $\sigma \in G(\mathbb{C}(Z))$ is defined by

$$\sigma := \alpha^{-1} \beta'' \alpha' \beta' \alpha' \beta \alpha,$$

then $\sigma \in G(\mathbb{C}[Z])$ and $\sigma_{Z \rightarrow 0} = (X + \frac{(-1)^c}{bc+1} \binom{a(bc+1)}{c} Y^{a+c(ab-1)}, Y)$.

To obtain Theorem 2 it is enough to remark that the multidegree of σ is generically equal to $(\deg \beta'', \deg \beta', \deg \beta) = (a + (c-1)(ab-1), b, a)$ and that $\sigma_{Z \rightarrow 0}$ is a triangular automorphism of degree $a + c(ab-1)$, so that $d(\sigma_{Z \rightarrow 0}) = (a + c(ab-1))$.

The proof of Theorem 3 is based on the following result:

Lemma 1. Let $a \geq 1$, $b \geq 0$ be integers. Let $S(T) = \sum_{n \geq 0} s_n T^n \in \mathbb{C}[[T]]$ be the power series defined by the functional equation

$$S(T) = (1 + TS^b(T))^a,$$

then

$$(i) \quad S(T) = \sum_{n \geq 0} \frac{1}{bn+1} \binom{a(bn+1)}{n} T^n \text{ and}$$

$$(ii) \quad S(-T(1+T)^{ab-1}) = \frac{1}{(1+T)^a}.$$

Proof. (i) Let us set $u(T) := TS^b(T)$, then $\frac{u(T)}{(1+u(T))^{ab}} = T$, hence u is a local analytic diffeomorphism around 0 with inverse v , where v is defined by $v(W) := \frac{W}{(1+W)^{ab}}$.

It is clear that $s_0 = 1$.

If \oint denotes integration around a little circle around the origin and if $n > 0$ is an integer, it comes out (by the Lagrange formula):

$$\begin{aligned} s_n &= \frac{1}{2\pi i} \oint \frac{S(T)}{T^{n+1}} dT = \frac{1}{2\pi i} \oint \frac{(1+u(T))^a}{T^{n+1}} dT \\ &= \frac{1}{2\pi i} \oint \frac{(1+W)^a}{v^{n+1}(W)} v'(W) dW \quad (\text{setting } T = v(W)) \\ &= \frac{a}{n} \frac{1}{2\pi i} \oint \frac{(1+W)^{a-1}}{v^n(W)} dW \quad (\text{integrating by parts}) \\ &= \frac{a}{n} \frac{1}{2\pi i} \oint \frac{(1+W)^{abn+a-1}}{W^n} dW = \frac{a}{n} \binom{abn+a-1}{n-1} \\ &= \frac{a}{a(bn+1)} \binom{a(bn+1)}{n} = \frac{1}{bn+1} \binom{a(bn+1)}{n}. \end{aligned}$$

(ii) Let us set $h(T) := -T(1+T)^{ab-1}$. Then h is a local analytic diffeomorphism around 0. Let us denote by k its inverse. It is sufficient to show that $S(T) = \frac{1}{(1+k(T))^a}$.

For this, let us set $\tilde{S}(T) := \frac{1}{(1+k(T))^a}$ and let us show that \tilde{S} satisfies the same functional equation than S , that it is to say $\tilde{S}(T) = (1 + T\tilde{S}^b(T))^a$.

This comes from the equality $\frac{1}{1+k(T)} = 1 + \frac{T}{(1+k(T))^{ab}}$, which comes itself from the equality $h(k(T)) = T$. \square

Remarks. 1. Let us set $\lambda = ab$. One could show that the radius of convergence of the power series $S(T)$ is equal to $+\infty$ (resp. 1, resp. $\frac{(\lambda-1)^{\lambda-1}}{\lambda^\lambda}$), if $\lambda = 0$ (resp. $\lambda = 1$, resp. $\lambda \geq 2$).

2. It is pleasant to explain the Lagrange formula by complex analysis; nevertheless, it would have been possible to use a purely algebraic version of this formula (see Corollary 5.4.3. p. 42 in [11]), which would have allowed us to replace the field \mathbb{C} by any field of characteristic 0.

3. Using the functional equation satisfied by $S(T)$, one could show that $S(T) \in \mathbb{Z}[[T]]$. Hence, there exists a version of Theorem 3 where the field \mathbb{C} is replaced by any field (possibly of positive characteristic).

Proof of Theorem 3.

Let us set $P(Y) := Y^a S_{<c}(-ZY^{ab-1}) \in \mathbb{C}[Z][Y]$,

where $S_{<c}(T)$ denotes the series $S(T)$ truncated at the order c ,

that is to say $S_{<c}(T) = \sum_{n=0}^{c-1} s_n T^n$.

We have $\beta'' = (X - P(Y), Y)$ by (i) of Lemma 1.

and if we set $\begin{cases} g := Y + Z(Z^c X + Y^a)^b \\ f := X + Z^{-c}(Y^a - P(g)) \end{cases}$ (1),
then $\sigma = (f, g)$.

We prove that $\sigma \in G(\mathbb{C}[Z])$ using the proposition of § 4.4. It is clear that $\text{Jac } \sigma = 1$ and that $g \in \mathbb{C}[Z][X, Y]$, hence to show that σ is an automorphism of $\mathbb{C}[Z][X, Y]$, it is sufficient to show that $Y^a - P(g)$ is divisible by Z^c (in the ring $\mathbb{C}[Z][X, Y]$), which can be written using congruences $Y^a - P(g) \equiv 0 [Z^c]$.

But, to compute $\sigma_{Z \rightarrow 0}$, we need to compute $Y^a - P(g)$ modulo Z^{c+1} . Therefore, let us carry out directly this computation.

We have $g \equiv Y(1 + ZY^{ab-1}) [Z^{c+1}]$, hence

$$\begin{aligned} Y^a - P(g) &= Y^a - g^a S_{<c}(-Zg^{ab-1}) \\ &\equiv Y^a Q(ZY^{ab-1}) [Z^{c+1}] \end{aligned} \quad (2)$$

where $Q(T) := 1 - (1+T)^a S_{<c}(-T(1+T)^{ab-1})$.

But, by (ii) of Lemma 1, we have:

$$(1+T)^a S(-T(1+T)^{ab-1}) = 1, \text{ hence}$$

$$\begin{aligned}
Q(T) &= (1+T)^a S_{\geq c}(-T(1+T)^{ab-1}), \quad \text{with } S_{\geq c}(T) = \sum_{n \geq c} s_n T^n \\
&\equiv (-1)^c s_c T^c [T^{c+1}], \\
&\equiv \frac{(-1)^c}{bc+1} \binom{a(bc+1)}{c} T^c [T^{c+1}], \quad \text{by (i) of Lemma 1.}
\end{aligned}$$

whence $Y^a - P(g) \equiv \frac{(-1)^c}{bc+1} \binom{a(bc+1)}{c} Z^c Y^{a+c(ab-1)} [Z^{c+1}]$ by (2). By (1) this shows that $f \in \mathbb{C}[Z][X, Y]$ and $\sigma \in G(\mathbb{C}[Z])$ by the proposition of § 4.4. Furthermore we have: $f \equiv X + \frac{(-1)^c}{bc+1} \binom{a(bc+1)}{c} Y^{a+c(ab-1)} [Z]$ by (1) and $g \equiv Y [Z]$ which shows that $\sigma_{Z \rightarrow 0} = (X + \frac{(-1)^c}{bc+1} \binom{a(bc+1)}{c} Y^{a+c(ab-1)}, Y)$. \square

6. The collapse power of a multidegree.

Definition. If $d = (d_1, \dots, d_l)$ is a multidegree with $l \geq 1$, let us define the collapse power of d by $\text{cp}(d) := \max\{k \in \mathbb{Z}, k \geq 2 \text{ and } G_{(k)} \cap \overline{G_d} \neq \emptyset\}$.

Using the well known fact that if the multidegree of σ is (d_1, \dots, d_l) , then the multidegree of σ^{-1} is (d_l, \dots, d_1) , it is easy to show the next result

Lemma 2. *If (d_1, \dots, d_l) is a multidegree, then*

$$\text{cp}(d_1, \dots, d_l) = \text{cp}(d_l, \dots, d_1).$$

Theorem 2 yields a lower bound for $\text{cp}(d_1, d_2, d_3)$.

Theorem 4. *If $(d_1, d_2, d_3) \in D$ is a multidegree such that $d_1 - d_3$ is a multiple of $d_2 \min(d_1, d_3) - 1$, then $\text{cp}(d_1, d_2, d_3) \geq \max(d_1, d_3) + d_2 \min(d_1, d_3) - 1$.*

Proof. By Lemma 2 and since the assumption in Theorem 4 is invariant by exchanging d_1 and d_3 , we can assume that $d_1 \geq d_3$. We set $a = d_3$, $b = d_2$ and $c = 1 + (d_1 - d_3)/(d_2 d_3 - 1)$ then $\max(d_1, d_3) + d_2 \min(d_1, d_3) - 1 = a + c(ab-1)$ and Theorem 4 follows from Theorem 2. \square

Acknowledgments. We would like to thank Mireille Bousquet-Melou for her indications concerning Lemma 1.

References.

- [1] H. Bass, E.H. Connell, D. Wright, The Jacobian Conjecture: Reduction of degree and formal expansion of the inverse, Bull. A.M.S. (1982), 287-330.
- [2] E. Edo, S. Vénéreau, Length 2 variables and transfer, Ann. Pol. Math 76 (2001), 67-76.
- [3] A. van den Essen, Polynomial automorphisms and the Jacobian Conjecture, Progress in Math. 190 (2000) Birkhäuser Verlag, Basel, Boston, Berlin.
- [4] S. Friedland, J. Milnor, Dynamical properties of plane polynomial automorphisms, Ergod. Th. Dyn. Syst 9 (1989), 67-99.
- [5] J-P. Furter, On the variety of automorphisms of the affine plane, J. Algebra 195 (1997), no 2, 604-623.
- [6] J-P. Furter, On the length of polynomial automorphisms of the affine plane, Math. Annalen 322 (2002), 401-411.
- [7] H.W.E. Jung, Über ganze birationale Transformationen der Ebene, J. Reine Angew. Math. 184 (1942), 161-174.
- [8] W. van der Kulk, On polynomial rings in two variables, Nieuw. Arch. Wisk. (3) 1 (1953), 33-41.
- [9] M. Nagata, On automorphism group of $k[x, y]$, Lecture Notes in Math., Kyoto Univ., 5, (1972).
- [10] I.R. Shafarevich, On some infinite-dimensional groups II, Math. USSR Izv, 18 (1982), 214-226.
- [11] R. Stanley, Enumerative Combinatorics, Vol. 2, Cambridge Studies Adv. Math. 62 (1999), Cambridge Univ. Press, Cambridge.

Liste des prépublications

99-1 Monique Jeanblanc et Nicolas Privault. A complete market model with Poisson and Brownian components. A paraître dans *Proceedings of the Seminar on Stochastic Analysis, Random Fields and Applications*, Ascona, 1999.

99-2 Laurence Cherfils et Alain Miranville. Generalized Cahn-Hilliard equations with a logarithmic free energy. A paraître dans *Revista de la Real Academia de Ciencias*.

99-3 Jean-Jacques Prat et Nicolas Privault. Explicit stochastic analysis of Brownian motion and point measures on Riemannian manifolds. *Journal of Functional Analysis* **167** (1999) 201-242.

99-4 Changgui Zhang. Sur la fonction q -Gamma de Jackson. A paraître dans *Acta Mathematica*.

99-5 Nicolas Privault. A characterization of grand canonical Gibbs measures by duality. A paraître dans *Potential Analysis*.

99-6 Guy Wallet. La variété des équations surstables. A paraître dans *Bulletin de la Société Mathématique de France*.

99-7 Nicolas Privault et Jiang-Lun Wu. Poisson stochastic integration in Hilbert spaces. *Annales Mathématiques Blaise Pascal*, **6** (1999) 41-61.

99-8 Augustin Fruchard et Reinhard Schäfke. Sursabilité et résonance.

99-9 Nicolas Privault. Connections and curvature in the Riemannian geometry of configuration spaces. *C. R. Acad. Sci. Paris, Série I* **330** (2000) 899-904.

99-10 Fabienne Marotte et Changgui Zhang. Multisommabilité des séries entières solutions formelles d'une équation aux q -différences linéaire analytique. A paraître dans *Annales de l'Institut Fourier*, 2000.

99-11 Knut Aase, Bernt Øksendal, Nicolas Privault et Jan Ubøe. White noise generalizations of the Clark-Haussmann-Ocone theorem with application to mathematical finance. *Finance and Stochastics*, **4** (2000) 465-496.

00-01 Eric Benoît. Canards en un point pseudo-singulier nœud. A paraître dans *Bulletin de la Société Mathématique de France*.

00-02 Nicolas Privault. Hypothesis testing and Skorokhod stochastic integration. *Journal of Applied Probability*, **37** (2000) 560-574.

00-03 Changgui Zhang. La fonction théta de Jacobi et la sommabilité des séries entières q -Gevrey, I. *C. R. Acad. Sci. Paris, Série I* **331** (2000) 31-34.

00-04 Guy Wallet. Déformation topologique par changement d'échelle.

00-05 Nicolas Privault. Quantum stochastic calculus for the uniform measure and Boolean convolution. A paraître dans *Séminaire de Probabilités XXXV*.

00-06 Changgui Zhang. Sur les fonctions q -Bessel de Jackson.

00-07 Laure Coutin, David Nualart et Ciprian A. Tudor. Tanaka formula for the fractional Brownian motion. A paraître dans *Stochastic Processes and their Applications*.

00-08 Nicolas Privault. On logarithmic Sobolev inequalities for normal martingales. *Annales de la Faculté des Sciences de Toulouse* **9** (2000) 509-518.

01-01 Emanuelle Augeraud-Veron et Laurent Augier. Stabilizing endogenous fluctuations by fiscal policies; Global analysis on piecewise continuous dynamical systems. A paraître dans *Studies in Nonlinear Dynamics and Econometrics*

01-02 Delphine Boucher. About the polynomial solutions of homogeneous linear differential equations depending on parameters. A paraître dans *Proceedings of the 1999 International Symposium on Symbolic and Algebraic Computation: ISSAC 99, Sam Dooley Ed., ACM, New York 1999.*

01-03 Nicolas Privault. Quasi-invariance for Lévy processes under anticipating shifts.

01-04 Nicolas Privault. Distribution-valued iterated gradient and chaotic decompositions of Poisson jump times functionals.

01-05 Christian Houdré et Nicolas Privault. Deviation inequalities: an approach via covariance representations.

01-06 Abdallah El Hamidi. Remarques sur les sentinelles pour les systèmes distribués

02-01 Eric Benoît, Abdallah El Hamidi et Augustin Fruchard. On combined asymptotic expansions in singular perturbation.

02-02 Rachid Bebbouchi et Eric Benoît. Equations différentielles et familles bien nées de courbes planes.

02-03 Abdallah El Hamidi et Gennady G. Laptev. Nonexistence of solutions to systems of higher-order semilinear inequalities in cone-like domains.

02-04 Hassan Lakhel, Youssef Ouknine, et Ciprian A. Tudor. Besov regularity for the indefinite Skorohod integral with respect to the fractional Brownian motion: the singular case.

02-05 Nicolas Privault et Jean-Claude Zambrini. Markovian bridges and reversible diffusions with jumps.

02-06 Abdallah El Hamidi et Gennady G. Laptev. Existence and Nonexistence Results for Reaction-Diffusion Equations in Product of Cones.

02-07 Guy Wallet. Nonstandard generic points.

02-08 Gilles Baily-Maitre. On the monodromy representation of polynomials.

02-09 Abdallah El Hamidi. Necessary conditions for local and global solvability of nondiagonal degenerate systems.

02-10 Abdallah El Hamidi et Amira Obeid. Systems of Semilinear higher order evolution inequalities on the Heisenberg group.

03-01 Abdallah El Hamidi et Gennady G. Laptev. Non existence de solutions d'inéquations semilinéaires dans des domaines coniques.

03-02 Eris Benoît et Marie-Joëlle Rochet. A continuous model of biomass size spectra governed by predation and the effects of fishing on them.

03-03 Catherine Stenger: On a conjecture of Wolfgang Wasow concerning the nature of turning points.

03-04 Christian Houtré et Nicolas Privault. Surface measures and related functional inequalities on configuration spaces.

03-05 Abdallah El Hamidi et Mokhtar Kirane. Nonexistence results of solutions to systems of semilinear differential inequalities on the Heisenberg group.

03-06 Uwe Franz, Nicolas Privault et René Schott. Non-Gaussian Malliavin calculus on real Lie algebras.

04-01 Abdallah El Hamidi. Multiple solutions to a nonlinear elliptic equation involving Paneitz type operators.

04-02 Mohamed Amara, Amira Obeid et Guy Vallet. Relaxed formulation and existence result of the degenerated elliptic small disturbance model.

04-03 Hippolyte d'Albis et Emmanuelle Augeraud-Veron. Competitive Growth in a Life-cycle Model: Existence and Dynamics

04-04 Sadjia Aït-Mokhtar: Third order differential equations with fixed critical points.

04-05 Mokhtar Kirane et Nasser-eddine Tatar. Asymptotic Behavior for a Reaction Diffusion System with Unbounded Coefficients.

04-06 Mokhtar Kirane, Eric Nabana et Stanislav I. Pohozaev. Nonexistence of Global Solutions to an Elliptic Equation with a Dynamical Boundary Condition.

04-07 Khaled M. Furati, Nasser-eddine Tatar and Mokhtar Kirane. Existence and asymptotic behavior for a convection Problem.

04-08 José Alfredo López-Mimbela et Nicolas Privault. Blow-up and stability of semilinear PDE's with gamma generator.

04-09 Abdallah El Hamidi. Multiple solutions with changing sign energy to a non-linear elliptic equation.

04-10 Sadjia Aït-Mokhtar: A singularly perturbed Riccati equation.

04-11 Mohamed Amara, Amira Obeid et Guy Vallet. Weighted Sobolev spaces for a degenerated nonlinear elliptic equation.

04-12 Abdallah El Hamidi. Existence results to elliptic systems with nonstandard growth conditions.

04-13 Eric Edo et Jean-Philippe Furter: Some families of polynomial automorphisms.