



Prépublications du Département de Mathématiques

Université de La Rochelle
Avenue Michel Crépeau
17042 La Rochelle Cedex 1
<http://www.univ-lr.fr/labo/lmca/>

Existence and nonexistence results for higher-order semilinear
evolution inequalities with critical potential

Abdallah El Hamidi et Gennady G. Laptev

Septembre 2004

Classification: 35R45 (35G25 35K55 35L70)

Mots clés: Keywords: blow-up, nonexistence, differential inequality, potential.

Existence and nonexistence results for higher-order semilinear evolution inequalities with critical potential

A. El Hamidi & G. G. Laptev

Abstract

We prove nonexistence results for higher-order semilinear evolution equations and inequalities of the form $\frac{\partial^k u}{\partial t^k} - \Delta u + \frac{\lambda}{|x|^2} u \geq |u|^q$ in $\mathbb{R}^N \times (0, \infty)$, where $\lambda \geq -\left(\frac{N-2}{2}\right)^2$. This problem can be seen as a higher-order evolution version of the nonlinear Wheeler-De Witt equation which appears in the theory of quantum cosmology.

In order to show that our result is sharp in the parabolic case, we establish the existence of positive solutions to the semilinear equation $\frac{\partial u}{\partial t} - \Delta u + \frac{\lambda}{|x|^2} u \geq u^q$ in $\mathbb{R}^N \times (0, \infty)$, for $\lambda \geq 0$. The nonexistence results are based on the test function method, developed by Mitidieri, Pohozaev, Tesei and Véron. The existence result is established by the construction of an explicit global solution of the semilinear parabolic inequality.

Keywords: blow-up, nonexistence, differential inequality, potential

MSC 2000: 35R45 (35G25 35K55 35L70)

1 Introduction

The problem of nonexistence of global solutions to nonlinear partial differential equations is of great interest. We cannot cite all the results in this direction and we refer the reader to [13] and the survey paper [1].

In this note we study the nonexistence of global solutions for higher-order (with respect to t) semilinear partial differential inequalities with critical potential

$$\begin{cases} \frac{\partial^k u}{\partial t^k} - \Delta u + \frac{\lambda}{|x|^2} u \geq |u|^q, & (x, t) \in \mathbb{R}^N \times (0, \infty), \quad \lambda \geq -\left(\frac{N-2}{2}\right)^2, \quad k \geq 1, \\ \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \geq 0, & x \in \mathbb{R}^N, \quad N \geq 3. \end{cases} \quad (1)$$

The well-known operator $-\Delta + \frac{\lambda}{|x|^2}$ (see [24]) appears naturally in the nonlinear Wheeler-De Witt equation which deals with the minisuperspace model in quantum cosmology. We can refer the reader to [4] and the references therein for a complete description of the model. Our problem can be seen then as a higher-order evolution version of the nonlinear Wheeler-De Witt equation.

The nonexistence results hold for q less or equal than critical exponents which depend on N , λ and k . For the parabolic case ($k = 1$), we show that our result is sharp. For that, we construct a global nonnegative solution to the semilinear inequality

$$\frac{\partial u}{\partial t} - \Delta u + \frac{\lambda}{|x|^2} u \geq |u|^q.$$

As for parabolic case ($k = 1$), the paper by Qi Zhang [23] contains an almost complete characterization (for positive solutions) of the relation between the critical exponent of the equation

$$\frac{\partial u}{\partial t} - \Delta u + V(x)u = u^q \quad (2)$$

and the potentials V behaving like $a/(1 + |x|^b)$, where a and b are real numbers. The main Qi Zhang's result is that the relation is quantized depending on b and when $V(x) \sim a/(1 + |x|^2)$ we are at a border-line case where the critical exponent can vary in $(1, \infty]$, and in all other cases the critical exponent takes only three values: 1, ∞ and $1 + \frac{2}{N}$ (the classical Fujita number). So, $V(x)$ with $b = 2$ plays a critical role and is called "the critical potential".

Unfortunately, for general potential $V(x) \sim a/(1 + |x|^2)$ it seems unrealistic to clear cut relation between the critical number and the potential. In the present paper we investigate very particular case [24] when

$$V(x) = \frac{\lambda}{|x|^2}.$$

We want to draw the reader's attention to the fact that we consider solutions without any sign assumption.

Definition 1. Let $u(x, t) \in C(\mathbb{R}^N \times [0, \infty))$ and the locally integrable traces $\frac{\partial^i u}{\partial t^i}(x, 0)$, $i = 1, \dots, k-1$, are well defined. The function $u(x, t)$ is called a weak solution to problem (1) if, for any nonnegative test-function $\varphi(x, t)$, $\frac{\partial^k \varphi}{\partial t^k} \in C(\mathbb{R}^N \times [0, \infty))$, $-\Delta \varphi + \frac{\lambda}{|x|^2} \varphi \in L_1(\mathbb{R}^N \times (0, \infty))$ with compact support, the integral inequality

$$\begin{aligned} \int_0^\infty \int_{\mathbb{R}^N} u \left((-1)^k \frac{\partial^k \varphi}{\partial t^k} - \Delta \varphi + \frac{\lambda}{|x|^2} \varphi \right) dx dt \geq \int_0^\infty \int_{\mathbb{R}^N} |u|^q \varphi dx dt \\ + \sum_{i=1}^{k-1} (-1)^i \int_{\mathbb{R}^N} \frac{\partial^{k-1-i} u}{\partial t^{k-1-i}}(x, 0) \frac{\partial^i \varphi}{\partial t^i}(x, 0) dx + \int_{\mathbb{R}^N} \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \varphi(x, 0) dx \end{aligned} \quad (3)$$

holds.

Let us introduce the parameters

$$s^* = \frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \lambda}, \quad s_* = -\frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \lambda}. \quad (4)$$

Note that s^* and s_* verify

$$\left(-\Delta + \frac{\lambda}{|x|^2}\right) |x|^{s^*} = \left(-\Delta + \frac{\lambda}{|x|^2}\right) |x|^{-s_*} = 0, \quad \text{for any } x \in \mathbb{R}^N \setminus \{0\}.$$

Our result for problem (1) is the following

Theorem 1. *Let*

$$\lambda \geq 0 \quad \text{and} \quad 1 < q \leq q^* = 1 + \frac{2}{s^* + 2/k}$$

or

$$0 > \lambda \geq -\left(\frac{N-2}{2}\right)^2 \quad \text{and} \quad 1 < q \leq q^* = 1 + \frac{2}{-s_* + 2/k}.$$

Then the problem (1) has no nontrivial global solution.

2 Auxiliary estimates

In this section we obtain some estimates depending on the parameter ρ , $\rho \rightarrow \infty$. These estimates play a fundamental role in the test function method [13, 16].

Let us consider the “standard cut-off function” $\zeta(y) \in C^\infty(\mathbb{R}_+)$ with the following properties:

$$0 \leq \zeta(y) \leq 1, \quad \zeta(y) = \begin{cases} 1 & \text{if } 0 \leq y \leq 1, \\ 0 & \text{if } y \geq 2. \end{cases}$$

For the function

$$\eta(y) = (\zeta(y))^{kp_0}$$

with some positive $p_0 > 1$ and $k \in \mathbb{N}$, by direct calculation one can obtain the estimates (for $1 < p \leq p_0$)

$$\begin{aligned} |\eta'(y)|^p &= (kp_0)^p \zeta^{kp_0(p-1)} \zeta^{kp_0-p} |\zeta'|^p \leq c_\eta \eta^{p-1}(y), \\ |\eta''(y)|^p &\leq (kp_0)^p \zeta^{kp_0(p-1)} \zeta^{kp_0-2p} ((kp_0-1)|\zeta'|^2 + \zeta|\zeta''|)^p \leq c_\eta \eta^{p-1}(y), \quad \dots \\ |\eta^{(k)}(y)|^p &\leq c_\eta \eta^{p-1}(y), \end{aligned}$$

with a positive constant c_η .

Now let us introduce the change of variables $y = t/\rho^\theta$, with $\theta > 0$, $\rho > 1$. For the function $\eta(t/\rho^\theta)$ we have

$$\text{supp} \left| \eta \left(\frac{t}{\rho^\theta} \right) \right| = \{0 \leq t \leq 2\rho^\theta\}, \quad \text{supp} \left| \frac{d^k \eta(t/\rho^\theta)}{dt^k} \right| = \{\rho^\theta \leq t \leq 2\rho^\theta\},$$

and

$$\int_{\text{supp} \left| \frac{d^k \eta(t/\rho^\theta)}{dt^k} \right|} \frac{\left| \frac{d^k \eta(t/\rho^\theta)}{dt^k} \right|^p}{\eta^{p-1}(t/\rho^\theta)} dt \leq c_\eta \rho^{-\theta(kp-1)}. \quad (5)$$

The parameter θ will be chosen later.

For the variable x we introduce the functions $\eta(r/\rho)$,

$$\xi(x) \equiv \xi(r) = r^s, \quad (6)$$

and

$$\psi_\rho(x) \equiv \psi_\rho(r) = r^s \eta \left(\frac{r}{\rho} \right). \quad (7)$$

For the derivatives of the function $\psi_\rho(r)$ we have:

$$\begin{aligned} \left| \frac{\partial \psi_\rho}{\partial r} \right|^p &\leq \left| s r^{s-1} \eta \left(\frac{r}{\rho} \right) + r^s \eta' \left(\frac{r}{\rho} \right) \frac{1}{\rho} \right|^p \\ &\leq c \eta^{p-1} \left(\frac{r}{\rho} \right) r^{(s-1)p} \left(1 + \frac{r^p}{\rho^p} \right), \\ \left| \frac{\partial^2 \psi_\rho}{\partial r^2} \right|^p &\leq \left| s(s-1) r^{s-2} \eta \left(\frac{r}{\rho} \right) + \frac{2s r^{s-1}}{\rho} \eta' \left(\frac{r}{\rho} \right) + \frac{r^s}{\rho^2} \eta'' \left(\frac{r}{\rho} \right) \right|^p \\ &\leq c \eta^{p-1} \left(\frac{r}{\rho} \right) r^{(s-2)p} \left(1 + \frac{r^p}{\rho^p} + \frac{r^{2p}}{\rho^{2p}} \right), \end{aligned}$$

here c does not depend on r and ρ . Using these estimates we arrive at the inequality for the operator $A = \Delta - \frac{\lambda}{|x|^2}$:

$$\begin{aligned} |A\psi_\rho(x)|^p &= \left| \Delta \psi_\rho(x) - \frac{\lambda}{|x|^2} \psi_\rho(x) \right|^p \\ &= \left| \frac{\partial^2 \psi_\rho}{\partial r^2} + \frac{N-1}{r} \frac{\partial \psi_\rho}{\partial r} - \frac{\lambda}{r^2} \psi_\rho(x) \right|^p \\ &\leq c \left| \frac{\partial^2 \psi_\rho}{\partial r^2} \right|^p + c \frac{1}{r^p} \left| \frac{\partial \psi_\rho}{\partial r} \right|^p + c \frac{1}{r^{2p}} |\psi_\rho|^p \\ &\leq c \eta^{p-1} \left(\frac{r}{\rho} \right) r^{(s-2)p} \left(1 + \frac{r^p}{\rho^p} + \frac{r^{2p}}{\rho^{2p}} \right) \\ &\leq c \psi_\rho^{p-1}(x) r^{s-2p} \left(1 + \frac{r^p}{\rho^p} + \frac{r^{2p}}{\rho^{2p}} \right). \end{aligned} \tag{8}$$

If $\lambda \geq 0$ then we choose $s = s_* \geq 0$. Due to $A(r^{s_*}) = 0$, we have $A\psi_\rho = 0$ for $r \leq \rho$ and $\text{supp } |A\psi_\rho| \subset \{\rho \leq r \leq 2\rho\}$. On the set $\text{supp } |A\psi_\rho|$ the estimate $1 + \frac{r^p}{\rho^p} + \frac{r^{2p}}{\rho^{2p}} \leq c$ holds, where c does not depend on r and ρ . Therefore, it follows from (8) (for $\rho \leq r \leq 2\rho$) that

$$|A\psi_\rho(x)|^p \leq c \psi_\rho^{p-1}(x) \rho^{s_*-2p};$$

and we get

$$\int_{\text{supp } |A\psi_\rho|} \frac{|A\psi_\rho(x)|^p}{\psi_\rho^{p-1}(x)} dx \leq c \int_\rho^{2\rho} \frac{\psi_\rho^{p-1}(x)}{\psi_\rho^{p-1}(x)} \rho^{s_*-2p} r^{N-1} dr \leq c \psi_\rho^{s_*-2p+N}. \tag{9}$$

For the general test-function

$$\varphi_\rho(x, t) = \eta \left(\frac{t}{\rho^\theta} \right) \psi_\rho(x) \tag{10}$$

we obtain the inequality

$$\begin{aligned} \int_{\text{supp } |A\varphi_\rho|} \frac{|A\varphi_\rho(x, t)|^p}{\varphi_\rho^{p-1}(x, t)} dx dt &\leq \int_0^{2\rho^\theta} \eta(t/\rho^\theta) dt \int_{\text{supp } |A\psi_\rho|} \frac{|A\psi_\rho|^p}{\psi_\rho^{p-1}} dx \\ &\leq c_\varphi \rho^{\theta+s_*-2p+N}. \end{aligned} \tag{11}$$

Analogously, using (5), we obtain:

$$\begin{aligned}
& \int_{\text{supp} \left| \frac{\partial^k \varphi_\rho}{\partial t^k} \right|} \left| \frac{\partial^k \varphi_\rho(x, t)}{\partial t^k} \right|^p \frac{1}{\varphi_\rho^{p-1}(x, t)} dx dt \\
& \leq \int_{\text{supp} \left| \frac{d^k \eta(t/\rho^\theta)}{dt^k} \right|} \left| \frac{d^k \eta(t/\rho^\theta)}{dt^k} \right|^p \frac{1}{\eta^{p-1}(t/\rho^\theta)} dt \int_{0 < |x| < 2\rho} \psi_\rho(x) dx \\
& \leq c_\eta \rho^{-\theta(kp-1)} c \int_0^{2\rho} r^{s_*+N-1} dr \leq c_\varphi \rho^{s_*+N-\theta(kp-1)}. \tag{12}
\end{aligned}$$

For $\theta = 2/k$, the powers in these two estimates are equal:

$$s_* + N - \theta(kp - 1) = \theta + s_* - 2p + N \equiv s_* - 2p + N + 2/k.$$

Finally, we have

$$J_p \equiv \int_{\text{supp} |(-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A\varphi_\rho|} \frac{\left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A\varphi_\rho \right|^p}{\varphi_\rho^{p-1}} dx dt \leq c_0 \rho^{s_* - 2p + N + 2/k}. \tag{13}$$

If $-\left(\frac{N-2}{2}\right)^2 \leq \lambda < 0$, we take $s = -s^* \leq 0$. Therefore, $A(r^{-s^*}) = 0$, $A\psi_\rho = 0$ for $r \leq \rho$ and $\text{supp} |A\psi_\rho| = \{\rho \leq r \leq 2\rho\}$,

$$\int_{\text{supp} |A\psi_\rho|} \frac{|A\psi_\rho(x)|^p}{\psi_\rho^{p-1}(x)} dx \leq c \int_\rho^{2\rho} \frac{\psi_\rho^{p-1}(x)}{\psi_\rho^{p-1}(x)} \rho^{-s^* - 2p} r^{N-1} dr \leq c_\psi \rho^{-s^* - 2p + N} \tag{14}$$

and

$$J_p \equiv \int_{\text{supp} |(-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A\varphi_\rho|} \frac{\left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A\varphi_\rho \right|^p}{\varphi_\rho^{p-1}} dx dt \leq c_0 \rho^{-s^* - 2p + N + 2/k}. \tag{15}$$

3 Proof of theorem 1

Let $u(x, t)$ be a global nontrivial solution of problem (1). From Definition 1 with the test function $\varphi(x, t) = \varphi_\rho(x, t)$, defined by (10) with $p = q' > 1$, $s = s_*$ or $s = -s^*$, and $\theta = 2/k$, using the equalities $\frac{\partial^i \varphi_\rho}{\partial t^i}(x, 0) \equiv 0$, $i = 1, \dots, k-1$, we obtain

$$\begin{aligned}
& \int_{\mathbb{R}^N} \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \varphi_\rho(x, 0) dx + \int_0^\infty \int_{\mathbb{R}^N} |u|^q \varphi_\rho dx dt \\
& \leq \int_{\text{supp} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A\varphi_\rho \right|} u \left((-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A\varphi_\rho \right) dx dt. \tag{16}
\end{aligned}$$

As for the last integral in (16), using the Hölder inequality, we find that

$$\begin{aligned}
& \int_{\mathbb{R}^N} \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \varphi_\rho(x, 0) dx + \int_0^\infty \int_{\mathbb{R}^N} |u|^q \varphi_\rho dx dt \\
&= \int_{\mathbb{R}^N} \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \varphi_\rho(x, 0) dx + \int_{\text{supp} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right|} |u|^q \varphi_\rho dx dt \\
&+ \int_{\varphi_\rho(x,t)=\xi(x)} |u|^q \xi dx dt \\
&\leq \int_{\text{supp} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right|} |u| \cdot \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right| dx dt \\
&\leq \left(\int_{\text{supp} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right|} |u|^q \varphi_\rho dx dt \right)^{1/q} J_{q'}^{1/q'}. \tag{17}
\end{aligned}$$

If $\lambda \geq 0$, using the estimate (13) (with $p = q'$), we have

$$\int_{\varphi_\rho(x,t)=\xi(x)} |u|^q \xi(x) dx dt \leq J_{q'} \leq c_0 \rho^{s_* - 2q' + N + 2/k}. \tag{18}$$

Now we pass to the limit as $\rho \rightarrow \infty$. In the case

$$s_* - 2q' + N + 2/k \leq 0 \tag{19}$$

this implies

$$\int_0^\infty \int_{\mathbb{R}^N} |u|^q \xi dx dt \leq c_0.$$

Then by the inequality $\varphi_\rho \leq \xi$ and taking into account the general properties of Lebesgue integral we have

$$\int_{\text{supp} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right|} |u|^q \varphi_\rho dx dt \leq \int_{\text{supp} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right|} |u|^q \xi dx dt = \varepsilon(\rho) \rightarrow 0$$

as $\rho \rightarrow \infty$. Then from the inequality (17) we finally get

$$\int_{\varphi_\rho(x,t)=\xi(x)} |u|^q \xi dx dt \leq \varepsilon^{1/q}(\rho) c_0^{1/q'} \rightarrow 0$$

as $\rho \rightarrow \infty$, and $\int_0^\infty \int_{\mathbb{R}^N} |u|^q \xi dx dt = 0$, that is, the solution $u(x, t)$ must be trivial under condition (19), which is equivalent to the condition of Theorem 1 for $\lambda \geq 0$.

For $\lambda < 0$ we take $s = -s^*$ and the estimate (15) (with $p = q'$) gives

$$\int_{\varphi_\rho(x,t)=\xi(x)} |u|^q \xi(x) dx dt \leq J_{q'} \leq c_0 \rho^{-s^* - 2q' + N + 2/k}.$$

In the case $-s^* - 2q' + N + 2/k \leq 0$, it implies the nonexistence of nontrivial solution.

4 Generalizations

In this section we present possible generalizations. For simplicity we restrict our attention to the case $\lambda \geq 0$.

Let us consider the inhomogeneous problem [22]

$$\begin{cases} \frac{\partial^k u}{\partial t^k} - \Delta u + \frac{\lambda}{|x|^2} u \geq |u|^q + w(x), & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \geq 0, & x \in \mathbb{R}^N, \end{cases} \quad (20)$$

with $w(x) \in L^1_{\text{loc}}(\mathbb{R}^N)$, $w(x) \geq 0$. We understand the weak solution of this problem in the sense of Definition 1 with the extra term $\int_0^\infty \int_{\mathbb{R}^N} w(x) \varphi \, dx dt$.

Theorem 2. *Let $\lambda \geq 0$ and*

$$1 < q < \frac{s^* + 2}{s^*} = 1 + \frac{2}{s^*}.$$

Then the problem (20) has no nontrivial global solution for any arbitrary small $w(x) \geq 0$, $w(x) \not\equiv 0$.

Proof. Following the proof of Theorem 1, we obtain the a priori estimate

$$\begin{aligned} & \int_0^{\rho^{2/k}} \int_{\{|x| < \rho\}} w(x) \xi \, dx dt + \int_{\{|x| < \rho\}} \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \xi(x) \, dx \\ & + \iint_{\varphi_\rho(x, t) = \xi(x)} |u|^q \xi \, dx dt \leq c_1 \rho^{-2q' + s_* + N + 2/k}, \end{aligned}$$

hence

$$c_1 \rho^{-2q' + s_* + N + 2/k} \geq \int_0^{\rho^{2/k}} \int_{\{|x| < \rho\}} w(x) \xi(x) \, dx dt \geq \rho^{2/k} c_w \quad (21)$$

with ρ such, that

$$\int_{\{|x| < \rho\}} w(x) \xi(x) \, dx \geq c_w \equiv \text{const} > 0.$$

If $-2q' + s_* + N < 0$, then the inequality (21) cannot hold as $\rho \rightarrow \infty$. \square

Finally, we extend our nonexistence results to the following systems

$$\begin{cases} \frac{\partial^k u}{\partial t^k} - \Delta u + \frac{\lambda}{|x|^2} u \geq |v|^{q_1}, & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ \frac{\partial^k v}{\partial t^k} - \Delta v + \frac{\lambda}{|x|^2} v \geq |u|^{q_2}, & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \geq 0, \quad \frac{\partial^{k-1} v}{\partial t^{k-1}}(x, 0) \geq 0, & x \in \mathbb{R}^N. \end{cases} \quad (22)$$

We precise that the sharpness of the result which follows is not established.

Theorem 3. *Let $\lambda \geq 0$, $q_1 > 1$, $q_2 > 1$ and*

$$\max\{\gamma_1, \gamma_2\} \geq \frac{s^* + 2/k}{2}, \quad \text{where} \quad \gamma_1 = \frac{q_1 + 1}{q_1 q_2 - 1}, \quad \gamma_2 = \frac{q_2 + 1}{q_1 q_2 - 1}.$$

Then (22) has no nontrivial global solution.

Proof. In this theorem the concept of solution is understood in the weak sense of Definition 1. Using the Hölder inequality, from the definition we obtain

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \varphi_\rho(x, 0) dx + \int_0^\infty \int_{\mathbb{R}^N} |v|^{q_1} \varphi_\rho dx dt \\ \leq \left(\iint_{\text{supp}} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right| |u|^{q_2} \varphi_\rho dx dt \right)^{1/q_2} J_{q_2}^{1/q_2'}, \end{aligned} \quad (23)$$

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{\partial^{k-1} v}{\partial t^{k-1}}(x, 0) \varphi_\rho(x, 0) dx + \int_0^\infty \int_{\mathbb{R}^N} |u|^{q_2} \varphi_\rho dx dt \\ \leq \left(\iint_{\text{supp}} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right| |v|^{q_1} \varphi_\rho dx dt \right)^{1/q_1} J_{q_1}^{1/q_1'}, \end{aligned} \quad (24)$$

where from (13) we have

$$J_{q_1} \leq c_0 \rho^{-2q_1' + s_* + N + 2/k}, \quad J_{q_2} \leq c_0 \rho^{-2q_2' + s_* + N + 2/k}.$$

We substitute (24) in (23). Then

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{\partial^{k-1} u}{\partial t^{k-1}}(x, 0) \psi_\rho(x) dx + \int_0^\infty \int_{\mathbb{R}^N} |v|^{q_1} \varphi_\rho dx dt \\ \leq \left(\iint_{\text{supp}} \left| (-1)^k \frac{\partial^k \varphi_\rho}{\partial t^k} - A \varphi_\rho \right| |v|^{q_1} \varphi_\rho dx dt \right)^{1/(q_1 q_2)} J_{q_1}^{1/(q_1' q_2)} J_{q_2}^{1/q_2'}, \end{aligned}$$

hence

$$\int_0^\infty \int_{\mathbb{R}^N} |v|^{q_1} \varphi_\rho dx dt \leq \left(J_{q_1}^{q_1-1} J_{q_2}^{q_2(q_2-1)} \right)^{1/(q_1 q_2 - 1)} \leq C \rho^{s_* + 2/k - 2\gamma_1}. \quad (25)$$

Therefore, there is no nontrivial $v(x, t)$ if

$$s_* + 2/k - 2\gamma_1 \leq 0.$$

Analogously, substituting (23) to (24), we deduce

$$\int_0^\infty \int_{\mathbb{R}^N} |u|^{q_2} \varphi_\rho dx dt \leq \left(J_{q_2}^{q_2-1} J_{q_1}^{q_1(q_1-1)} \right)^{1/(q_1 q_2 - 1)} \leq C \rho^{s_* + 2/k - 2\gamma_2}, \quad (26)$$

so that there is no nontrivial $u(x, t)$ if $s_* + 2/k - 2\gamma_2 \leq 0$. It is evidently that if $u(x, t) \equiv 0$, then $v(x, t) \equiv 0$, and vice versa. Finally we get the general nonexistence condition

$$\max\{\gamma_1, \gamma_2\} \geq \frac{s_* + 2/k}{2}.$$

□

5 Existence Results

In this section, we will limit ourselves to the problem

$$\begin{cases} \frac{\partial u}{\partial t} - \Delta u + \frac{\lambda}{|x|^2} u \geq |u|^q & \text{in } \mathbb{R}^N \times]0, +\infty[, \\ u(x, 0) = u_0(x) \geq 0 & \text{in } \mathbb{R}^N, \end{cases} \quad (27)$$

with $\lambda \geq 0$. In this case, Theorem 1 shows that if

$$1 < q \leq q^* = 1 + \frac{2}{N + s_*}$$

then the problem (27) has no global nontrivial solution [3, 12]. We complete this nonexistence result by the existence one:

Theorem 4. *If*

$$q > q^* \equiv 1 + \frac{2}{N + s_*}$$

then nontrivial global solutions of (27) exist.

Proof. Let v be a positive solution of

$$\begin{cases} \frac{\partial v}{\partial t} - \Delta v + \frac{\lambda}{|x|^2} v = 0 & \text{in } \mathbb{R}^N \times]0, +\infty[, \\ v(x, 0) = v_0(x) \geq 0 & \text{in } \mathbb{R}^N, \end{cases}$$

and let the function w defined on $\mathbb{R}^N \times]0, +\infty[$ by $w(x, t) = \alpha(t)v(x, t)$, where the function α has to be defined. If α is selected such that

$$\alpha'(t) = (\alpha(t))^q \left[\sup_{x \in \mathbb{R}^N} v(x, t) \right]^{q-1} = (\alpha(t))^q \|v(\cdot, t)\|_{L^\infty(\mathbb{R}^N)}^{q-1},$$

then w is a solution of (27) on its interval of definition. Let us set α be the solution of the Cauchy problem

$$\begin{cases} \alpha'(t) = (\alpha(t))^q \|v(\cdot, t)\|_{L^\infty(\mathbb{R}^N)}^{q-1}, & t > 0, \\ \alpha(0) = \alpha_0 > 0. \end{cases} \quad (28)$$

It is easy to see that the solution of (28) is global if, and only if,

$$\int_0^{+\infty} \|v(\cdot, t)\|_{L^\infty(\mathbb{R}^N)}^{q-1} dt < +\infty \quad (29)$$

and

$$0 < \alpha_0 < \left((q-1) \int_0^{+\infty} \|v(\cdot, t)\|_{L^\infty(\mathbb{R}^N)}^{q-1} dt \right)^{\frac{-1}{q-1}}.$$

At this stage, we will construct the function v on $\mathbb{R}^N \times]0, +\infty[$. Consider the function v defined on $\mathbb{R}^N \times]0, +\infty[$ by

$$v(x, t) = \frac{1}{t+1} \frac{1}{|x|^{\frac{1}{2}(N-2)}} I_\nu \left(\frac{|x|}{2(t+1)} \right) \exp \left(-\frac{|x|^2 + 1}{4(t+1)} \right),$$

where $\nu = s_* + \frac{N-2}{2}$ and I_ν is the modified Bessel function of order ν [21]. One can observe by direct calculation that the function v is a positive solution of

$$\frac{\partial v}{\partial t} - \Delta v + \frac{\lambda}{|x|^2} v = 0 \text{ in } \mathbb{R}^N \times]0, +\infty[.$$

Recall that the asymptotic behaviour of I_ν in the neighborhood of 0 and $+\infty$ is given respectively by [21]

$$I_\nu(z) \sim \frac{z^\nu}{2^\nu \Gamma(\nu + 1)} \text{ as } z \rightarrow 0^+ \quad (30)$$

and

$$I_\nu(z) \sim \frac{\exp(z)}{\sqrt{2\pi z}} \text{ as } z \rightarrow +\infty. \quad (31)$$

Moreover, if the following estimate

$$\limsup_{t \rightarrow +\infty} (t+1)^{\frac{1}{q^*-1}} \|v(\cdot, t)\|_{L^\infty(\mathbb{R}^N)} < +\infty \quad (32)$$

holds then the condition (29) will be satisfied for any $q > q^*$. Indeed, it suffices to remark that

$$\int_0^{+\infty} (t+1)^{-\frac{q-1}{q^*-1}} dt < +\infty, \text{ for any } q > q^*.$$

We will show now that the estimate (32) is satisfied. First, since

$$\lim_{|x| \rightarrow 0} v(x, t) = \lim_{|x| \rightarrow \infty} v(x, t) = 0, \quad \forall t > 0,$$

then, for any $t > 0$, there exists $r^*(t) \in]0, +\infty[$, such that

$$v(r^*(t), t) = \|v(\cdot, t)\|_{L^\infty(\mathbb{R}^N)}.$$

Let

$$\mathcal{V}(t) = (t+1)^{\frac{1}{q^*-1}} v(r^*(t), t).$$

Using the fact that

$$\frac{1}{q^* - 1} = \frac{N + s_*}{2},$$

we can write

$$\mathcal{V}(t) = (t+1)^{\frac{s_*}{2}} \left(\frac{r^*(t)}{t+1}\right)^{-\frac{N-2}{2}} I_\nu\left(\frac{r^*(t)}{2(t+1)}\right) \exp\left(-\frac{(r^*(t))^2 + 1}{4(t+1)}\right).$$

If we set

$$y^*(t) = \frac{r^*(t)}{2(t+1)},$$

then

$$\mathcal{V}(t) = e^{-\frac{1}{4(t+1)}} (t+1)^{\frac{s_*}{2}} (y^*(t))^{-\frac{N-2}{2}} I_\nu(y^*(t)) e^{-(t+1)(y^*(t))^2}.$$

Suppose that there is a sequence $(t_k)_{k \in \mathbb{N}} \rightarrow +\infty$ such that

$$\lim_{t_k \rightarrow +\infty} \mathcal{V}(t_k) = +\infty.$$

Three cases can arise:

- Case 1: There exists a subsequence, also denoted by $(t_k)_{k \in \mathbb{N}}$, such that

$$\lim_{t_k \rightarrow +\infty} y^*(t_k) = +\infty.$$

In this case

$$\mathcal{V}(t_k) \sim \text{const} \cdot (t_k + 1)^{\frac{s^*}{2}} (y^*(t_k))^{-\frac{N-1}{2}} e^{y^*(t_k)} e^{-(t_k+1)(y^*(t_k))^2} \quad \text{as } t_k \rightarrow +\infty,$$

which implies that

$$\lim_{t_k \rightarrow \infty} \mathcal{V}(t_k) = 0.$$

- Case 2: There exists a subsequence, also denoted by $(t_k)_{k \in \mathbb{N}}$, such that

$$\lim_{t_k \rightarrow +\infty} y^*(t_k) = 0.$$

In this case

$$\mathcal{V}(t_k) \sim \text{const} \cdot [(t_k + 1)(y^*(t_k))^2]^{\frac{s^*}{2}} e^{-(t_k+1)(y^*(t_k))^2} \quad \text{as } t_k \rightarrow +\infty,$$

which implies that $\mathcal{V}(t_k)$ is bounded, since the function $z \mapsto z^{\frac{s^*}{2}} e^{-z}$ is bounded on \mathbb{R}^+ .

- Case 3: There are two constants A and B such that the sequence $(y^*(t_k))_{k \in \mathbb{N}}$ satisfies

$$0 < A \leq y^*(t_k) \leq B < +\infty.$$

In this case, the expression $\mathcal{V}(t_k)$ is clearly bounded.

Whence, there is no subsequence of $(t_k)_{k \in \mathbb{N}} \rightarrow +\infty$ such that

$$\lim_{t_k \rightarrow +\infty} \mathcal{V}(t_k) = +\infty,$$

which implies that there is no sequence $(t_k)_{k \in \mathbb{N}} \rightarrow +\infty$ such that

$$\lim_{t_k \rightarrow +\infty} \mathcal{V}(t_k) = +\infty.$$

This ends the proof. □

References

- [1] DENG K. & LEVINE H.A., The role of critical exponents in blow-up theorems: the sequel. *J. Math. Anal. Appl.* **243** (2000), 85–126.
- [2] C. Bandle & M. Essén, On positive solutions of Emden equations in cone-like domains. *Arch. Rational Mech. Anal.* 112 (1990), no. 4, 319–338.
- [3] C. Bandle & H. Levine, A Fujita type phenomena for reaction-diffusion equations with convection like terms. *Differential Integral Equations* 7 (1994), no. 5-6, 1169–1193.

- [4] H. Berestycki & M. J. Esteban, Existence and bifurcation of solutions for an elliptic degenerate problem. *J. Differential Equations* 134 (1997), no. 1, 1–25.
- [5] El HAMIDI, A., Necessary conditions for local and global solvability of non-diagonal degenerate systems. *Z. Anal. Anwendungen* **22** (2003), 109–127.
- [6] El HAMIDI, A. & LAPTEV, G. G., Nonexistence of solutions to systems of higher-order semilinear inequalities in cone-like domains. *Electron. J. Differential Equations* **97** (2002)
- [7] El HAMIDI, A. & OBEID, A., Systems of semilinear higher-order evolution inequalities on the Heisenberg group. *J. Math. Anal. Appl.* **280** (2003), 77–90.
- [8] KARTSATOS A.G. & KURTA V.V., On the critical Fujita exponents for solutions of quasilinear parabolic inequalities. *J. Math. Anal. Appl.* **269** (2002), 73–86.
- [9] KURTA V.V., On the absence of positive solutions to semilinear elliptic equations. *Proc. Steklov Inst. Math.* **227** (1999), 162–169.
- [10] LAPTEV G.G., On the absence of solutions to a class of singular semilinear differential inequalities. *Proc. Steklov Inst. Math.* **232** (2001), 223–235.
- [11] LAPTEV G.G., Nonexistence results for higher-order evolution partial differential inequalities. *Proc. Amer. Math. Soc.* **131** (2003), 415–423.
- [12] H. Levine & P. Meier, The value of the critical exponent for reaction-diffusion equations in cones. *Arch. Rational Mech. Anal.* 109 (1990), no. 1, 73–80.
- [13] MITIDIERI E. & POHOZAEV S.I., Nonexistence of global positive solutions to quasilinear elliptic inequalities. *Dokl. Russ. Acad. Sci.* **57** (1998), 250–253.
- [14] MITIDIERI E. & POHOZAEV S.I., Nonexistence of weak solutions for some degenerate elliptic and parabolic problems on \mathbb{R}^n . *J. Evolution Equations* **1** (2001), 189–220.
- [15] MITIDIERI E. & POHOZAEV S.I., Nonexistence of weak solutions for some degenerate and singular hyperbolic problems on \mathbb{R}^N . *Proc. Steklov Inst. Math.* **232** (2001), 240–259.
- [16] POHOZAEV S.I. & TESEI A., Critical exponents for the absence of solutions for systems of quasilinear parabolic inequalities. *Differ. Uravn.* **37** (2001), 521–528.
- [17] POHOZAEV S.I. & VERON L., Nonexistence results of solutions of semilinear differential inequalities on the Heisenberg group. *Manuscripta math.* **102** (2000), 85–99.
- [18] SAMARSKII A.A., GALAKTIONOV V.A., KURDYUMOV S.P. & MIKHAILOV A.P., *Blow-up in Quasilinear Parabolic Equations*, Walter de Gruyter, Berlin/New York, 1995.
- [19] SATTINGER D.H., *Topics in Stability and Bifurcation Theory*. Lect. Notes Math., **309**, Springer, New York, N. Y., 1973.
- [20] SATTINGER D.H., Monotone methods in nonlinear elliptic and parabolic boundary value problems. *Indiana U. Math. J.* **21** (1972), 979–1000.

- [21] WATSON G.N., *A treatise on the Theory of Bessel Functions*. Cambridge University Press, London/New York, 1966.
- [22] ZHANG QI, Blow-up results for nonlinear parabolic equations on manifolds. *Duke Math. J.* **97** (1999), 515–539.
- [23] ZHANG QI, The quantizing effect of potentials on the critical number of reaction–diffusion equations. *J. Differential Equations* **170** (2001), 188–214.
- [24] ZHANG QI, Global lower bound for the heat kernel of $-\Delta + \frac{c}{|x|^2}$. *Proc. Amer. Math. Soc.* **129** (2001), 1105–1112.

ABDALLAH EL HAMIDI
Université de La Rochelle,
Laboratoire de Mathématiques et Applications
Avenue Michel Crépeau
F-17000 La Rochelle
e-mail: aelhamid@univ-lr.fr

GENNADY G. LAPTEV
Department of Function Theory
Steklov Mathematical Institute
Gubkina 8, 119991 Moscow, Russia
e-mail: laptev@home.tula.net

Liste des prépublications

- 99-1 Monique Jeanblanc et Nicolas Privault. A complete market model with Poisson and Brownian components. A paraître dans *Proceedings of the Seminar on Stochastic Analysis, Random Fields and Applications*, Ascona, 1999.
- 99-2 Laurence Cherfils et Alain Miranville. Generalized Cahn-Hilliard equations with a logarithmic free energy. A paraître dans *Revista de la Real Academia de Ciencias*.
- 99-3 Jean-Jacques Prat et Nicolas Privault. Explicit stochastic analysis of Brownian motion and point measures on Riemannian manifolds. *Journal of Functional Analysis* **167** (1999) 201-242.
- 99-4 Changgui Zhang. Sur la fonction q -Gamma de Jackson. A paraître dans *Aequationes Math.*
- 99-5 Nicolas Privault. A characterization of grand canonical Gibbs measures by duality. A paraître dans *Potential Analysis*.
- 99-6 Guy Wallet. La variété des équations surstables. A paraître dans *Bulletin de la Société Mathématique de France*.
- 99-7 Nicolas Privault et Jiang-Lun Wu. Poisson stochastic integration in Hilbert spaces. *Annales Mathématiques Blaise Pascal*, **6** (1999) 41-61.
- 99-8 Augustin Fruchard et Reinhard Schäfke. Sursabilité et résonance.
- 99-9 Nicolas Privault. Connections and curvature in the Riemannian geometry of configuration spaces. *C. R. Acad. Sci. Paris, Série I* **330** (2000) 899-904.
- 99-10 Fabienne Marotte et Changgui Zhang. Multisommabilité des séries entières solutions formelles d'une équation aux q -différences linéaire analytique. A paraître dans *Annales de l'Institut Fourier*, 2000.
- 99-11 Knut Aase, Bernt Øksendal, Nicolas Privault et Jan Ubøe. White noise generalizations of the Clark-Haussmann-Ocone theorem with application to mathematical finance. *Finance and Stochastics*, **4** (2000) 465-496.
- 00-01 Eric Benoît. Canards en un point pseudo-singulier nœud. A paraître dans *Bulletin de la Société Mathématique de France*.
- 00-02 Nicolas Privault. Hypothesis testing and Skorokhod stochastic integration. *Journal of Applied Probability*, **37** (2000) 560-574.
- 00-03 Changgui Zhang. La fonction théta de Jacobi et la sommabilité des séries entières q -Gevrey, I. *C. R. Acad. Sci. Paris, Série I* **331** (2000) 31-34.
- 00-04 Guy Wallet. Déformation topologique par changement d'échelle.
- 00-05 Nicolas Privault. Quantum stochastic calculus for the uniform measure and Boolean convolution. A paraître dans *Séminaire de Probabilités XXXV*.
- 00-06 Changgui Zhang. Sur les fonctions q -Bessel de Jackson.

- 00-07 Laure Coutin, David Nualart et Ciprian A. Tudor. Tanaka formula for the fractional Brownian motion. A paraître dans *Stochastic Processes and their Applications*.
- 00-08 Nicolas Privault. On logarithmic Sobolev inequalities for normal martingales. *Annales de la Faculté des Sciences de Toulouse* **9** (2000) 509-518.
- 01-01 Emanuelle Augeraud-Veron et Laurent Augier. Stabilizing endogenous fluctuations by fiscal policies; Global analysis on piecewise continuous dynamical systems. A paraître dans *Studies in Nonlinear Dynamics and Econometrics*
- 01-02 Delphine Boucher. About the polynomial solutions of homogeneous linear differential equations depending on parameters. A paraître dans *Proceedings of the 1999 International Symposium on Symbolic and Algebraic Computation: ISSAC 99, Sam Dooley Ed., ACM, New York 1999*.
- 01-03 Nicolas Privault. Quasi-invariance for Lévy processes under anticipating shifts.
- 01-04 Nicolas Privault. Distribution-valued iterated gradient and chaotic decompositions of Poisson jump times functionals.
- 01-05 Christian Houdré et Nicolas Privault. Deviation inequalities: an approach via covariance representations.
- 01-06 Abdallah El Hamidi. Remarques sur les sentinelles pour les systèmes distribués
- 02-01 Eric Benoît, Abdallah El Hamidi et Augustin Fruchard. On combined asymptotic expansions in singular perturbation.
- 02-02 Rachid Bebbouchi et Eric Benoît. Equations différentielles et familles bien nées de courbes planes.
- 02-03 Abdallah El Hamidi et Gennady G. Laptev. Nonexistence of solutions to systems of higher-order semilinear inequalities in cone-like domains.
- 02-04 Hassan Lakhel, Youssef Ouknine, et Ciprian A. Tudor. Besov regularity for the indefinite Skorohod integral with respect to the fractional Brownian motion: the singular case.
- 02-05 Nicolas Privault et Jean-Claude Zambrini. Markovian bridges and reversible diffusions with jumps.
- 02-06 Abdallah El Hamidi et Gennady G. Laptev. Existence and Nonexistence Results for Reaction-Diffusion Equations in Product of Cones.
- 02-07 Guy Wallet. Nonstandard generic points.
- 02-08 Gilles Bailly-Maitre. On the monodromy representation of polynomials.
- 02-09 Abdallah El Hamidi. Necessary conditions for local and global solvability of nondiagonal degenerate systems.
- 02-10 Abdallah El Hamidi et Amira Obeid. Systems of Semilinear higher order evolution inequalities on the Heisenberg group.
- 03-01 Abdallah El Hamidi et Gennady G. Laptev. Non existence de solutions d'inéquations semilinéaires dans des domaines coniques.

- 03-02 Eris Benoît et Marie-Joëlle Rochet. A continuous model of biomass size spectra governed by predation and the effects of fishing on them.
- 03-03 Catherine Stenger: On a conjecture of Wolfgang Wasow concerning the nature of turning points.
- 03-04 Christian Houdré et Nicolas Privault. Surface measures and related functional inequalities on configuration spaces.
- 03-05 Abdallah El Hamidi et Mokhtar Kirane. Nonexistence results of solutions to systems of semilinear differential inequalities on the Heisenberg group.
- 03-06 Uwe Franz, Nicolas Privault et René Schott. Non-Gaussian Malliavin calculus on real Lie algebras.
- 04-01 Abdallah El Hamidi. Multiple solutions to a nonlinear elliptic equation involving Paneitz type operators.
- 04-02 Mohamed Amara, Amira Obeid et Guy Vallet. Relaxed formulation and existence result of the degenerated elliptic small disturbance model.
- 04-03 Hippolyte d'Albis et Emmanuelle Augeraud-Veron. Competitive Growth in a Life-cycle Model: Existence and Dynamics
- 04-04 Sadjia Aït-Mokhtar: Third order differential equations with fixed critical points.
- 04-05 Mokhtar Kirane et Nasser-eddine Tatar. Asymptotic Behavior for a Reaction Diffusion System with Unbounded Coefficients.
- 04-06 Mokhtar Kirane, Eric Nabana et Stanislav I. Pohozaev. Nonexistence of Global Solutions to an Elliptic Equation with a Dynamical Boundary Condition.
- 04-07 Khaled M. Furati, Nasser-eddine Tatar and Mokhtar Kirane. Existence and asymptotic behavior for a convection Problem.
- 04-08 José Alfredo López-Mimbela et Nicolas Privault. Blow-up and stability of semilinear PDE's with gamma generator.
- 04-09 Abdallah El Hamidi. Multiple solutions with changing sign energy to a nonlinear elliptic equation.
- 04-10 Sadjia Aït-Mokhtar: A singularly perturbed Riccati equation.
- 04-11 Mohamed Amara, Amira Obeid et Guy Vallet. Weighted Sobolev spaces for a degenerated nonlinear elliptic equation.
- 04-12 Abdallah El Hamidi. Existence results to elliptic systems with nonstandard growth conditions.
- 04-13 Eric Edo et Jean-Philippe Furter: Some families of polynomial automorphisms.
- 04-14 Laurence Cherfils et Yavdat Il'yasov. On the stationary solutions of generalized reaction diffusion equations with p & q - Laplacian.

- 04-15 Jean-Christophe Breton et Youri Davydov. Local limit theorem for supremum of an empirical processes for i.i.d. random variables.
- 04-16 Jean-Christophe Breton, Christian Houdré et Nicolas Privault. Dimension free and infinite variance tail estimates on Poisson space.
- 04-17 Abdallah El Hamidi et Gennady G. Laptev. Existence and nonexistence results for higher-order semilinear evolution inequalities with critical potential.