



Prépublications du Département de Mathématiques

Université de La Rochelle
Avenue Michel Crépeau
17042 La Rochelle Cedex 1
<http://www.univ-lr.fr/labo/lmca/>

Compactness and quasilinear problems with critical exponents

Abdallah El Hamidi et Jean-Michel Rakotoson

Juin 2005

Classification: 35J60,35J25,35J65,35J85.

Mots clés: Quasilinear problems, compactness, critical Sobolev exponents.

2005/04

Dedicated to Professor Roger Temam for his 65th anniversary.

Compactness and quasilinear problems with critical exponents

A. EL Hamidi(1)

(1) Laboratoire de Mathématiques, Université de La Rochelle
Av. Michel Crépeau, 17042 LA ROCHELLE Cedex 09 - France

J.M. Rakotoson(2)

(2) Laboratoire de Mathématiques - Université de Poitiers - SP2MI -
Boulevard Marie et Pierre Curie, Téléport 2
BP30179 86962 FUTUROSCOPE CHASSENEUIL Cedex - France.

Keywords : Quasilinear problems, compactness, critical Sobolev exponents.

Abstract

A compactness result is revised in order to prove the pointwise convergence of the gradients of a sequence of solutions to a general quasilinear inequality (anisotropic or not, degenerate or not) and for an arbitrary open set. Combining this result with the well-known Brézis-Lieb lemma, we derive simple proofs of Palais-Smale properties in many optimization problems especially on unbounded domains.

1 Introduction.

In his recent works [3, 4], the first author observed that the pointwise convergence obtained by the second author for measure data problems [12, 13, 14, 15], combined with the well-known Brézis-Lieb lemma [2], can be helpful for proving the strong convergence of the gradient in the variational problems with critical exponents even on arbitrary domains. Notice that the concentration-compactness principle due to Lions [11] and the concentration-compactness principle at infinity by Bianchi et al. [1] are widely used to overcome the difficulty due to the lack of compactness. In the case of bounded domains and compact manifolds, the Struwe decomposition is also very useful to recover compactness in nonlinear elliptic problems with critical exponent [17, 6].

For the reader's convenience and to have a self-contained paper, we rephrase, in a more general framework, compactness results established in [12, 13, 15] and *reproduce* the proofs given therein (for instance), since they are stated for bounded sets with usual Leray-Lions operators, and for T-sets.

As examples of applications of our result, we recover in particular recent results in [7] on quasilinear elliptic equations involving critical sobolev exponents. There is a large literature on critical exponents see for instance [19, 20, 8, 7] and references therein. Two others examples are

given, but our principal result can be applied to a large class of quasilinear elliptic problems where there holds a lack of compactness.

As a corollary of our result, we can state

Lemma 1 *Let \hat{a} be a Caratheodory function from $\mathbb{R}^N \times \mathbb{R} \times \mathbb{R}^N$ into \mathbb{R}^N satisfying the usual Leray-Lions growth and monotonicity conditions. Let (u_n) be a bounded sequence of $W_{\text{loc}}^{1,p}(\mathbb{R}^N) = \{v \in L_{\text{loc}}^p(\mathbb{R}^N), |\nabla v| \in L_{\text{loc}}^p(\mathbb{R}^N)\}$, with $1 < p < +\infty$, (f_n) be a bounded sequence of $L_{\text{loc}}^1(\mathbb{R}^N)$ and (g_n) be a sequence of $W_{\text{loc}}^{-1,p'}(\mathbb{R}^N)$ tending strongly to zero. Assume that (u_n) satisfies:*

$$(H1) \quad \int_{\mathbb{R}^N} \hat{a}(x, u_n(x), \nabla u_n(x)) \cdot \nabla \varphi \, dx = \int_{\mathbb{R}^N} f_n \varphi \, dx + \langle g_n, \varphi \rangle,$$

$$\forall \varphi \in W_{\text{comp}}^{1,p}(\mathbb{R}^N) = \left\{ v \in W^{1,p}(\mathbb{R}^N), \text{ with compact support} \right\}, \varphi \text{ bounded.}$$

Then

1. there exists a function u such that $u_n(x) \rightarrow u(x)$ a.e. in \mathbb{R}^N ,
2. $u \in W_{\text{loc}}^{1,p}(\mathbb{R}^N)$,
3. there exists a subsequence, still denoted (u_n) , such that

$$\nabla u_n(x) \rightarrow \nabla u(x) \text{ a.e. in } \mathbb{R}^N.$$

This lemma 1 generalizes in particular some results of Boccardo-Murat [18]. Thus, we give an alternate proof for critical exponents equations instead of the concentration-compactness principle of P.L. Lions [11] and the concentration-compactness at infinity of Bianchi and al. [1] for such problems.

The above lemma will be used to show that suitable Palais-Smale sequences are in fact relatively compact. Let us recall the following

Définition 1 (Palais Smale sequence)

Let X be a real Banach space and $I : X \rightarrow \mathbb{R}$ a functional mapping which is Gâteaux differentiable. We say that a sequence $(u_n)_{n \geq 0}$ is a Palais-Smale sequence if:

- (a) there is a real number c such that $\lim_{n \rightarrow +\infty} I(u_n) = c$.
- (b) $\lim_{n \rightarrow +\infty} I'(u_n) = 0$ strongly in the dual X' .

Remark 1 *In our application X will be a reflexive Banach space, the conditions (a) and (b) will (often) imply the boundedness of (u_n) , from which we will have statement 1. and 2. (with additional compactness). The hypothesis (H1) will often follow from (b).*

2 Notations and compactness result.

Let Ω be an arbitrary open set of \mathbb{R}^N , we shall denote by $\omega \subset\subset \Omega$ any relatively compact open subset ω of Ω (that is $\bar{\omega} \subset \Omega$, where $\bar{\omega}$ is the closure of ω).

Let $1 < p_i < +\infty$, $i = 1, \dots, N$, we set

$$p = \min\{p_i, 1 \leq i \leq N\},$$

$$\mathbb{L}_{\text{loc}}^{\vec{p}}(\Omega) = \prod_{i=1}^N L_{\text{loc}}^{p_i}(\Omega), \quad \vec{p} = (p_1, \dots, p_N),$$

$$W_{\text{loc}}^{1, \vec{p}}(\Omega) = \left\{ v \in \bigcup_{i=1}^N L_{\text{loc}}^{p_i}(\Omega) : \nabla v \in \mathbb{L}_{\text{loc}}^{\vec{p}}(\Omega) \right\}.$$

We remark that $\bigcup_{i=1}^N L_{\text{loc}}^{p_i}(\Omega) = L_{\text{loc}}^p(\Omega)$.

For given $q \in (1, +\infty)$, we denote by $q' := \frac{q}{q-1}$ its conjugate exponent. We shall use the following globally real Lipschitz functions:

$$\text{For } \varepsilon > 0, \sigma \in \mathbb{R}, S_\varepsilon(\sigma) = \begin{cases} \sigma & \text{if } |\sigma| \leq \varepsilon \\ \varepsilon \text{sign}(\sigma) & \text{otherwise,} \end{cases}$$

and $\sigma^k := S_k(\sigma)$ for $k \geq 1$.

We shall consider a nonlinear map $\hat{a} : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ satisfying:

(L1) $\hat{a}(x, \cdot, \cdot)$ is a continuous map for almost every x and for all $(\sigma, \xi) \in \mathbb{R} \times \mathbb{R}^N$, $\hat{a}(\cdot, \sigma, \xi)$ is measurable (such a property is called Caratheodory property),

(L2) \hat{a} maps bounded sets of $W_{\text{loc}}^{1, \vec{p}}(\Omega)$ into bounded sets of $\prod_{i=1}^N L_{\text{loc}}^{p'_i}(\Omega)$,

and for almost all $x \in \Omega$, for all (σ, ξ) in $\mathbb{R} \times \mathbb{R}^N$, $\hat{a}(x, \sigma, \xi) \cdot \xi \geq 0$,

for almost every $x \in \Omega$ and for all $v \in W_{\text{loc}}^{1, \vec{p}}(\Omega)$, the mapping $u \mapsto \hat{a}(x, u, \nabla v)$ is continuous

from $W^{1, \vec{p}}(\omega)$ -weak into $\prod_{i=1}^N L^{p'_i}(\omega)$ -strong, for all $\omega \subset\subset \Omega$,

(L3) for almost every $x \in \Omega$, for all $(\sigma, \xi_i) \in \mathbb{R} \times \mathbb{R}^N$, $i = 1, 2$,

$$[\hat{a}(x, \sigma, \xi_1) - \hat{a}(x, \sigma, \xi_2)] [\xi_1 - \xi_2] > 0, \text{ for } \xi_1 \neq \xi_2.$$

(L4) if for some $x \in \Omega$, there is a sequence $(\sigma_n, \xi_{1n}) \in \mathbb{R} \times \mathbb{R}^N$, $\xi_2 \in \mathbb{R}^N$ such that $[\hat{a}(x, \sigma_n, \xi_{1n}) - \hat{a}(x, \sigma_n, \xi_2)] [\xi_{1n} - \xi_2]$ and σ_n are bounded as $n \rightarrow +\infty$ then $|\xi_{1n}|$ remains in a bounded set of \mathbb{R} as $n \rightarrow +\infty$.

We start with this result concerning the convergence almost everywhere of the gradients:

Theorem 1

Let (u_n) be a bounded sequence of $W_{\text{loc}}^{1, \vec{p}}(\Omega)$. Then

(i) There is a subsequence still denoted (u_n) and a function $u \in W_{\text{loc}}^{1, \vec{p}}(\Omega)$ such that $u_n(x) \rightarrow u(x)$ a.e. in Ω as $n \rightarrow +\infty$.

(ii) If furthermore, we assume (L1)-(L4) and that $\forall \varphi \in C_c^\infty(\Omega)$, $\forall k \geq k_0 > 0$:

$$\limsup_{n \rightarrow +\infty} \int_{\Omega} \widehat{a}(x, u_n(x), \nabla u_n(x)) \cdot \nabla(\varphi S_\varepsilon(u_n - u^k)) \leq o(1)$$

as $\varepsilon \rightarrow 0$ then there exists a subsequence still denoted (u_n) such that

$$\nabla u_n(x) \rightarrow \nabla u(x) \text{ a.e. in } \Omega.$$

Remark 2

2.1. The function $o(1)$ in (ii) might depend on k and φ .

2.2. As in [15], one can give a more general framework using T -sets instead of $W_{\text{loc}}^{1, p}(\Omega)$ but in view of our applications to variational problems derived from Euler equations, this framework seems not be appropriate.

2.3. The proof of (ii) is the same as in [13, 14, 15], for convenience we reproduce it here.

2.4. The assumption (L2) is satisfied if $p_1 = \dots = p_N = p$ and for all $\omega \subset\subset \Omega$, there is a constant $c_\omega > 0$ and a function $a_0 \in L^{p'}(\omega)$ such that for almost every $x \in \omega$, for all $(\sigma, \xi) \in \mathbb{R} \times \mathbb{R}^N$:

$$|\widehat{a}(x, \sigma, \xi)| \leq c_\omega \left[|\sigma|^{p-1} + |\xi|^{p-1} + a_0(x) \right].$$

and (L4) is true if $\widehat{a}(x, \sigma, \xi) \cdot \xi \geq c_\omega^1 |\xi|^p - c_\omega^2$, $c_\omega^1 > 0$.

2.5. Bounded sets in $W_{\text{loc}}^{1, \vec{p}}(\Omega)$ will be bounded in

$$W^{1, \vec{p}}(\omega) = \left\{ v \in \bigcup_{i=1}^N L^{p_i}(\omega), \nabla v \in \prod_{j=1}^N L^{p_j}(\omega) \right\}, \text{ for every } \omega \subset\subset \Omega.$$

2.6. If $p_1 = \dots = p_N$, we adopt the usual notation for Sobolev spaces: $W^{1, p}(\omega)$ and we use 2.4.

Proof.

(i) Let $(\omega_j)_{j \geq 0}$ be a sequence of bounded relatively compact subsets of Ω such that $\overline{\omega}_j \subset \omega_{j+1}$ and $\bigcup_{j=0}^{+\infty} \omega_j = \Omega$. Since $(u_n)_{n \geq 0}$ is bounded in $W^{1, \vec{p}}(\omega_j)$, by the usual embeddings, we deduce that there is a subsequence $u_{n(j)}$ and a function u in $W^{1, \vec{p}}(\omega_j)$ such that $u_{n(j)}(x) \xrightarrow{n \rightarrow +\infty} u(x)$. We conclude with the usual diagonal Cantor process.

(ii) Let $\varphi \in C_c^\infty(\Omega)$, $0 \leq \varphi \leq 1$, $\varphi = 1$ on ω_j and $\text{supp}(\varphi) \subset \omega_{j+1}$, and set

$$\Delta(u_n, u)(x) = \left[\widehat{a}(x, u_n(x), \nabla u_n(x)) - \widehat{a}(x, u_n(x), \nabla u(x)) \right] \nabla(u_n - u)(x).$$

Then one has:

(ii.1) $\Delta(u_n, u)(x) \geq 0$ a.e. on Ω (due to (L3)),

(ii.2) $\sup_n \int_{\omega_{j+1}} \Delta(u_n, u) dx$ is finite (since (u_n) is in a bounded set of $W_{\text{loc}}^{1, \overline{p}}(\Omega)$ and the growth condition (L2)).

Let us show that $\lim_n \int_{\Omega} \varphi \Delta(u_n, u)^{\frac{1}{p}} dx = 0$. On one hand, we have

$$\int_{\Omega} \varphi \Delta(u_n, u)^{\frac{1}{p}} dx = \int_{\{|u|>k\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx + \int_{\{|u|\leq k\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx. \quad (1)$$

By the Hölder inequality and noticing that

$$\text{meas}\{x \in \omega_{j+1} : |u| > k\} \leq \frac{c(j)}{k^p},$$

one deduces that

$$\int_{\{|u|>k\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx \leq \frac{c_1(j)}{k^{p-1}}. \quad (2)$$

$(c_m(j))$ are different constants depending on j and φ but independent of n , ε and k). While for the second integral, we have

$$\int_{\{|u|\leq k\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx = \int_{\{|u|\leq k\} \cap \{|u_n - u|\leq \varepsilon\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx + \int_{\{|u|\leq k\} \cap \{|u_n - u|>\varepsilon\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx. \quad (3)$$

Moreover, the second term in the right hand side of the last inequality satisfies

$$\int_{\{|u|\leq k\} \cap \{|u_n - u|>\varepsilon\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx \leq c_2(j) \text{meas}\{x \in \omega_{j+1} : |u_n - u|(x) > \varepsilon\}^{1 - \frac{1}{p}}$$

and since (u_n) converges to u in measure, we deduce that, for n sufficiently large, $\text{meas}\{x \in \omega_{j+1} : |u_n - u|(x) > \varepsilon\} \leq \varepsilon$. It follows that

$$\limsup_{n \rightarrow +\infty} \int_{\{|u|\leq k\} \cap \{|u_n - u|>\varepsilon\}} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx \leq c_2(j) \varepsilon^{1 - \frac{1}{p}}. \quad (4)$$

Setting $A_{n,k}^\varepsilon = \omega_{j+1} \cap \{|u| \leq k\} \cap \{|u_n - u| \leq \varepsilon\}$, we obtain from the Hölder inequality:

$$\int_{A_{n,k}^\varepsilon} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx \leq c_3(j) \left(I_{n,k}^1(\varepsilon) - I_{n,k}^2(\varepsilon) \right)^{\frac{1}{p}}, \quad (5)$$

with

$$I_{n,k}^1(\varepsilon) = \int_{A_{n,k}^\varepsilon} \widehat{a}(x, u_n, \nabla u_n) \cdot \nabla(u_n - u) \varphi dx,$$

$$I_{n,k}^2(\varepsilon) = \int_{\{|u|\leq k\}} \widehat{a}(x, u_n, \nabla u) \cdot \nabla S_\varepsilon(u_n - u) \varphi dx.$$

Since $\widehat{a}(x, u_n, \nabla u) \rightarrow \widehat{a}(x, u, \nabla u)$ strongly in $\prod_{i=1}^N L^{p'_i}(\omega_{j+1})$ (by the last statement of (L2)) and $\nabla S_\varepsilon(u_n - u) \rightarrow 0$ in $\prod_{i=1}^N L^{p_i}(\omega_{j+1})$ -weak, we deduce that

$$\lim_{n \rightarrow +\infty} I_{n,k}^2(\varepsilon) = 0, \quad (6)$$

while for the term $I_{n,k}^1(\varepsilon)$, we get:

$$\begin{aligned} I_{n,k}^1(\varepsilon) &\leq \int_{\Omega} \widehat{a}(x, u_n, \nabla u_n) \cdot \nabla(\varphi S_\varepsilon(u_n - u^k)) \\ &\quad - \int_{\Omega} \widehat{a}(x, u_n, \nabla u_n) \cdot \nabla \varphi S_\varepsilon(u_n - u^k) dx. \end{aligned} \quad (7)$$

Since

$$\left| \int_{\Omega} \widehat{a}(x, u_n, \nabla u_n) \cdot \nabla \varphi S_\varepsilon(u_n - u^k) dx \right| \leq c_4(j)\varepsilon, \quad (8)$$

the assumption (ii) implies then

$$\limsup_{n \rightarrow +\infty} I_{n,k}^1(\varepsilon) \leq c_4(j)\varepsilon + o(1) \text{ as } \varepsilon \rightarrow 0. \quad (9)$$

Combining relations (5), (6) and (9), it follows:

$$\limsup_{n \rightarrow +\infty} \int_{A_{n,k}^\varepsilon} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx \leq o(1) \text{ as } \varepsilon \rightarrow 0. \quad (10)$$

From relation (1), (2), (3), (4), and (10), we deduce:

$$\limsup_{n \rightarrow +\infty} \int_{\Omega} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx \leq o(1) \text{ (as } \varepsilon \rightarrow 0) + O\left(\frac{1}{k^{p-1}}\right). \quad (11)$$

Letting first $\varepsilon \rightarrow 0$ and then k to infinity, we then obtain:

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \Delta(u_n, u)^{\frac{1}{p}} \varphi dx = 0.$$

From which we deduce that for a subsequence $(u_{j_n})_{n \geq 0}$,

$$\Delta(u_{j_n}, u)(x) \rightarrow 0 \text{ a.e. on } \omega_j,$$

i.e. there is a subset Λ_j of ω_j such that $meas(\Lambda_j) = 0$ and

$$\Delta(u_{j_n}, u)(x) \rightarrow 0 \quad \forall x \in \omega_j \setminus \Lambda_j.$$

Then, due to the hypothesis (L4), the sequence $(|\nabla u_{j_n}|)$ is bounded in \mathbb{R} . We claim that

$$\forall x \in \omega_j \setminus \Lambda_j, \quad \nabla u_{j_n}(x) \rightarrow \nabla u(x).$$

Suppose that the last claim is false [10, 9], then there are $x \in \omega_j \setminus \Lambda_j$, $z^*(x) \neq \nabla u(x)$ and a subsequence $(|\nabla u_{j'_n}|)$ verifying

$$\nabla u_{j'_n}(x) \rightarrow z^*(x).$$

Since \widehat{a} is Caratheodory, the continuity implies that

$$\begin{cases} \left[\widehat{a}(x, u(x), z^*(x)) - \widehat{a}(x, u(x), \nabla u(x)) \right] \cdot (z^*(x) - \nabla u(x)) = 0 \\ z^*(x) \neq \nabla u(x), \end{cases}$$

which is in contradiction with the hypothesis (L3). The proof is achieved by the diagonal process of Cantor. \diamond

Proof of Lemma 1 (Corollary of theorem 1)

Here, we have $p_1 = \dots = p_N = p$. Since (u_n) belongs to a bounded set of $W_{\text{loc}}^{1,p}(\mathbb{R}^N)$, statement (i) of Theorem 1 implies that there is a function u and a subsequence still denoted by (u_n) such that

$$u_n(x) \xrightarrow{n \rightarrow +\infty} u(x) \text{ a.e. in } \mathbb{R}^N,$$

and

$$u \in W_{\text{loc}}^{1,p}(\mathbb{R}^N).$$

Then for all $\varphi \in C_c^\infty(\mathbb{R}^N)$, $\varphi S_\varepsilon(u_n - u^k)$ is an element of $W_{\text{comp}}^{1,p}(\mathbb{R}^N)$ and

$$\left| \int_{\mathbb{R}^N} f_n \varphi S_\varepsilon(u_n - u^k) dx \right| \leq \varepsilon |\varphi|_\infty |f_n|_{L^1(\omega)} \leq c(\varphi)\varepsilon, \quad (12)$$

(for every φ such that $\text{supp}(\varphi) \subset \omega$, $\bar{\omega}$ is a compact of \mathbb{R}^N), and

$$|\langle g_n, \varphi S_\varepsilon(u_n - u^k) \rangle| \leq |g_n|_{W^{-1,p'}(\omega)} |\varphi S_\varepsilon(u_n - u^k)|_{W^{1,p}(\mathbb{R}^N)}.$$

Using the fact that $|\varphi S_\varepsilon(u_n - u^k)|_{W^{1,p}(\mathbb{R}^N)}$ is bounded independently of ε, n, k and that $|g_n|_{W^{-1,p'}(\omega)} \xrightarrow{n \rightarrow \infty} 0$, it holds:

$$\limsup_n \int_{\mathbb{R}^N} \widehat{a}(x, u_n, \nabla u_n) \cdot \nabla(\varphi S_\varepsilon(u_n - u^k)) dx \leq O(\varepsilon).$$

Finally, Theorem 1 ends the proof. \diamond

Remark 3 *Many extensions of the Theorem 1 can be made (for instance on manifolds or on measure spaces). Here, we choose the above framework for the applications we made here. Nevertheless, one can use Theorem 1 for weighted spaces choosing correctly the open set Ω and the map \widehat{a} . Some results in that direction have been already made by Marchi [16], and also by Fengquan Li, Zhao Huixiu [5] using the method of [12, 13].*

3 Some examples of applications.

3.1 Example 1.

We start by recovering a recent result of H. Ohya [7] using this alternate proof (without concentration-compactness principles) to show that we simplify the author's proof.

For this, we recall a part of the author's framework.

Let $1 < p < N$, $p^* = \frac{Np}{N-p}$ and let Ω be an unbounded open set with smooth boundary if $\partial\Omega \neq \emptyset$, $\theta(x)$, $a(x)$, $K(x)$ be three non negative functions with the additional regularity that $\theta \in C^2(\Omega)$, $a \in L^r(\Omega)$ for some $r \in \left[\frac{N}{p}, +\infty\right]$ and K is such that $e^{(p-p^*)\theta(x)}K(x) = V(x)$, being bounded.

The author defined the following sets and quantities

$$\begin{aligned} L^p(\theta, \Omega) &= \left\{ u \in L^p(\Omega); \int_{\Omega} e^{p\theta(x)} |u|^p dx < +\infty \right\}, \\ W^{1,p}(\theta, \Omega) &= \left\{ u \in W_0^{1,p}(\Omega) : \int_{\Omega} e^{p\theta} (|u|^p + |\nabla u|^p) < +\infty \right\}, \\ \lambda_1 &= \inf_{u \in W^{1,p}(\theta, \Omega) \setminus \{0\}} \left\{ \int_{\Omega} e^{p\theta(x)} |\nabla u|^p dx / \int_{\Omega} e^{p\theta(x)} a(x) |u|^p dx \right\} > 0, \\ I_{\theta}(u) &= \frac{1}{p} \int_{\Omega} e^{p\theta(x)} (|\nabla u|^p - \lambda a(x) |u|^p) dx - \frac{1}{p^*} \int_{\Omega} e^{p\theta(x)} K(x) |u|^{p^*} dx, \end{aligned}$$

for $u \in W^{1,p}(\theta, \Omega)$. The author showed, under some hypotheses on θ and Ω , that $W^{1,p}(\theta, \Omega)$ is embedded continuously (resp. compactly) in $L^q(\theta, \Omega)$ provided that $q \in [p, p^*]$ (resp. $q \in [p, p^*)$). Moreover, in this situation the Poincaré inequality holds true, which implies that the semi-norm $\left(\int_{\Omega} e^{p\theta} |\nabla u|^p dx \right)^{\frac{1}{p}}$ is in fact a norm on $W^{1,p}(\theta, \Omega)$. We shall prove the following Palais Smale property:

Theorem 2

Under the above property, for every $\lambda < \lambda_1$, any Palais-Smale sequence $(u_m)_m$ of I_{θ} on $X = W^{1,p}(\theta, \Omega)$ satisfying :

- (a) $I_{\theta}(u_m) \rightarrow b_{\theta}$
- (b) $I'_{\theta}(u_m) \rightarrow 0$ in $(W^{1,p}(\theta, \Omega))'$ (dual of X)

contains a convergent subsequence in $W^{1,p}(\theta, \Omega)$ provided that

$$0 < b_{\theta} < b_{\theta}^* = \frac{1}{N} S^{\frac{N}{p}} |V|_{\infty}^{-\frac{N-p}{p}},$$

where

$$S = \inf_{u \in W_0^{1,p}(\Omega) \setminus \{0\}} \left\{ \int_{\Omega} |\nabla u|^p dx / \left(\int_{\Omega} |u|^{p^*} dx \right)^{\frac{p}{p^*}} \right\}.$$

Remark 4 *Theorem 2 corresponds to Theorem 4.1 in [7].*

Proof of theorem 2

The sequence (u_m) is bounded in X as it was observed in [7]. On one hand one has

$$I_{\theta}(u_m) - \frac{1}{p^*} \langle I'_{\theta}(u_m), u_m \rangle = \frac{1}{N} \|u_m\|_X^p - \frac{\lambda}{N} \int_{\Omega} a(x) |u_m|^p e^{p\theta} dx$$

and applying the Poincaré-Sobolev inequality, there is a positive constant c such that

$$\frac{1}{N} \left(1 - \frac{\lambda}{\lambda_1}\right) \|u_m\|_X^p \leq I_\theta(u_m) - \frac{1}{p^*} \langle I'_\theta(u_m), u_m \rangle \leq c.$$

Thus, $\int_\Omega e^{p\theta(x)} K(x) |u_m|^{p^*} dx$ and $\int_\Omega e^{p\theta(x)} a(x) |u_m|^p dx$ are bounded independently of m . If we set

$$\widehat{a}(x, \xi) = e^{p\theta(x)} |\xi|^{p-2} \xi, \quad x \in \Omega, \quad \xi \in \mathbb{R}^N,$$

then \widehat{a} satisfies conditions (L1)-(L4) (since $\theta \in C(\Omega)$). With the definition of X , we may then apply the first statement of Theorem 1 to conclude that there is an element $u \in X$ and subsequence still denoted (u_m) such that

$$u_m(x) \rightarrow u(x) \text{ a.e. in } \Omega.$$

Furthermore since $S_\varepsilon(u_m - u^k) \in X$ and $\|S_\varepsilon(u_m - u^k)\|_X \leq \|u_m\|_X + \|u\|_X$, we get for all $\varphi \in C_c^\infty(\Omega)$:

$$\begin{aligned} & \int_\Omega \widehat{a}(x, \nabla u_m) \cdot \nabla (\varphi S_\varepsilon(u_m - u^k)) dx \\ & \leq \lambda \varepsilon \left(\int_\Omega e^{p\theta(x)} |u_m|^p a(x) dx \right)^{\frac{1}{p'}} \left(\int_\Omega \varphi^p e^{p\theta(x)} a(x) dx \right)^{\frac{1}{p}} \\ & \quad + \varepsilon \left(\int_\Omega e^{p\theta(x)} |u_m|^{p^*} K(x) dx \right)^{\frac{1}{p^{*'}}} \left(\int_\Omega \varphi^{p^*} e^{p\theta(x)} K(x) dx \right)^{\frac{1}{p^*}} \\ & \quad + c \|I'_\theta(u_m)\|_{X'} (\|u_m\|_X + \|u\|_X). \end{aligned}$$

Thus,

$$\limsup_{m \rightarrow +\infty} \int_\Omega \widehat{a}(x, \nabla u_m) \cdot \nabla (\varphi S_\varepsilon(u_m - u^k)) dx \leq O(\varepsilon).$$

Thus, from Theorem 1, passing if necessary to a subsequence, $\nabla u_m(x) \xrightarrow{m \rightarrow +\infty} \nabla u(x)$ a.e. in Ω .
>From Vitali's theorem, we then deduce that u is a critical point of I_θ , that is,

$$\forall \phi \in X, \quad \langle I'_\theta(u), \phi \rangle = 0.$$

At this stage, we apply the Brézis-Lieb lemma [2], that is from the equations,

$$\langle I'_\theta(u), u \rangle = 0, \quad \langle I'_\theta(u_m), u_m \rangle = o(1),$$

we then have

$$\lim_{m \rightarrow +\infty} \int_\Omega e^{p\theta(x)} |\nabla(u - u_m)|^p dx = \lim_{m \rightarrow +\infty} \int_\Omega e^{p\theta(x)} |u_m - u|^{p^*}(x) K(x) dx =: \ell.$$

(Note that from the integrability of a and the boundedness of (u_m) in X , we deduce that $\lim_{m \rightarrow +\infty} \int_\Omega a(x) |u_m - u|^p e^{p\theta(x)} dx = 0$ (see Proposition 1 below).)

Since $V(x) = e^{(p-p^*)\theta(x)} K(x)$ is bounded, we deduce

$$\ell \leq |V|_\infty \int_\Omega e^{p^*\theta(x)} |u_m - u|^{p^*}(x) dx + o(1).$$

From the definition of S , one obtains $\ell \leq |V|_\infty S^{\frac{p^*}{p}} \ell^{\frac{p^*}{p}}$.

If $\ell = 0$, the proof is done.

If $\ell \neq 0$ then $\ell \geq S^{\frac{N}{p}} |V|_\infty^{-\frac{N-p}{p}}$. Since, $\lim_m I_\theta(u_m) = b_\theta$, thus
 $b_\theta = \frac{1}{N} \int_\Omega |u|^{p^*} K(x) e^{p\theta(x)} dx + \frac{1}{N} \ell + o(1)$ (still using the Brézis-Lieb's lemma).
Then $b_\theta \geq \frac{1}{N} \ell \geq \frac{1}{N} S^{\frac{N}{p}} |V|_\infty^{-\frac{N-p}{p}} = b_\theta^*$, which gives a contradiction. Thus necessarily $\ell = 0$. \diamond

In the next example, we give a classical existence result with a new proof based on our approach.

3.2 Example 2.

In this paragraph, we are concerned with the existence of (at least) one positive solution to the elliptic problem

$$-\Delta_p u = \lambda a(x) |u|^{p-2} u + |u|^{p^*-2} u \text{ in } \mathbb{R}^N, \quad (13)$$

where the function a satisfies the following conditions: $a \geq 0$ on \mathbb{R}^N , $a \not\equiv 0$ and $a \in L^{\frac{N}{p}}(\mathbb{R}^N)$. The parameter λ is assumed to be positive, $1 < p < N$ and $N \geq 3$.

Consider the Euler-Lagrange functional associated to Problem (13) defined by

$$J_\lambda(u) := \frac{1}{p} \int_{\mathbb{R}^N} |\nabla u|^p dx - \frac{\lambda}{p} \int_{\mathbb{R}^N} a(x) |u|^p dx - \frac{1}{p^*} \int_{\mathbb{R}^N} |u|^{p^*} dx$$

which is of class $C^1(\mathcal{D}^{1,p}(\mathbb{R}^N))$. We recall that $\mathcal{D}^{1,p}(\mathbb{R}^N)$ is the completion of the space $\mathcal{D}(\mathbb{R}^N)$ with respect to the norm

$$\|\varphi\| := \left(\int_{\mathbb{R}^N} |\nabla \varphi|^p dx \right)^{\frac{1}{p}}.$$

The space $\mathcal{D}^{1,p}(\mathbb{R}^N)$ can also be seen as

$$\mathcal{D}^{1,p}(\mathbb{R}^N) = \left\{ \varphi \in L^{p^*}(\mathbb{R}^N) : |\nabla \varphi| \in L^p(\mathbb{R}^N) \right\}.$$

By solutions of Problem (13) we understand critical points of the functional J_λ . Remark that the functional J_λ is bounded neither above nor below on $\mathcal{D}^{1,p}(\mathbb{R}^N)$. Then, to find possible critical points of J_λ , we limit the study to the corresponding Nehari manifold which contains all critical points of J_λ . We recall that the Nehari manifold associated to J_λ , denoted by \mathcal{N}_{J_λ} , is defined by

$$\mathcal{N}_{J_\lambda} := \{ \varphi \in \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\} : J'_\lambda(\varphi)(\varphi) = 0 \}.$$

In the sequel, we will set $\|u\|_{a,p} := \left(\int_{\mathbb{R}^N} a(x) |u|^p dx \right)^{\frac{1}{p}}$.

Lemma 2 *For every $\lambda > 0$, the functional J_λ is bounded below on the Nehari manifold \mathcal{N}_{J_λ} .*

Proof. For every $u \in \mathcal{N}_{J_\lambda}$, it holds

$$J_\lambda(u) = \left(\frac{1}{p} - \frac{1}{p^*} \right) \|u\|_{p^*}^{p^*} = \frac{1}{N} \|u\|_{p^*}^{p^*},$$

and this ends the proof. \diamond

As before, due to the integrability of a , the compact embedding $D^{1,p}(\mathbb{R}^N) \subset L_{\text{loc}}^p(\mathbb{R}^N)$, the Vitali's theorem implies the following statement:

Lemma 3 *The functional*

$$\begin{aligned} \mathcal{D}^{1,p}(\mathbb{R}^N) &\longrightarrow \mathbb{R} \\ u &\longmapsto \int_{\mathbb{R}^N} a(x) |u|^p dx \end{aligned}$$

is weakly continuous.

We recall that the Nehari manifold can be characterized more explicitly by

$$\mathcal{N}_{J_\lambda} := \left\{ t\varphi; (t, \varphi) \in (\mathbb{R} \setminus \{0\}) \times (\mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\}) : \frac{d}{dt} J_\lambda(t\varphi) = 0 \right\}.$$

For this reason, we introduce the modified functional

$$\tilde{J}_\lambda(t, u) := J_\lambda(tu), \text{ on } \mathbb{R} \times \mathcal{D}^{1,p}(\mathbb{R}^N).$$

Since we are interested in positive solutions to Problem (13), we restrict ourselves in what follows to $t > 0$.

A direct computation shows that for every $u \in \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\}$, there is a unique value $\lambda(u)$ of λ defined by

$$\lambda(u) = \frac{\|\nabla u\|_p^p}{\|u\|_{a,p}^p},$$

such that for every $\lambda \in (0, \lambda(u))$, one has $t(u, \lambda)u \in \mathcal{N}_{J_\lambda}$, where

$$t(u, \lambda) = \left(\frac{\|\nabla u\|_p^p - \lambda \|u\|_{a,p}^p}{\|u\|_{p^*}^{p^*}} \right)^{\frac{1}{p^*-p}}. \quad (14)$$

We introduce

$$\lambda_1 := \inf_{u \in \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\}} \lambda(u)$$

which is not other than the first eigenvalue to the nonlinear eigenvalue problem

$$-\Delta_p u = \lambda a(x) |u|^{p-2} u \text{ in } \mathbb{R}^N.$$

>From Lemma (3), we get clearly that the characteristic value λ_1 is positive. Now, for every $\lambda \in (0, \lambda_1)$ one has more precisely

$$\mathcal{N}_{J_\lambda} = \{t(u, \lambda)u : u \in \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\}\}.$$

For every $\lambda \in (0, \lambda_1)$, we introduce

$$\alpha(\lambda) := \inf_{u \in \mathcal{N}_{J_\lambda}} J_\lambda(u) = \inf_{u \in \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\}} J_\lambda(t(u, \lambda)u).$$

It is not difficult to see that the functional

$$\begin{aligned} \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\} &\longrightarrow \mathbb{R} \\ u &\longmapsto J_\lambda(t(u, \lambda)u) \end{aligned}$$

is 0-homogeneous. Then we get

$$\alpha(\lambda) = \inf_{u \in \mathbb{S}} J_\lambda(t(u, \lambda)u), \quad (15)$$

where \mathbb{S} is the unit sphere in $\mathcal{D}^{1,p}(\mathbb{R}^N)$. In this particular case, we have in fact

$$\alpha(\lambda) := \inf_{u \in \mathbb{S}} \left(\frac{1}{p} - \frac{1}{p^*} \right) \left(\frac{\|\nabla u\|_p^p - \lambda \|u\|_{a,p}^p}{\|u\|_{p^*}^p} \right)^{\frac{p^*}{p^* - p}}.$$

In what follows, we will write $(PS)_c$ to denote a Palais-Smale sequence of J_λ with the level $c \in \mathbb{R}$.

Lemma 4 *Let $\lambda \in (0, \lambda_1)$. There exists a minimizing sequence of (15) denoted by $(u_n) \subset \mathbb{S}$ such that:*

- (i) $0 < \liminf_{n \rightarrow \infty} t(u_n, \lambda) \leq \limsup_{n \rightarrow \infty} t(u_n, \lambda) < +\infty$.
- (ii) $(t(u_n, \lambda)u_n)$ is a bounded Palais-Smale sequence for J_λ .

Proof. Let us denote $U_n := t(u_n, \lambda)u_n$, it holds obviously that $t(u_n, \lambda) = \|\nabla U_n\|_p$. On one hand, one has

$$\left(\frac{1}{p} - \frac{1}{p^*} \right) \left(\|\nabla U_n\|_p^p - \lambda \|U_n\|_{a,p}^p \right) = \alpha(\lambda) + o_n(1).$$

Then,

$$\|\nabla U_n\|_p^p = \lambda \|U_n\|_{a,p}^p + N\alpha(\lambda) + o_n(1).$$

Applying the Hölder and Young inequalities to $\|U_n\|_{a,p}^p$, we conclude that $\|\nabla U_n\|_p$ is bounded.

On the other hand, for every $u \in \mathbb{S}$, one gets

$$J_\lambda(tu) \geq \frac{t^p}{p} \left(1 - \frac{\lambda}{\lambda_1} \right) - \frac{t^{p^*}}{p^* S^{\frac{p^*}{p}}},$$

where S is the best Sobolev constant in the embedding $\mathcal{D}^{1,p}(\mathbb{R}^N) \subset L^{p^*}(\mathbb{R}^N)$. It follows that

$$\exists \delta(\lambda) > 0 \text{ such that } \alpha(\lambda) > \delta(\lambda) > 0, \quad (16)$$

and consequently $\liminf_{n \rightarrow \infty} \|\nabla U_n\|_p > 0$. This ends the claim (i).

For every $u \in \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\}$ and $\lambda \in (0, \lambda_1)$, we have $\partial_t \tilde{J}_\lambda(t(u, \lambda), u) = 0$ and $\partial_{tt} \tilde{J}_\lambda(t(u, \lambda), u) < 0$. The implicit function theorem implies that $t(u, \lambda)$ is C^1 with respect to u since J_λ is. Let us introduce the C^1 functional \mathcal{J}_λ defined on \mathbb{S} by

$$\mathcal{J}_\lambda(u) = \tilde{J}_\lambda(t(u, \lambda), u) \equiv J_\lambda(t(u, \lambda)u).$$

Then

$$\alpha(\lambda) = \inf_{u \in \mathbb{S}} \mathcal{J}_\lambda(u).$$

Using the Ekeland variational principle on the complete manifold $(\mathbb{S}, \|\cdot\|)$ to the functional \mathcal{J}_λ , there exists a minimizing sequence of (15) denoted by $(u_n) \subset \mathbb{S}$ such that:

$$|\mathcal{J}'_\lambda(u_n)(\varphi_n)| \leq \frac{1}{n} \|\varphi_n\|, \text{ for every } \varphi_n \in \mathbb{T}_{u_n}\mathbb{S},$$

where $\mathbb{T}_{u_n}\mathbb{S}$ is the tangent space to \mathbb{S} at the point u_n . Moreover, for every $\varphi_n \in \mathbb{T}_{u_n}\mathbb{S}$, one has

$$\begin{aligned} \mathcal{J}'_\lambda(u_n)(\varphi_n) &= \partial_t \tilde{\mathcal{J}}_\lambda(t(u_n, \lambda), u_n) t'(u_n, \lambda)(\varphi_n) + \partial_u \tilde{\mathcal{J}}_\lambda(t(u_n, \lambda), u_n)(\varphi_n), \\ &= \partial_u \tilde{\mathcal{J}}_\lambda(t(u_n, \lambda), u_n)(\varphi_n), \end{aligned}$$

since $\partial_t \tilde{\mathcal{J}}_\lambda(t(u_n, \lambda), u_n) \equiv 0$, where $t'(u_n, \lambda)$ denotes the derivative of $t(\cdot, \lambda)$ with respect to its first variable at the point (u_n, λ) .

Furthermore, let

$$\begin{aligned} \pi : \mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\} &\longrightarrow \mathbb{R} \times \mathbb{S} \\ u &\longmapsto \left(\|u\|, \frac{u}{\|u\|} \right) := (\pi_1(u), \pi_2(u)). \end{aligned}$$

Applying Hölder's inequality, we get for every $(u, \varphi) \in (\mathcal{D}^{1,p}(\mathbb{R}^N) \setminus \{0\}) \times \mathcal{D}^{1,p}(\mathbb{R}^N)$:

$$\begin{cases} |\pi'_1(u)(\varphi)| &\leq \|\varphi\|, \\ \|\pi'_2(u)(\varphi)\| &\leq 2 \frac{\|\varphi\|}{\|u\|}. \end{cases}$$

>From (i), there is a positive constant C such that

$$t(u_n, \lambda) \geq C, \quad \forall n \in \mathbb{N}.$$

Then for every $\varphi \in \mathcal{D}^{1,p}(\mathbb{R}^N)$, there are $\varphi_n^1 \in \mathbb{R}$ and $\varphi_n^2 \in \mathbb{T}_{u_n}\mathbb{S}$ such that $|\varphi_n^1| \leq \|\varphi\|$, $\|\varphi_n^2\| \leq \frac{2}{C} \|\varphi\|$ and

$$\begin{aligned} J'_\lambda(t(u_n, \lambda)u_n)(\varphi) &= \partial_t \tilde{\mathcal{J}}_\lambda(t(u_n, \lambda), u_n)(\varphi_n^1) + \partial_u \tilde{\mathcal{J}}_\lambda(t(u_n, \lambda), u_n)(\varphi_n^2), \\ &= \partial_u \tilde{\mathcal{J}}_\lambda(t(u_n, \lambda), u_n)(\varphi_n^2), \\ &= \mathcal{J}'_\lambda(u_n)(\varphi_n^2). \end{aligned}$$

Therefore,

$$\begin{aligned} J'_\lambda(t(u_n, \lambda)u_n)(\varphi) &\leq \frac{1}{n} \|\varphi_n^2\| \\ &\leq \frac{2}{nC} \|\varphi\|. \end{aligned}$$

We easily conclude that

$$\lim_{n \rightarrow \infty} J'_\lambda(U_n) = 0 \text{ in } \mathcal{D}^{-1,p'}(\mathbb{R}^N),$$

which achieves the proof. ◇

As a consequence of Theorem 4 (see Example 3 with $\rho = \rho_1 = 1 = K$, $p_c = p^* = \frac{Np}{N-p}$), one has :

Lemma 5 Let $\lambda \in (0, \lambda_1)$ and $(U_n) \subset \mathcal{D}^{1,p}(\mathbb{R}^N)$ be a Palais-Smale sequence for J_λ such that $U_n \rightharpoonup U$ in $\mathcal{D}^{1,p}(\mathbb{R}^N)$. Then, passing if necessary to a subsequence, we get

$$\nabla U_n \longrightarrow \nabla U \text{ a.e. in } \mathbb{R}^N. \quad (17)$$

Lemma 6 Let $\lambda \in (0, \lambda_1)$ and $(U_n) \subset \mathcal{D}^{1,p}(\mathbb{R}^N)$ be a $(PS)_c$ sequence for J_λ . If

$$0 < c < \frac{1}{N} S^{\frac{N}{p}} \quad (18)$$

then Problem (13) has a nontrivial solution.

Proof. >From theorem 4, J_λ satisfies the Palais-Smale conditions if $0 < c < \frac{1}{N} S^{\frac{N}{p}}$. Thus, any weak limit U of U_n in $\mathcal{D}^{1,p}(\mathbb{R}^N)$ satisfies then $J'_\lambda(U) = 0$, $J_\lambda(U) = c > 0$. Consequently, U is nontrivial critical point of J_λ . \diamond

Lemma 7 For every $\lambda \in (0, \lambda_1)$ we have $0 < \alpha(\lambda) < \frac{1}{N} S^{\frac{N}{p}}$.

Proof. Let

$$\Psi_\varepsilon(x) := \frac{\left(N\varepsilon^{\frac{N-p}{p-1}}\right)^{\frac{N-p}{p^2}}}{\left(\varepsilon + |x|^{\frac{p}{p-1}}\right)^{\frac{N-p}{p}}}, \quad x \in \mathbb{R}^N, \quad \varepsilon > 0.$$

It is well known that

$$\|\nabla \Psi_\varepsilon\|_p^p = \|\Psi_\varepsilon\|_{p^*}^{p^*} = S^{N/p}.$$

Moreover,

$$J_\lambda(t(\Psi_\varepsilon), \lambda)\Psi_\varepsilon := \left(\frac{1}{p} - \frac{1}{p^*}\right) \left(\frac{\|\nabla \Psi_\varepsilon\|_p^p - \lambda \|\Psi_\varepsilon\|_{a,p}^p}{\|\Psi_\varepsilon\|_{p^*}^p}\right)^{\frac{p^*}{p^*-p}} < \frac{1}{N} S^{N/p}.$$

Then, using (16) we obtain

$$0 < \alpha(\lambda) < \frac{1}{N} S^{N/p}.$$

\diamond

Theorem 3 For every $\lambda \in (0, \lambda_1)$, Problem (13) admits at least one positive solution.

Proof. From the preceding lemmas, it is clear that Problem (13) possesses a solution u which is nontrivial, since $\alpha(\lambda) \neq 0$. Since J_λ is even in u , that $u \in \mathcal{D}^{1,p}(\mathbb{R}^N)$ implies that $|u| \in \mathcal{D}^{1,p}(\mathbb{R}^N)$ and $t(u, \lambda) = t(|u|, \lambda)$, we conclude that Problem (13) has a nontrivial nonnegative solution. The maximum principle achieves the proof. \diamond

3.3 Example 3.

Similar examples on weighted spaces can be made. Let ρ, ρ_1 be two nonnegative continuous functions on \mathbb{R}^N and assume that the closed set $F = \{x \in \mathbb{R}^N : \rho(x) = 0\} \cup \{x \in \mathbb{R}^N : \rho_1(x) = 0\}$

is of measure zero. We set $\Omega = \mathbb{R}^N \setminus F$.

For every $1 < p < +\infty$, we define

$$Y := D^{1,p}(\mathbb{R}^N, \rho) = \left\{ u \in L^p_{\text{loc}}(\Omega) : \int_{\mathbb{R}^N} |\nabla u|^p \rho(x) dx < +\infty \right\}.$$

Let $p_c > p$ so that

$$S_\rho = \inf_{Y \setminus \{0\}} \left\{ \int_{\mathbb{R}^N} |\nabla u|^p \rho(x) dx / \left(\int_{\mathbb{R}^N} |u|^{p_c} \rho_1(x) dx \right)^{\frac{p}{p_c}} \right\} > 0.$$

We set $\frac{1}{N_c} = \frac{1}{p} - \frac{1}{p_c}$ and let

$$a \in L^{\frac{N_c}{p}}_+(\mathbb{R}^N, \rho_1) = \left\{ f \geq 0 : \int_{\mathbb{R}^N} f(x)^{\frac{N_c}{p}} \rho_1(x) dx < +\infty \right\},$$

and the Euler-Lagrange functional defined on Y by

$$I_\rho(u) = \int_{\mathbb{R}^N} |\nabla u|^p \rho(x) dx - \frac{\lambda}{p} \int_{\mathbb{R}^N} |u|^p a(x) \rho_1(x) dx - \frac{1}{p_c} \int_{\mathbb{R}^N} K(x) |u|^{p_c} \rho_1(x) dx,$$

where K is a positive function in $L^\infty(\mathbb{R}^N)$. As previously, we have

$$\lambda_{1\rho} = \inf \left(\int_{\mathbb{R}^N} |\nabla u|^p \rho(x) / \int_{\mathbb{R}^N} |u|^p a(x) \rho_1(x) dx \right) > 0,$$

(see Proposition 1).

Theorem 4

Let $\lambda < \lambda_{1\rho}$, and (u_m) be a Palais-Smale sequence of I_ρ satisfying

- (a) $I_\rho(u_m) \rightarrow b_\rho$,
- (b) $I'_\rho(u_m) \rightarrow 0$ in Y' .

Then

1. (u_m) contains a weakly convergent subsequence to a function $u \in Y$ satisfying

$$I'_\rho(u) = 0.$$

2. This subsequence is strongly convergent in Y provided that

$$0 < b_\rho < b_\rho^* = \frac{1}{N_c} S_\rho^{\frac{N_c}{p}} \|K\|_\infty^{-\frac{N_c-p}{p}}.$$

We first need the following

Proposition 1

Under the above assumptions, if $(u_m)_m$ is weakly convergent to u in Y then

$$\lim_m \int_{\mathbb{R}^N} a(x) |u_m - u|^p \rho_1(x) dx = 0.$$

Proof

Since $\int_{\mathbb{R}^N} |u_m|^{p_c} \rho_1(x) dx$ is bounded uniformly with respect to m , then for every subset A of \mathbb{R}^N

$$\int_A a(x) |u_m - u|^p \rho_1(x) dx \leq c \left(\int_A a(x)^{\frac{N_c}{p}} \rho_1(x) dx \right)^{\frac{p}{N_c}} < +\infty.$$

But $Y \subset W_{\text{loc}}^{1,p}(\Omega)$ and then Y is compactly embedded in $L_{\text{loc}}^p(\Omega)$.

Thus using the Vitali's theorem one has, for every bounded set ω in \mathbb{R}^N ,

$$\lim_{m \rightarrow +\infty} \int_{\omega} a(x) |u_m - u|^p \rho_1(x) dx = 0.$$

(Recall that $\Omega = \mathbb{R}^N \setminus F$ and $\text{meas}(F) = 0$). Combining the two relations and the integrability of a , we conclude

$$\lim_m \int_{\mathbb{R}^N} a(x) |u_m - u|^p \rho_1(x) dx = 0.$$

◇

Notice that the previous proposition implies that $\lambda_{1\rho} > 0$.

The proof of Theorem 4 is similar to that of Theorem 2, we sketch it as the following:

Sketch of the proof of Theorem 4

Setting $\|u_m\|_Y^p = \int_{\mathbb{R}^N} |\nabla u_m|^p \rho(x) dx$, it follows that

$$\|u_m\|_Y^p \left(1 - \frac{\lambda}{\lambda_{1\rho}} \right) \frac{1}{N_c} \leq I_{\rho}(u_m) - \frac{1}{p_c} \langle I'_{\rho}(u_m), u_m \rangle.$$

Thus $\|u_m\|_Y$ remains in a bounded set of \mathbb{R} which implies that

$$\int_{\mathbb{R}^N} a(x) |u_m|^p \rho_1(x) dx \text{ and } \int_{\mathbb{R}^N} |u_m|^{p_c} K(x) \rho_1(x) dx$$

are in a bounded set of \mathbb{R} . Thus $\int_{\mathbb{R}^N} |u_m|^{p_c} \rho_1(x) dx$ is bounded. We deduce in particular that (u_m) remains in a bounded set of $W_{\text{loc}}^{1,p}(\Omega)$ (since for every $\omega \subset\subset \Omega$, $\rho_{\omega} = \inf_{\omega} \rho(x) > 0$ and $\rho_{1\omega} = \inf_{\omega} \rho_1(x) > 0$). We may appeal the first statement (i) of Theorem 1: There exist $u \in D^{1,p}(\mathbb{R}^N, \rho)$ and a subsequence still denoted by $(u_m)_m$ such that $u_m(x) \rightarrow u(x)$ a.e. in \mathbb{R}^N . Moreover since $S_{\varepsilon}(u_m - u^k) \in Y$, if we set $\hat{a}(x, \xi) = \rho(x) |\xi|^{p-2} \xi$, for all $x \in \Omega$ and $\xi \in \mathbb{R}^N$, then the conditions (L1)-(L4) are satisfied. Furthermore, $\forall \varphi \in C_c^{\infty}(\Omega)$, choosing $\omega \subset\subset \Omega$ such that $\text{supp } \varphi \subset \omega$, $\sup_{\omega} \rho(x)$ and $\sup_{\omega} \rho_1(x)$ are finite, it holds then:

$$\int_{\Omega} \hat{a}(x, \nabla u_m(x)) \cdot \nabla(\varphi S_{\varepsilon}(u_m - u^k)) dx \leq c(\varphi)\varepsilon + \|I'_{\rho}(u_m)\|_{Y'} (\|u_m\|_Y + \|u\|_Y).$$

Consequently, statement (ii) of Theorem 1 is also satisfied so, we conclude that (for a subsequence) $\nabla u_m(x) \rightarrow \nabla u(x)$ a.e. on \mathbb{R}^N . From Vitali's theorem, we deduce that

$$\forall \varphi \in D^{1,p}(\mathbb{R}^N) \langle I'_{\rho}(u), \varphi \rangle = 0.$$

To conclude that $\|u_m - u\|_Y \rightarrow 0$, we use the Brézis-Lieb lemma to get

$$\int_{\mathbb{R}^N} |\nabla u_m - \nabla u|^p \rho(x) dx = - \int_{\mathbb{R}^N} |\nabla u|^p \rho(x) dx + \int_{\mathbb{R}^N} |\nabla u_m|^p \rho(x) dx + o(1),$$

$$\begin{aligned} \int_{\mathbb{R}^N} |u_m - u|^{p_c} K(x) \rho_1(x) dx &= - \int_{\mathbb{R}^N} |u|^{p_c} K(x) \rho_1(x) dx \\ &\quad + \int_{\mathbb{R}^N} |\nabla u_m|^{p_c} K(x) \rho_1(x) dx + o(1) \end{aligned}$$

Since $\langle I'_\rho(u), u \rangle = 0$, $\langle I'_\rho(u_m), u_m \rangle = o(1)$, the above equalities imply that

$$\lim_{m \rightarrow \infty} \|u_m - u\|_Y^p = \lim_m \int_{\mathbb{R}^N} |u_m - u|^{p_c} K(x) \rho_1(x) dx =: \ell_0$$

(noticing that $\lim_m \int_{\mathbb{R}^N} a(x) |u_m - u|^p \rho_1(x) dx = 0$ (see Proposition 1)).

Since $K \in L^\infty(\mathbb{R}^N)$

$$\int_{\mathbb{R}^N} |u_m - u|^{p_c} K(x) \rho_1(x) dx \leq |K|_\infty S_p^{\frac{p_c}{p}} \left(\int_{\mathbb{R}^N} |\nabla(u_m - u)|^p \rho(x) dx \right)^{\frac{p_c}{p}}.$$

Thus

$$\ell_0 \leq |K|_\infty S_p^{\frac{p_c}{p}} \ell_0^{\frac{p_c}{p}}.$$

If $\ell_0 = 0$ the proof is done.

If $\ell_0 \neq 0$ then $\ell_0 \geq S_p^{\frac{N_c}{p}} |K|_\infty^{-\frac{N_c-p}{p}}$. Since

$$\lim_m I_\rho(u_m) = b_\rho : b_\rho \geq \frac{1}{N_c} \ell_0 \geq \frac{1}{N_c} S_p^{-\frac{N_c-p}{p}} |K|_\infty^{-\frac{N_c-p}{p}} = b_\rho^*,$$

which leads to a contradiction. \diamond

Theorem 5 *There exists $\lambda^* \in (0, \lambda_{1\rho})$ such that the problem*

$$-\operatorname{div}(\rho(x)|\nabla u|^{p-2}\nabla u) = \rho_1(x) (\lambda a(x)|u|^{p-2}u + K(x)|u|^{p_c-2}u) \quad \text{in } \mathbb{R}^N \quad (19)$$

has a nontrivial positive solution for every $\lambda \in (\lambda^, \lambda_{1\rho})$.*

Sketch of the proof.

Let $\lambda \in (0, \lambda_{1\rho})$ and Φ be a positive eigenfunction associated to the positive eigenvalue $\lambda_{1\rho}$. It follows that

$$I_\rho(t(\Phi, \lambda)\Phi) = \frac{C}{N_c} \left(1 - \frac{\lambda}{\lambda_{1\rho}}\right)^{N_c/p},$$

where C is a positive constant depending on the data of Problem 19 and $t(\cdot, \cdot)$ is defined in the same manner as (14) with minor changes. It is clear that there exists $\lambda^* \in (0, \lambda_{1\rho})$ such that $\forall \lambda \in (\lambda^*, \lambda_{1\rho})$:

$$0 < \frac{C}{N_c} \left(1 - \frac{\lambda}{\lambda_{1\rho}}\right)^{N_c/p} < \frac{1}{N_c} S_p^{\frac{N_c}{p}} |K|_\infty^{-\frac{N_c-p}{p}}.$$

Following the different steps of the previous example with slight changes, we get the result. \diamond

Acknowledgments. This work was done during the first author's "Délégation CNRS" at the "Équipe Analyse, Géométrie et Modélisation", Université de Cergy-Pontoise. He would like to thank them for the financial support and the nice working atmosphere during his stay.

References

- [1] **G. Bianchi, J. Chabrowski, A. Szulkin** *On symmetric solutions of an elliptic equation with a nonlinearity involving critical Sobolev exponent.* Nonlinear Anal. 25 (1995), no. 1, 41-59.
- [2] **H. Brézis, E. Lieb** *A relation between pointwise convergence of functions and convergence of functional* Proceedings of the AMS, vol 88, N°3 (July 1983).
- [3] **A. EL Hamidi** *Sur les conditions nécessaires et suffisantes d'existence de solutions pour des problèmes nonlinéaires,* Habilitation à Diriger des Recherches, Université de La Rochelle, (2004).
- [4] **A. EL Hamidi** *Multiple solutions with changing sign energy in a nonlinear equation,* Commun. Pure Appl. Anal. Vol 3, N° 2 , (2004) p.253-260.
- [5] **Fengquan Li, Zhao Huixiu** *Anisotropic parabolic equations with measure data,* J. Partial Diff. Eqs, vol 14 (2001) p. 21-30.
- [6] **E. Hebey, F. Robert** *Coercivity and Struwe's compactness for Paneitz type operators with constant coefficients,* Calc. Var. Partial Differential Equations 13 (2001), no. 4, 491-517
- [7] **H. Ohya** *Existence results for some quasilinear elliptic equations involving critical Sobolev exponents,* Adv. Diff. Equ. Vol 9, N° 11-12 (2004) p. 1339-1368.
- [8] **P. De Napoli, M. C. Mariani** *Mountain pass solutions to equations of p -Laplacian type,* Nonlinear Analysis, Vol 54 (2003) p.1205-1219.
- [9] **J.L. Lions** *Quelques méthodes de résolution de problèmes aux limites nonlinéaires,* Dunod Paris, (1969).
- [10] **J. Leray, J.L. Lions** *Quelques résultats de Visik sur les problèmes elliptiques non linéaires par les méthodes de Minty-Browder,* Bull. Soc. Math. France, Vol 93 (1965) p. 97-107.
- [11] **P.L. Lions** *The concentration-compactness principle in the calculus of variations. The limit case, I, II,* Rev. Mat. Iberoamericana, 1, 145-201 and 45-121, (1985).
- [12] **J.M. Rakotoson** *Compactness results for quasilinear elliptic problems,* Applied Math. Letters vol 4, (1991) p.33-35.
- [13] **J.M. Rakotoson** *Quasilinear elliptic problems with measure as data,* Diff. Int. Equa., Vol 4 N° 3 (1991) p.449-457.
- [14] **J.M. Rakotoson** *Equations et inéquations avec données mesures,* C.R.A.S, t 314 (1992) p. 105-107.
- [15] **J.M. Rakotoson** *Generalized solution in a new type sets for problems with measures as data,* Diff. Int. Equ., Vol 6, (1993) p. 27-36.
- [16] **S. Marchi** *Note on a paper of J.M. Rakotoson,* Equationi a derivate parziali, Istituto Lombardo (Rend. Sc.) A 127, (1991) p. 25-31.

- [17] **M. Struwe** *A global compactness result for elliptic boundary value problems involving limiting nonlinearities*, Math. Z. 187 (1984), no. 4, p. 511–517.
- [18] **L. Boccardo, F. Murat** *Almost everywhere convergence of the gradients of solutions to elliptic and parabolic equations*, Nonlinear Analysis, 19 (1992), p. 581–597.
- [19] **F. Demengel, E. Hebey** *On some nonlinear equations involving the p -Laplacian with critical Sobolev growth*, Adv. Differential Equations 3 (1998), no. 4, p. 533–574.
- [20] **F. Demengel, E. Hebey** *On some nonlinear equations involving the p -Laplacian with critical Sobolev growth and perturbation terms*, Appl. Anal. 72 (1999), no. 1-2, p. 75–109.

Liste des prépublications

- 99-1 Monique Jeanblanc et Nicolas Privault. A complete market model with Poisson and Brownian components. A paraître dans *Proceedings of the Seminar on Stochastic Analysis, Random Fields and Applications*, Ascona, 1999.
- 99-2 Laurence Cherfils et Alain Miranville. Generalized Cahn-Hilliard equations with a logarithmic free energy. A paraître dans *Revista de la Real Academia de Ciencias*.
- 99-3 Jean-Jacques Prat et Nicolas Privault. Explicit stochastic analysis of Brownian motion and point measures on Riemannian manifolds. *Journal of Functional Analysis* **167** (1999) 201-242.
- 99-4 Changgui Zhang. Sur la fonction q -Gamma de Jackson. A paraître dans *Aequationes Math.*
- 99-5 Nicolas Privault. A characterization of grand canonical Gibbs measures by duality. A paraître dans *Potential Analysis*.
- 99-6 Guy Wallet. La variété des équations surstables. A paraître dans *Bulletin de la Société Mathématique de France*.
- 99-7 Nicolas Privault et Jiang-Lun Wu. Poisson stochastic integration in Hilbert spaces. *Annales Mathématiques Blaise Pascal*, **6** (1999) 41-61.
- 99-8 Augustin Fruchard et Reinhard Schäfke. Sursabilité et résonance.
- 99-9 Nicolas Privault. Connections and curvature in the Riemannian geometry of configuration spaces. *C. R. Acad. Sci. Paris, Série I* **330** (2000) 899-904.
- 99-10 Fabienne Marotte et Changgui Zhang. Multisommabilité des séries entières solutions formelles d'une équation aux q -différences linéaire analytique. A paraître dans *Annales de l'Institut Fourier*, 2000.
- 99-11 Knut Aase, Bernt Øksendal, Nicolas Privault et Jan Ubøe. White noise generalizations of the Clark-Haussmann-Ocone theorem with application to mathematical finance. *Finance and Stochastics*, **4** (2000) 465-496.
- 00-01 Eric Benoît. Canards en un point pseudo-singulier nœud. A paraître dans *Bulletin de la Société Mathématique de France*.
- 00-02 Nicolas Privault. Hypothesis testing and Skorokhod stochastic integration. *Journal of Applied Probability*, **37** (2000) 560-574.
- 00-03 Changgui Zhang. La fonction thêta de Jacobi et la sommabilité des séries entières q -Gevrey, I. *C. R. Acad. Sci. Paris, Série I* **331** (2000) 31-34.
- 00-04 Guy Wallet. Déformation topologique par changement d'échelle.
- 00-05 Nicolas Privault. Quantum stochastic calculus for the uniform measure and Boolean convolution. A paraître dans *Séminaire de Probabilités XXXV*.
- 00-06 Changgui Zhang. Sur les fonctions q -Bessel de Jackson.
- 00-07 Laure Coutin, David Nualart et Ciprian A. Tudor. Tanaka formula for the fractional Brownian motion. A paraître dans *Stochastic Processes and their Applications*.
- 00-08 Nicolas Privault. On logarithmic Sobolev inequalities for normal martingales. *Annales de la Faculté des Sciences de Toulouse* **9** (2000) 509-518.
- 01-01 Emanuelle Augeraud-Veron et Laurent Augier. Stabilizing endogenous fluctuations by fiscal policies; Global analysis on piecewise continuous dynamical systems. A paraître dans *Studies in Nonlinear Dynamics and Econometrics*

- 01-02 Delphine Boucher. About the polynomial solutions of homogeneous linear differential equations depending on parameters. A paraître dans *Proceedings of the 1999 International Symposium on Symbolic and Algebraic Computation: ISSAC 99, Sam Dooley Ed., ACM, New York 1999*.
- 01-03 Nicolas Privault. Quasi-invariance for Lévy processes under anticipating shifts.
- 01-04 Nicolas Privault. Distribution-valued iterated gradient and chaotic decompositions of Poisson jump times functionals.
- 01-05 Christian Houdré et Nicolas Privault. Deviation inequalities: an approach via covariance representations.
- 01-06 Abdallah El Hamidi. Remarques sur les sentinelles pour les systèmes distribués
- 02-01 Eric Benoît, Abdallah El Hamidi et Augustin Fruchard. On combined asymptotic expansions in singular perturbation.
- 02-02 Rachid Bebbouchi et Eric Benoît. Equations différentielles et familles bien nées de courbes planes.
- 02-03 Abdallah El Hamidi et Gennady G. Laptev. Nonexistence of solutions to systems of higher-order semilinear inequalities in cone-like domains.
- 02-04 Hassan Lakhel, Youssef Ouknine, et Ciprian A. Tudor. Besov regularity for the indefinite Skorohod integral with respect to the fractional Brownian motion: the singular case.
- 02-05 Nicolas Privault et Jean-Claude Zambrini. Markovian bridges and reversible diffusions with jumps.
- 02-06 Abdallah El Hamidi et Gennady G. Laptev. Existence and Nonexistence Results for Reaction-Diffusion Equations in Product of Cones.
- 02-07 Guy Wallet. Nonstandard generic points.
- 02-08 Gilles Bailly-Maitre. On the monodromy representation of polynomials.
- 02-09 Abdallah El Hamidi. Necessary conditions for local and global solvability of nondiagonal degenerate systems.
- 02-10 Abdallah El Hamidi et Amira Obeid. Systems of Semilinear higher order evolution inequalities on the Heisenberg group.
- 03-01 Abdallah El Hamidi et Gennady G. Laptev. Non existence de solutions d'inéquations semilinéaires dans des domaines coniques.
- 03-02 Eris Benoît et Marie-Joëlle Rochet. A continuous model of biomass size spectra governed by predation and the effects of fishing on them.
- 03-03 Catherine Stenger: On a conjecture of Wolfgang Wasow concerning the nature of turning points.
- 03-04 Christian Houdré et Nicolas Privault. Surface measures and related functional inequalities on configuration spaces.
- 03-05 Abdallah El Hamidi et Mokhtar Kirane. Nonexistence results of solutions to systems of semilinear differential inequalities on the Heisenberg group.
- 03-06 Uwe Franz, Nicolas Privault et René Schott. Non-Gaussian Malliavin calculus on real Lie algebras.
- 04-01 Abdallah El Hamidi. Multiple solutions to a nonlinear elliptic equation involving Paneitz type operators.
- 04-02 Mohamed Amara, Amira Obeid et Guy Vallet. Relaxed formulation and existence result of the degenerated elliptic small disturbance model.
- 04-03 Hippolyte d'Albis et Emmanuelle Augeraud-Veron. Competitive Growth in a Life-cycle Model: Existence and Dynamics

- 04-04 Sadjia Aït-Mokhtar: Third order differential equations with fixed critical points.
- 04-05 Mokhtar Kirane et Nasser-eddine Tatar. Asymptotic Behavior for a Reaction Diffusion System with Unbounded Coefficients.
- 04-06 Mokhtar Kirane, Eric Nabana et Stanislav I. Pohozaev. Nonexistence of Global Solutions to an Elliptic Equation with a Dynamical Boundary Condition.
- 04-07 Khaled M. Furati, Nasser-eddine Tatar and Mokhtar Kirane. Existence and asymptotic behavior for a convection Problem.
- 04-08 José Alfredo López-Mimbela et Nicolas Privault. Blow-up and stability of semilinear PDE's with gamma generator.
- 04-09 Abdallah El Hamidi. Multiple solutions with changing sign energy to a nonlinear elliptic equation.
- 04-10 Sadjia Aït-Mokhtar: A singularly perturbed Riccati equation.
- 04-11 Mohamed Amara, Amira Obeid et Guy Vallet. Weighted Sobolev spaces for a degenerated nonlinear elliptic equation.
- 04-12 Abdallah El Hamidi. Existence results to elliptic systems with nonstandard growth conditions.
- 04-13 Eric Edo et Jean-Philippe Furter: Some families of polynomial automorphisms.
- 04-14 Laurence Cherfils et Yavdat Il'yasov. On the stationary solutions of generalized reaction diffusion equations with p & q - Laplacian.
- 04-15 Jean-Christophe Breton et Youri Davydov. Local limit theorem for supremum of an empirical processes for i.i.d. random variables.
- 04-16 Jean-Christophe Breton, Christian Houdré et Nicolas Privault. Dimension free and infinite variance tail estimates on Poisson space.
- 04-17 Abdallah El Hamidi et Gennady G. Laptev. Existence and nonexistence results for higher-order semilinear evolution inequalities with critical potential.
- 05-01 Mokhtar Kirane et Nasser-eddine Tatar. Nonexistence of Solutions to a Hyperbolic Equation with a Time Fractional Damping.
- 05-02 Mokhtar Kirane et Yamina Laskri. Nonexistence of Global Solutions to a Hyperbolic Equation with a Time Fractional Damping.
- 05-03 Mokhtar Kirane, Yamina Laskri et Nasser-eddine Tatar. Critical Exponents of Fujita Type for Certain Evolution Equations and Systems with Spatio-Temporal Fractional Derivatives.
- 05-04 Abdallah El Hamidi et Jean-Michel Rakotoson. Compactness and quasilinear problems with critical exponents