



## Prépublications du Département de Mathématiques

Université de La Rochelle  
Avenue Michel Crépeau  
17042 La Rochelle Cedex 1  
<http://www.univ-lr.fr/labo/lmca>

# Convergence in variation of the joint laws of multiple stable stochastic integrals

J-C Breton

Septembre 2006

**Classification:** 60G52, 60G57, 60H05, 60F99.

**Mots clés:** Convergence in variation, Multiple stochastic integrals, Stable process, LePage representation, Method of superstructure.

2006/05

# Convergence in variation of the joint laws of multiple stable stochastic integrals

Jean-Christophe BRETON\*

Laboratoire de Mathématiques et Applications  
Avenue Michel Crépeau, Université de La Rochelle  
17042 La Rochelle cedex, France  
E-mail: jcbreton@univ-lr.fr

## Abstract

In this note, we are interested in the regularity in the sense of total variation of the joint laws of multiple stable stochastic integrals. Namely, we show that the convergence

$$\mathcal{L}(I_{d_1}(f_1^n), \dots, I_{d_p}(f_p^n)) \xrightarrow{\text{var}} \mathcal{L}(I_{d_1}(f_1), \dots, I_{d_p}(f_p)), \quad n \rightarrow +\infty$$

holds true as long as each kernel  $f_i^n$  converges when  $n \rightarrow +\infty$  to  $f_i$  in the Lorentz-type space  $L^\alpha(\log_+)^{d_i-1}([0, 1]^{d_i})$  for  $1 \leq i \leq p$ . This result generalizes [4] from the one-dimensional case to the joint law case. It generalizes also [6] from the Wiener-Itô setting to the stable setting and [5] in the study of joint law of multiple stable integrals.

**Keywords:** Convergence in variation, Multiple stochastic integrals, Stable process, LePage representation, Method of superstructure.

**AMS classification:** 60G52, 60G57, 60H05, 60F99.

## 1 Introduction

In this paper, we deal with the regularity of the joint laws of multiple stable integrals (MSIs)

$$I_d(f) = \int_{[0,1]^d} f dM^d \tag{1.1}$$

with respect to their integrand  $f$ . Here and in the sequel,  $M$  is an  $\alpha$ -stable random measure on  $([0, 1], \mathcal{B}([0, 1]))$  defined for  $0 < \alpha < 2$  on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ :

$$M(A) \stackrel{\mathcal{L}}{=} S_\alpha \left( \lambda(A)^{1/\alpha}, \frac{\int_A \beta d\lambda}{\lambda(A)}, 0 \right), \quad A \in \mathcal{B}([0, 1])$$

---

\*Preprint

where  $\lambda$  is the Lebesgue measure and  $\beta : [0, 1] \rightarrow [-1, 1]$  is the skewness intensity of  $M$ , see Samorodnitsky-Taquq in [14, Section 3]. Moreover, for  $\alpha \geq 1$ , the measure  $M$  is assumed to be symmetric (that is:  $\beta = 0$ ).

The MSIs are a generalization of multiple Wiener-Itô integrals (MWI). A broad litterature is devoted to the study of MWIs and it is natural to investigate which properties of MWIs remain true in the stable case.

The MSI in (1.1) is defined for kernel  $f$  in a Lorentz-type space:

$$f \in L^\alpha(\log_+)^{d-1}([0, 1]^d) := \left\{ f : [0, 1]^d \rightarrow \mathbb{R} \mid \int_{[0,1]^d} |f|^\alpha (1 + \log_+ |f|)^{d-1} d\lambda^d \right\}$$

where  $\log_+ x := \log(x \vee 1)$ . The main feature of MSI is given by the Representation theorem which gives an insight into the discrete structure of MSI. It shows that  $I_d(f)$  can be represented in law by a multiple LePage-type series

$$S_d(f) = C_\alpha^{d/\alpha} \sum_{i_1, \dots, i_d > 0} \gamma_{i_1} \cdots \gamma_{i_d} \Gamma_{i_1}^{-1/\alpha} \cdots \Gamma_{i_d}^{-1/\alpha} f(V_{i_1}, \dots, V_{i_d}), \quad (1.2)$$

where  $C_\alpha = (\int_0^\infty x^{-\alpha} \sin x dx)^{-1}$  is a normalization factor,  $(\Gamma_i)_{i>0}$  is the sequence of arrival times of a standard Poisson process and  $(V_i, \gamma_i)_{i>0}$  are independent and identically distributed random vectors with  $V_i$  uniformly distributed on  $[0, 1]$  and  $\gamma_i = \pm 1$  with conditional laws

$$\mathbb{P}(\gamma_i = -1 \mid V_i) = \frac{1 - \beta(V_i)}{2}, \quad \mathbb{P}(\gamma_i = +1 \mid V_i) = \frac{1 + \beta(V_i)}{2}.$$

Moreover, the sequence  $(\Gamma_i)_{i>0}$  and  $(V_i, \gamma_i)_{i>0}$  are independent.

The Representation theorem shows that MSIs are also related to random multilinear forms (see [10]). For a complete account on the construction of MSI, we refer to [3] and references therein. The laws of MSIs have been studied by several authors. We briefly review some results on the law of MSIs.

In [13], the tails of  $I_d(f)$  is expressed in terms of  $f$ .

In [11], the regularity of the sample path of a process defined by an integral like in (1.1) is related to the smoothness of the kernel.

In [12], the independence of MSIs is studied in terms of the kernels, generalizing the MWI case of [15].

In [5], the existence of the densities for the joint laws of MSIs is studied, generalizing the MWI case of [8].

In this article, we go further in the study of the joint laws of MSIs than in [5] and we study their regularity in the sense of total variation norm. More precisely, given  $d_1, \dots, d_p$  the dimensions of  $p$  MSIs, we study the convergence in variation of the joint laws

$$(I_{d_1}(f_1^n), \dots, I_{d_p}(f_p^n)) \quad (1.3)$$

for integrands

$$f_1^n \rightarrow f_1 \text{ in } L^\alpha(\log_+)^{d_1-1}([0, 1]^{d_1}), \dots, f_p^n \rightarrow f_p \text{ in } L^\alpha(\log_+)^{d_p-1}([0, 1]^{d_p}). \quad (1.4)$$

This is a generalization of [4] which deals with one-dimensional law of MSIs ( $p = 1$  in our setting). This is also a generalization of [6] where the convergence in

variation of joint laws is investigated for MWIs ( $\alpha = 2$  in our setting). In this paper, we deal with arbitrary  $p \in \mathbb{N}^*$  and arbitrary  $\alpha \in (0, 2)$ .

Moreover, since the densities of the joint laws of MSIs exist (under rather broad conditions, see [5]), the convergence in variation of the law states also the convergence of the densities in  $L^1(\mathbb{R}^d)$ .

The paper is organized as follows. In Section 2, we start giving some notations yet used in [5]; they will be used all along this note. Next, we state the convergence result in Theorem 1. The sequel is devoted to the proof of Theorem 1. The problem is first reduced in Sections 3 and 4. In Section 5, we use the method of superstructure to reduce to the study of finite-dimensional functionals. Finally, in Section 6, the convergence in variation of these functionals is shown using the results of convergence in variation for smooth image-measures in [1] (see Proposition 1).

Note that the one-dimensional argument used in [4] (which states the one-dimensional counterpart of Theorem 1) can not be generalized in a multidimensional setting (at least easily). Actually, the proof of Theorem 1 relies on arguments yet used in [6] and in [5]. But this is not a simple rewriting of these arguments. Indeed, they have to be merged together: on the one hand, the method of stratification used in [5] is not sufficient to yield a convergence in variation, instead we use the method of superstructure, on the other hand, the argument in [6] relies on Gaussian analysis which has to be replaced by stable considerations. Moreover, new difficulties appear in the implementation of these merged arguments.

In the sequel a.s. stands for *almost surely*, a.e. for *almost everywhere*, i.i.d. for *independent and identically distributed*,  $:=$  means a definition,  $C$  is a finite and positive generic constant,  $\mu_A$  is the restriction to a measurable set  $A$  of a measure  $\mu$ ,  $\|\nu\|$  is the total variation of a signed measure  $\nu$ ,  $\xrightarrow{var}$  stands for the convergence of variation,  $\xrightarrow{P}$  for the convergence in probability  $P$  and finally bold characters are used for multi-indicial notations.

## 2 Convergence in variation of joint laws

In this study, we shall use the same background as in [5]. We begin by reminding of the notations of [5] that will be used all along this article:

- for  $i = 1, \dots, p$ ,  $N_i = d_1 + \dots + d_i$ ,  $N = N_p$ ,
- $\mathbf{a}^i = (a_0^i, \dots, a_p^i) \in \mathbb{N}^{p+1}$  a  $(p+1)$ -partition of  $d_i$ :  $d_i = |\mathbf{a}^i| = a_0^i + \dots + a_p^i$ ,
- $\mathbf{a} = (\mathbf{a}^1, \dots, \mathbf{a}^p) \in (\mathbb{N}^{p+1})^p$ ,
- $M_{\mathbf{a}} = (a_j^i)_{1 \leq i, j \leq p}$  a  $p$ -square matrix with  $d_i = \sum_{k=0}^p a_k^i$  for  $1 \leq i \leq p$  and  $b_k = \sum_{i=1}^p a_k^i$  for  $0 \leq k \leq p$ ,
- for  $\mathbf{b} = (b_1, \dots, b_p) \in \mathbb{N}^p$ :

$$E(\mathbf{b}) = \left\{ \mathbf{a} = (\mathbf{a}^1, \dots, \mathbf{a}^p) \mid a_1^i + \dots + a_p^i = d_i, \sum_{i=1}^p a_k^i = b_k, k = 1, \dots, p \right\},$$

- $\sigma_{\mathbf{a}}$  the permutation of  $\{1, \dots, N\}$  that sends for each  $i, k$

$$j = \sum_{u=1}^{k-1} b_u + \sum_{s=1}^{i-1} a_k^s + l, \quad l = 1, \dots, a_k^i,$$

to

$$\sigma_{\mathbf{a}}(j) = \sum_{v=1}^{i-1} d_i + \sum_{s=1}^{k-1} a_s^i + l,$$

- $U_{\mathbf{a}} : \mathbb{R}^N \longrightarrow \mathbb{R}^N$  associated to  $\sigma_{\mathbf{a}}$  by  $U_{\mathbf{a}}(t_1, \dots, t_N) = (t_{\sigma_{\mathbf{a}}(1)}, \dots, t_{\sigma_{\mathbf{a}}(N)})$ ,
- denoting  $\Pi_{b_1, \dots, b_d}$  the sub-group of  $\Pi_N$  of permutations preserving the following "b-blocks":  $(1, \dots, b_1), (b_1 + 1, \dots, b_1 + b_2), \dots, (b_1 + b_2 + \dots + b_{p-1} + 1, \dots, b_1 + b_2 + \dots + b_p = N)$ :

$$S_{b_1, \dots, b_d} \phi_{\mathbf{b}}(t) = \frac{b_1! \cdots b_d!}{N!} \sum_{\sigma \in \Pi_{b_1, \dots, b_d}} \sum_{\mathbf{a} \in E(\mathbf{b})} \prod_{i=1}^p \frac{d_i!}{a_0^i! \cdots a_p^i!} \det M_{\mathbf{a}} \phi(U_{\mathbf{a}}(t))$$

where  $\phi(t) = \phi(t_1, \dots, t_N) = f_1(t_1, \dots, t_{N_1}) \cdots f_p(t_{N_{p-1}+1}, \dots, t_{N_p})$ .

Note that the previous function  $S_{b_1, \dots, b_d} \phi_{\mathbf{b}}$  is symmetric in each  $\mathbf{b}$ -blocks.

The main result of this paper is:

**Theorem 1** *Let  $f_1^n, \dots, f_p^n$  be kernels converging respectively to  $f_1, \dots, f_p$  like in (1.4). Suppose moreover the limit functions  $f_1, \dots, f_p$  satisfy the following hypothesis:*

$$(\mathbf{H}) \begin{cases} S_{b_1, \dots, b_d} \phi_{\mathbf{b}} \neq 0 \text{ a.e. on } [0, 1]^N \text{ for some } \mathbf{b} = (b_1, \dots, b_p) \in (\mathbb{N}^*)^p \\ \text{with } |\mathbf{b}| = N = d_1 + \dots + d_p. \end{cases}$$

Then  $\mathcal{L}(I_{d_1}(f_1^n), \dots, I_{d_p}(f_p^n)) \xrightarrow{\text{var}} \mathcal{L}(I_{d_1}(f_1), \dots, I_{d_p}(f_p))$  when  $n \rightarrow +\infty$ .

Roughly speaking,  $(\mathbf{H})$  is a non-degeneracy condition dealing with how overlapped are the  $f_i$ 's. The same remark and the same examples as in [5] apply about condition  $(\mathbf{H})$ . In particular, we stress on that this condition is optimal in several examples and coincides with the condition for the same convergence for joint laws of MWIs, see [6]. For instance,

- for  $p = 1$  and  $d_1 = d$ ,  $(\mathbf{H})$  is satisfied with  $\mathbf{b} = d$  if  $f \neq 0$  a.e.
- for  $p > 0$  and  $d_1 = \dots = d_p = 1$ ,  $(\mathbf{H})$  is satisfied with  $\mathbf{b} = (1, 1)$  if  $\det\{(f_i(t_j))_{1 \leq i, j \leq p}\} \neq 0$  a.e.
- for  $p = 2$  and  $d_1 = d_2 = 2$ ,  $(\mathbf{H})$  is satisfied with  $\mathbf{b} = (2, 2)$  if  $f_1$  and  $f_2$  are not proportional a.e.

The global scheme of the proof is the following. First, we explain how to reduce the problem in Section 3 (see (3.9)). Using the representation of stable integrals by LePage series, we introduce on the Skorohod space some related functionals to be studied (see (3.6)). Next, approximating and localizing the problem in Section 4, we use the method of superstructure in Section 5 where the main point is to study the convergence in variation of measures under finite-dimensional mappings (see (5.24)). This is finally done in Section 6 with the following result from [1] and the study of some related coefficients (see (6.26)).

**Proposition 1 (Corollary 4 in [1])** *Let  $F_j, F \in W_{loc}^{p,1}(\mathbb{R}^n, \mathbb{R}^n)$ , where  $p \geq n$ , and let the mappings  $F_j$  converge to  $F$  with respect to the Sobolev norm  $\|\cdot\|_{p,1}$  on every ball. Assume that  $E \subset \{\det DF \neq 0\}$  is a set of finite Lebesgue measure. Then  $\lambda|_E F_j^{-1} \xrightarrow{var} \lambda|_E F^{-1}$ .*

### 3 Reduction of the problem

In this section, we describe the arguments yet used in [3] and in [5] to reduce the study of the convergence in variation of laws like in (1.3).

**Representation and stable stuff.** Like in [5], we first reduce the study to random multiple LePage series. From the Representation Theorem [3, Th. 3.2], we have like in (1.2) the following equality of joint laws

$$(S_{d_1}(f_1), \dots, S_{d_p}(f_p)) \stackrel{\mathcal{L}}{=} (I_{d_1}(f_1), \dots, I_{d_p}(f_p)). \quad (3.5)$$

Moreover, we have from [3, Sec. 4.1.2 and 4.2.3]:

**Proposition 2** *Let  $(I_{d_1}(f_1^n), \dots, I_{d_p}(f_p^n))$  be a vector of MSIs with kernels  $f_1^n, \dots, f_p^n$  converging like in (1.4). Then, when  $n \rightarrow +\infty$ , we have*

$$(I_{d_1}(f_1^n), \dots, I_{d_p}(f_p^n)) \xrightarrow{\mathbb{P}} (I_{d_1}(f_1), \dots, I_{d_p}(f_p)).$$

Thanks to (3.5), we shall actually study the joint law of  $(S_{d_1}(f_1^n), \dots, S_{d_p}(f_p^n))$ . For  $x$  in the Skorohod space  $\mathbb{D}$  (the space of *cadlag* functions on  $[0, 1]$ ), let  $\delta_x(t)$  be the jump of  $x$  at  $t$  and  $(t_i)_{i>0}$  is the list of its jump-times. We consider the multi-dimensional functional  $F = (F_1, \dots, F_p)$  with  $F_i : \mathbb{D} \rightarrow \mathbb{R}$  given by:

$$F_i(x) = \sum_{t_1, \dots, t_{d_i}} \delta_x(t_1) \cdots \delta_x(t_{d_i}) f_i(t_1, \dots, t_{d_i}) \quad (3.6)$$

whenever the multiple series is convergent, otherwise  $F_i(x) = 0$ .

In the sequel, we shall also consider the stable standard process  $\eta$  given by

$$\eta_t = M([0, t]), \quad t \in [0, 1].$$

The sample paths of  $\eta$  live in  $\mathbb{D}$ ; we denote its law by  $P$ . From the Representation Theorem (in a one-dimensional case), we have

$$\eta_t = \int_{[0,t]} \mathbf{1}_{[0,t]} dM \stackrel{\mathcal{L}}{=} C_\alpha^{1/\alpha} \sum_{i>0} \gamma_i \Gamma_i^{-1/\alpha} \mathbf{1}_{[0,t]}(V_i)$$

from which the following interpretations come

- $V_i, i > 0$ , are the jump-times of the stable process  $\eta$ ;
- $C_\alpha^{1/\alpha} \Gamma_i^{-1/\alpha}$  is the modulus of the jump at  $V_i$ , decreasingly ordered;
- $\gamma_i$  indicates the direction of the jump.

We deduce

$$\begin{aligned} F_i(\eta(\omega)) &= C_\alpha^{d_i/\alpha} \sum_{k_1, \dots, k_{d_i} > 0} (\gamma_{k_1} \Gamma_{k_1}^{-1/\alpha}) \cdots (\gamma_{k_{d_i}} \Gamma_{k_{d_i}}^{-1/\alpha}) f_i(V_{k_1}, \dots, V_{k_{d_i}}) \\ &= S_{d_i}(f_i)(\omega), \end{aligned}$$

so that

$$F(\eta) \stackrel{\mathcal{L}}{=} (S_{d_1}(f_1), \dots, S_{d_p}(f_p)).$$

We define also  $F^n$  from  $(f_1^n, \dots, f_p^n)$  like  $F$  from  $(f_1, \dots, f_p)$  in (3.6). The convergence in variation of the law of (1.3) actually rewrites, in our notations:

$$P(F^n)^{-1} \xrightarrow{\text{var}} PF^{-1}, \quad n \rightarrow +\infty. \quad (3.7)$$

In the sequel, we shall use the following result. This is a rewriting of Prop. 2 in terms of the functionals related to the corresponding MSIs.

**Proposition 3** *Let  $f_1^n, \dots, f_p^n$  be converging kernels like in (1.4). Then, with the previous notations, we have  $F^n \xrightarrow{P} F$  when  $n \rightarrow +\infty$ .*

**Approximation.** This procedure consists in the following straightforward result:

**Proposition 4 (Approximation)** *In order to prove (3.7), it is enough to see for all  $\varepsilon > 0$ , there is some measurable set  $\mathbb{D}(\varepsilon)$  in  $\mathbb{D}$  with  $P(\mathbb{D}(\varepsilon)) > 1 - \varepsilon$  and*

$$P_{\mathbb{D}(\varepsilon)}(F^n)^{-1} \xrightarrow{\text{var}} P_{\mathbb{D}(\varepsilon)}F^{-1}. \quad (3.8)$$

**Proof:** We have

$$\begin{aligned} &\|P(F^n)^{-1} - PF^{-1}\| \\ &\leq \|P(F^n)^{-1} - P_{\mathbb{D}(\varepsilon)}(F^n)^{-1}\| + \|P_{\mathbb{D}(\varepsilon)}(F^n)^{-1} - P_{\mathbb{D}(\varepsilon)}F^{-1}\| + \|P_{\mathbb{D}(\varepsilon)}F^{-1} - PF^{-1}\| \\ &\leq 2P(\mathbb{D}(\varepsilon)^c) + \|P_{\mathbb{D}(\varepsilon)}(F^n)^{-1} - P_{\mathbb{D}(\varepsilon)}F^{-1}\| \\ &\leq 2\varepsilon + \|P_{\mathbb{D}(\varepsilon)}(F^n)^{-1} - P_{\mathbb{D}(\varepsilon)}F^{-1}\|. \end{aligned}$$

But from (3.8), the last bound is bounded by  $3\varepsilon$  for  $n$  large enough. This ends the proof of the argument of approximation.  $\square$

**Localization.** Using the separability of  $\mathbb{D}(\varepsilon)$ , we localize the problem by the following result:

**Proposition 5 (Localization)** *In order to prove (3.8), it is enough to exhibit for all  $x \in \mathbb{D}(\varepsilon)$  some neighbourhood  $V(x)$  of  $x$  such that*

$$P_{V(x)}(F^n)^{-1} \xrightarrow{\text{var}} P_{V(x)}F^{-1}, \quad n \rightarrow +\infty. \quad (3.9)$$

**Proof:** Since  $\mathbb{D}(\varepsilon)$  is separable, there is a countable family  $\{x_i, i \in \mathbb{N}^*\}$  such that  $\mathbb{D}(\varepsilon) = \bigcup_{i=1}^{+\infty} V(x_i)$ . We have  $\lim_{k \rightarrow +\infty} P(\bigcup_{i=1}^k V(x_i)) = P(\mathbb{D}(\varepsilon))$  so that for any

fixed  $\epsilon > 0$  and  $k$  large enough, we have  $P(\mathbb{D}(\epsilon) \setminus A_k) < \epsilon$  where  $A_k = \bigcup_{i=1}^k V(x_i)$ . Therefore, for such a  $k$ , we have:

$$\begin{aligned}
& \|P_{\mathbb{D}(\epsilon)}(F^n)^{-1} - P_{\mathbb{D}(\epsilon)}F^{-1}\| \\
& \leq \|P_{\mathbb{D}(\epsilon)}(F^n)^{-1} - P_{A_k}(F^n)^{-1}\| + \|P_{A_k}(F^n)^{-1} - P_{A_k}F^{-1}\| + \|P_{A_k}(F)^{-1} - P_{\mathbb{D}(\epsilon)}F^{-1}\| \\
& \leq 2P(\mathbb{D}(\epsilon) \setminus A_k) + \|P_{A_k}(F^n)^{-1} - P_{A_k}F^{-1}\| \\
& \leq 2\epsilon + \sum_{i=1}^k \|P_{V(x_i)}(F^n)^{-1} - P_{V(x_i)}F^{-1}\| \\
& \leq 3\epsilon
\end{aligned}$$

where the last bound comes for  $n$  large enough from (3.9). Finally, we derive (3.8) and this proves the localization.  $\square$

Using localization, it is enough to prove (3.9). To do so, we shall use the method of superstructure. For a general description of this method, we refer to [9]. Like in [3, 5] with the method of stratification, these preliminary procedures of approximation and of localization are necessary in order to implement successfully the method of superstructure.

## 4 Approximation and localization

In this section, we exhibit the set  $\mathbb{D}(\epsilon)$  required in the approximation procedure and the neighbourhood  $V(x)$  required for  $P$ -almost all  $x \in \mathbb{D}(\epsilon)$  in the localization procedure. Actually, the approximation and the localization procedures are the same as in [5], we thus refer to the Section 3 of [5] for a precise description. Here, we only sketch the main steps.

**Approximation.** Let  $\mathbf{b}$  be given by hypothesis **(H)** in Theorem 1 and  $\tilde{t} = (\tilde{t}_1, \dots, \tilde{t}_N)$  be some Lebesgue point of  $A_{\mathbf{b}} = \{t \in [0, 1]^N \mid S_{b_1, \dots, b_p} \phi_b(t) \neq 0\} \in \mathcal{B}([0, 1]^N)$ . The Lebesgue measure of  $A_{\mathbf{b}}$  is positive by hypothesis **(H)**. There is no restriction in assuming  $\tilde{t}$  is chosen with its coordinates all distinct ( $\tilde{t}_i \neq \tilde{t}_j, i \neq j$ ). Let  $\epsilon > 0$  be fixed, there is a product neighbourhood  $V_\epsilon = U_1^\epsilon \times \dots \times U_N^\epsilon$  of  $\tilde{t}$  in  $[0, 1]^N$  satisfying

$$U_i^\epsilon \cap U_j^\epsilon = \emptyset, i \neq j \quad \text{and} \quad \frac{\lambda^N(V_\epsilon \cap A_{\mathbf{b}})}{\lambda^N(V_\epsilon)} \geq 1 - \epsilon. \quad (4.10)$$

We consider the following sets:

$$\tilde{\mathbb{D}}(\epsilon) = \{x \in \mathbb{D} \mid \text{for } i = 1, 2, \dots, N, x \text{ has at least one jump at a time in } U_i^\epsilon, \text{ the maximal modulus of these jumps being realized only once}\},$$

$$\mathbb{D}(\epsilon) = \{x \in \tilde{\mathbb{D}}(\epsilon) \mid x \text{ has an unique maximal jump on each } U_i^\epsilon \text{ at } T_{U_i^\epsilon}(x) \text{ with } T_\epsilon(x) := (T_{U_1^\epsilon}(x), \dots, T_{U_N^\epsilon}(x)) \in A_{\mathbf{b}}\}.$$

We recall from [5] the following result for the standard stable process  $\eta$ :

**Lemma 1** *The random vector  $T_\epsilon(\eta) = (T_{U_1^\epsilon}(\eta), \dots, T_{U_N^\epsilon}(\eta))$  is uniformly distributed on  $V_\epsilon$ . Moreover, for  $i \neq j$ ,  $T_{U_i^\epsilon}(\eta)$  and  $T_{U_j^\epsilon}(\eta)$  are independent.*



With (4.10), Lemma 1 gives:

$$P(\mathbb{D}(\varepsilon)) = P_{\mathbb{D}(\varepsilon)} T_\varepsilon^{-1}(A_{\mathbf{b}}) = \frac{\lambda^N(V_\varepsilon \cap A_{\mathbf{b}})}{\lambda^N(V_\varepsilon)} \geq 1 - \varepsilon.$$

The set  $\mathbb{D}(\varepsilon)$  is the set required in the procedure of approximation of  $\mathbb{D}$ .

**Localization.** For the sake of completeness of notations, we recall the localization procedure of [5]. Let  $x \in \mathbb{D}(\varepsilon)$  be fixed and denote for  $i = 1, \dots, N$ :

- $t_i = T_{U_i^\varepsilon}(x)$  the time of the largest jump of  $x$  in  $U_i^\varepsilon$ ;
- $t'_i$  the time of the second largest jump of  $x$  in  $U_i^\varepsilon$ ,  $|\delta_x(t'_i)| < |\delta_x(t_i)|$ ;
- $\varepsilon_0 = \frac{1}{2} \min_{i=1, \dots, N} |\delta_x(t_i)|$ .

Note that by Lemma 1, the jump-time  $t_i$  can be seen as a random variable on  $(\mathbb{D}(\varepsilon), P_{\mathbb{D}(\varepsilon)}/P(\mathbb{D}(\varepsilon)))$  whose law is uniform on  $U_i^\varepsilon$ .

By finiteness of the number of jumps of  $x$  larger than  $\varepsilon_0/2$ , we select  $\delta_1 > 0$  such that  $t_i$  is the unique time of  $\Delta'_i := (t_i - \delta_1, t_i + \delta_1) \subset U_i^\varepsilon$  where a jump larger than  $\varepsilon_0/2$  in modulus occurs. Let the following technical conditions be fulfilled:

$$\bullet \varepsilon_0/2 < \varepsilon_1 < \varepsilon_2 < \dots < \varepsilon_p < \varepsilon_0; \quad (4.11)$$

$$\bullet \delta_2 < \frac{1}{4} \min \left\{ \varepsilon_0, 2\delta_1, \inf_{i=1, \dots, N} \{|\delta_x(t_i)| - |\delta_x(t'_i)|\}, 2\varepsilon_1 - \varepsilon_0 \right\}; \quad (4.12)$$

$$\bullet \beta = \delta_1 - \delta_2 \quad (\delta_2 \leq \beta \leq \delta_1);$$

$$\bullet \Delta_i := (t_i - \beta, t_i + \beta) \subset \Delta'_i \subset U_i^\varepsilon.$$

In the sequel, we consider local field  $l = (l_x)_{x \in \mathbb{D}}$ . That is, for  $m \in \mathbb{N}^*$ ,  $\varepsilon > 0$ , reals  $\tau_i$ , and intervals  $\Delta_i = (a_i, b_i)$ , we define

$$l_x(t) = \sum_{s|\delta_x(s) > \varepsilon} \left( \sum_{i=1}^m \tau_i \mathbf{1}_{\Delta_i}(s) \mathbf{1}_{[s, \infty[}(t) \right)^+ - \sum_{s|\delta_x(s) < -\varepsilon} \left( \sum_{i=1}^m \tau_i \mathbf{1}_{\Delta_i}(s) \mathbf{1}_{[s, \infty[}(t) \right)^-$$

see [9, p. 163] for a precise definition of local fields. Roughly speaking, local fields  $(l_x)_x$  are admissible directions for stable processes. Moreover, we note

$$\omega_x(t) = \begin{cases} \tau_i & \text{if } t \in (a_i, b_i), \quad |\delta_x(t)| > \varepsilon, \quad \delta_x(t) \tau_i > 0, \\ 0 & \text{else,} \end{cases} \quad (4.13)$$

so that the jumps of  $x$  and  $x + c l_x$  are linked by  $\delta_{x+c l_x}(t) = \delta_x(t) + c \omega_x(t)$ .

Next, we associate the following sets to  $l$ :

- $A(l)^+$  the set of  $x \in \mathbb{D}$  such that for all  $i$  with  $\tau_i > 0$ ,  $x$  does not have jumps of length exactly  $\varepsilon$  on  $(a_i, b_i)$ ,  $\delta_x(a_i) < \varepsilon$ ,  $\delta_x(b_i) < \varepsilon$ , and  $x$  has at least one jump larger than  $\varepsilon$  on  $(a_i, b_i)$ ,
- $A(l)^-$  the set of  $x \in \mathbb{D}$  such that for all  $i$  with  $\tau_i < 0$ ,  $x$  does not have jumps of length exactly  $-\varepsilon$  on  $(a_i, b_i)$ ,  $\delta_x(a_i) > -\varepsilon$ ,  $\delta_x(b_i) > -\varepsilon$ , and  $x$  has at least one jump lower than  $-\varepsilon$  on  $(a_i, b_i)$ ,

$$A(l) = A(l)^+ \cap A(l)^-. \quad (4.14)$$

The set  $A(l)$  is suitable to study the local field  $l$ . In particular, it is shown in [5, Sec. A1, A2] that  $A(l)$  is open in  $\mathbb{D}$  and that the local field  $l$  is continuous on  $A(l)$ .

In order to apply the method of superstructure in a multi-dimensional setting, we consider  $p$  local fields  $l^i$ ,  $1 \leq i \leq p$ , and their open set  $A(l^i)$ , given like in (4.14). We select the  $p$  local fields with the following parameters: for  $i = 1, \dots, p$ ,

- $\varepsilon_i$  given by (4.11),
- $m_i = b_i$ , given by hypothesis **(H)**,
- $\Delta_j^i = \Delta_{b_1+\dots+b_{i-1}+j}$  for  $j = 1, \dots, b_i$ ,
- $\tau_j^i$  with the same sign as  $\delta_x(t_i)$  and with constant modulus  $\tau > 0$ .

In the sequel, we note  $\ell = (l^1, \dots, l^p)$ . We have  $x \in \tilde{A}(\ell) := \bigcap_{i=1}^p A(l^i)$ , open set. We shall apply the localization procedure with the following neighbourhood  $V(x)$ :

$$V(x) = B(x, \delta_2) \cap \tilde{A}(\ell) \cap \mathbb{D}(\varepsilon) \quad (4.15)$$

where  $\delta_2$  is given in (4.12).

Finally, with  $\mathbb{D}(\varepsilon)$  and  $V(x)$  given above, Proposition 4 and Proposition 5 applies and the proof of Theorem 1 reduces to (3.9). The convergence in (3.9) is tackled with the method of superstructure in the next sections.

## 5 Superstructure in $\mathbb{D}(\varepsilon)$

In order to prove (3.9), we use the method of superstructure in the neighbourhood  $V(x)$  of  $x$  defined in (4.15). For a general account on this method we refer to [9, Sec. 5]. Here, we only sketch the method.

This method applies to study the convergence  $PF_n^{-1} \xrightarrow{var} PF^{-1}$  when  $F_n$  and  $F$  are some functionals on some  $(\mathcal{Y}, P)$ . When we have a family of transformations  $(G_{\mathbf{c}})_{\mathbf{c} \in (\mathbb{R}_+)^p}$ , satisfying

$$PG_{\mathbf{c}}^{-1} \xrightarrow{var} P, \quad \mathbf{c} \rightarrow 0, \quad (5.16)$$

we define the following auxiliary measures and functionals on the product space  $\mathcal{Y}_\varepsilon = [0, \varepsilon]^p \times \mathcal{Y}$ ,  $\varepsilon > 0$ :

$$\begin{aligned} Q_\varepsilon &= \frac{1}{\varepsilon^p} \lambda_{[0, \varepsilon]^p} \otimes P, \\ F_\varepsilon(\mathbf{c}, y) &= F(G_{\mathbf{c}}(y)), \end{aligned}$$

(note that  $F_\varepsilon$  depends on  $\varepsilon$  only through its domain of definition  $\mathcal{Y}_\varepsilon$ ).

Since  $Q_\varepsilon F_\varepsilon^{-1} = \frac{1}{\varepsilon^p} \int_{[0, \varepsilon]^p} PG_{\mathbf{c}}^{-1} F^{-1} d\mathbf{c}$ , for the total variation, we derive:

$$\|Q_\varepsilon F_\varepsilon^{-1} - PF^{-1}\| \leq \frac{1}{\varepsilon^p} \int_{[0, \varepsilon]^p} \|P - PG_{\mathbf{c}}^{-1}\| d\mathbf{c}$$

and from (5.16) together with the dominated convergence

$$\lim_{\epsilon \rightarrow 0} \|Q_\epsilon F_\epsilon^{-1} - P F^{-1}\| = 0. \quad (5.17)$$

Next, we express  $Q_\epsilon F_\epsilon^{-1}$  as a mixture of finite-dimensional measures: we introduce  $\varphi_y(\mathbf{c}) = F(G_\mathbf{c}(y))$  for  $\mathbf{c} \in [0, \epsilon]^p$  and we have:

$$Q_\epsilon F_\epsilon^{-1} = \frac{1}{\epsilon^p} \int_{\mathcal{Y}} \lambda_{[0, \epsilon]^p} \varphi_y^{-1} dP. \quad (5.18)$$

In the sequel, we shall note  $\varphi_y = (\varphi_y^1, \dots, \varphi_y^p)$ . In our setting, in order to study the laws of  $(I_{d_1}(f_1^n), \dots, I_{d_p}(f_p^n))_n$  for  $f_i^n \rightarrow f$ ,  $n \rightarrow +\infty$ , in the space  $L^\alpha(\log_+)^{d_i-1}([0, 1]^{d_i})$ ,  $1 \leq i \leq p$ , we apply this method to  $\mathcal{Y} = \tilde{A}(\ell)$  equipped with the restricted probability law  $P_x := P_{V(x)}$  of the process  $\eta$  and to the functionals  $F$  and  $F^n$  given in (3.7). We use the family of transformations  $(G_\mathbf{c})_{\mathbf{c} \in (\mathbb{R}_+)^p}$  defined from the local fields  $l^i$ ,  $i = 1, \dots, p$ , by

$$G_\mathbf{c} : \begin{cases} \tilde{A}(\ell) & \longrightarrow \tilde{A}(\ell) \\ y & \longmapsto y + \langle \mathbf{c}, l_y \rangle \end{cases} \quad (5.19)$$

where  $\langle \mathbf{c}, l_y \rangle := c_1 l_y^1 + \dots + c_p l_y^p$  for  $\mathbf{c} = (c_1, \dots, c_p)$ . Moreover, with a similar notation, note that with  $\omega^i$  defined from  $l^i$  like in (4.13), we have for  $\mathbf{c} \in (\mathbb{R}_+)^p$ ,

$$\delta_{G_\mathbf{c}(y)}(t) = \delta_y(t) + \langle \mathbf{c}, \omega_y \rangle. \quad (5.20)$$

**Remark 1** The open set  $\tilde{A}(\ell)$  is invariant under  $G_{c_1, \dots, c_p}$ . Roughly speaking, this is because each term  $l_y^i$ ,  $1 \leq i \leq p$ , emphasizes the membership of  $A(l^i)$  and does not alter conditions to belong to  $A(l^j)$  for  $j \neq i$ . A more detailed justification is given in [3].

The condition (5.16) in our setting is satisfied for the family of transformations (5.19). Indeed, in Lemma 4.1 of [3], it is shown that  $(G_\mathbf{c})_\mathbf{c}$  defined an admissible semigroup in the sense of [9, p. 14]. Moreover, the conditional measures of  $P$  on the orbits of the semigroup  $(G_\mathbf{c})_\mathbf{c}$  have densities. Since  $(G_\mathbf{c})_\mathbf{c}$  acts as a translation, we derive (5.16) when  $\mathbf{c} \rightarrow 0$  (see (21.8) in [9] for a more detailed proof). The convergence in (5.16) is enough for our study but we have actually more:

**Lemma 2** *The convergence  $P^x G_\mathbf{c}^{-1} \xrightarrow{\text{var}} P^x$  is uniform with respect to  $x$  in  $\tilde{A}(\ell)$ .*

**Proof:** In this proof only, we use the setting described in Section 4 of [9]. First, we define an equivalence relation  $\mathcal{R}$  on  $\tilde{A}(\ell)$  by  $x_1 \mathcal{R} x_2$  if and only if there are  $\mathbf{c}^1, \mathbf{c}^2 \in (\mathbb{R}_+)^p$  such that  $G_{\mathbf{c}^1} x_1 = G_{\mathbf{c}^2} x_2$ . Let  $\Gamma$  be the partition given by  $\mathcal{R}$  and  $\pi : \tilde{A}(\ell) \rightarrow \tilde{A}(\ell)/\Gamma$  be the canonical projection. The equivalence classes  $\pi^{-1}(\gamma)$ ,  $\gamma \in \tilde{A}(\ell)/\Gamma$ , are called orbits of the semigroup  $(G_\mathbf{c})_\mathbf{c}$ . In Proposition 4.2 of [9], each orbit  $\pi^{-1}(\gamma)$  is shown to be isomorphic (via a mapping  $J_\gamma$ ) to a measurable set  $C_\gamma \subset \mathbb{R}^p$ . Next, we define a Lebesgue measure  $\lambda_\gamma$  on  $\pi^{-1}(\gamma)$  by

$$\lambda_\gamma(B) = \lambda_p(J_\gamma(B \cap \pi^{-1}(\gamma))), \quad B \in \mathcal{B}(\tilde{A}(\ell))$$

where  $\lambda_p$  is the Lebesgue measure on  $\mathbb{R}^p$ . Moreover, the mapping  $J_\gamma$  intertwines the action of the semigroup  $(G_\mathbf{c})_\mathbf{c}$  on the orbit  $\pi^{-1}(\gamma)$  with the action of the semigroup of translation  $(\tau_\mathbf{c})_\mathbf{c}$  on  $C_\gamma$ :

$$J_\gamma G_\mathbf{c} = \tau_\mathbf{c} J_\gamma.$$

Thus, we have

$$\begin{aligned}
\|P^x G_{\mathbf{c}}^{-1} - P^x\| &= \left\| \int_{\tilde{A}(\ell)/\Gamma} P_\gamma^x G_{\mathbf{c}}^{-1} dP_\Gamma(\gamma) - \int_{\tilde{A}(\ell)/\Gamma} P_\gamma^x dP_\Gamma(\gamma) \right\| \\
&\leq \int_{\tilde{A}(\ell)/\Gamma} \|P_\gamma^x G_{\mathbf{c}}^{-1} - P_\gamma^x\| dP_\Gamma(\gamma) \\
&\leq \int_{\tilde{A}(\ell)/\Gamma} \|P_\gamma^x J_\gamma^{-1} \tau_{\mathbf{c}}^{-1} J_\gamma - P_\gamma^x J_\gamma^{-1} J_\gamma\| dP_\Gamma(\gamma) \\
&\leq \int_{\tilde{A}(\ell)/\Gamma} \|P_\gamma^x J_\gamma^{-1} \tau_{\mathbf{c}}^{-1} - P_\gamma^x J_\gamma^{-1}\| dP_\Gamma(\gamma). \tag{5.21}
\end{aligned}$$

In Theorem 4.1 of [9], it is shown that the conditional measures  $(P_\gamma)$  of  $P$  on the orbits of the semigroup  $(G_{\mathbf{c}})_{\mathbf{c}}$  have densities. Thus  $P_\gamma J_\gamma^{-1}$  has also a density in  $C_\gamma \subset \mathbb{R}^p$ . But the translation operator is uniformly continuous in  $L^1(\mathbb{R}^p)$  (that is  $\lim_{h \rightarrow 0} \varphi(\cdot + h) = \varphi(\cdot)$  in  $L^1(\mathbb{R}^p)$  uniformly with respect to  $\varphi$ ).

Therefore,  $\lim_{\mathbf{c} \rightarrow 0} \|P_\gamma^x J_\gamma^{-1} \tau_{\mathbf{c}}^{-1} - P_\gamma^x J_\gamma^{-1}\| = 0$  holds uniformly with respect to both  $\gamma$  and  $x$ . Integrating with respect to  $\gamma \in \tilde{A}(\ell)/\Gamma$ , we derive that the right-hand side of (5.21) goes to 0 when  $\mathbf{c} \rightarrow 0$  uniformly with respect to  $x$ . Finally (5.16) holds uniformly with respect to  $x$  in  $\tilde{A}(\ell)$ .  $\square$

Applying the method of superstructure in this setting, we define as previously multi-dimensional auxiliary functionals  $F_\epsilon^n$  on  $\mathcal{Y}_\epsilon$  and we derive as in (5.17):

$$\|Q_\epsilon(F_\epsilon^n)^{-1} - P^x(F^n)^{-1}\| \leq \frac{1}{\epsilon^p} \int_{[0, \epsilon]^p} \|P^x - P^x G_{\mathbf{c}}^{-1}\| d\mathbf{c} \rightarrow 0, \quad \epsilon \rightarrow 0 \tag{5.22}$$

uniformly with respect to  $n \in \mathbb{N}$ . We have

$$\begin{aligned}
&\|P^x F^{-1} - P^x(F^n)^{-1}\| \\
&\leq \|P^x F^{-1} - Q_\epsilon F_\epsilon^{-1}\| + \|Q_\epsilon F_\epsilon^{-1} - Q_\epsilon(F_\epsilon^n)^{-1}\| + \|Q_\epsilon(F_\epsilon^n)^{-1} - P^x(F^n)^{-1}\|. \tag{5.23}
\end{aligned}$$

We deduce from (5.17) and (5.22) that the first and third terms in (5.23) can be chosen arbitrary small for  $\epsilon > 0$  small enough and uniformly with respect  $n$ . Note that even if we will not use this, from Lemma 2, it holds also uniformly with respect to  $x$ . Consequently, it remains to deal when  $\epsilon > 0$  is fixed with the term in the middle of the right-hand side of (5.23) when  $n \rightarrow +\infty$ . Moreover from (5.18) and its counterpart for index  $n$ , we can write:

$$\|Q_\epsilon F_\epsilon^{-1} - Q_\epsilon(F_\epsilon^n)^{-1}\| \leq \frac{1}{\epsilon^p} \int_{V(x)} \|\lambda_{[0, \epsilon]^p} \varphi_y^{-1} - \lambda_{[0, \epsilon]^p} \varphi_{n,y}^{-1}\| dP$$

with  $\varphi_{n,y}(\mathbf{c}) = (\varphi_{n,y}^1(\mathbf{c}), \dots, \varphi_{n,y}^p(\mathbf{c})) = F^n(G_{\mathbf{c}}(y))$ . Note that the domain of integration above is  $V(x)$  (and not  $\tilde{A}(\ell)$ ) because the method of superstructure is applied on  $\tilde{A}(\ell)$  with  $P_x = P_{V(x)}$ . It is enough now to show for  $P$ -almost all  $y \in V(x)$ ,

$$\lambda_{[0, \epsilon]^p} \varphi_{n,y}^{-1} \xrightarrow{\text{var}} \lambda_{[0, \epsilon]^p} \varphi_y^{-1}, \quad n \rightarrow +\infty. \tag{5.24}$$

This is done in Section 6 using Proposition 1.

## 6 Study of the conditional functionals

After some algebraic calculations, we re-express the functionals  $\varphi_{n,y}$  and  $\varphi_y$  as ordered polynomials. The conditional functional  $\varphi_y : \mathbb{R}^p \rightarrow \mathbb{R}^p$  is given by

$$\begin{aligned}\varphi_y(\mathbf{c}) &= (\varphi_{1,y}(\mathbf{c}), \dots, \varphi_{p,y}(\mathbf{c})) \\ &= F(y + c_1 l_y^1 + \dots + c_p l_y^p), \quad \mathbf{c} = (c_1, \dots, c_p).\end{aligned}$$

Moreover, we have

$$\begin{aligned}\varphi_{i,y}(\mathbf{c}) &= \varphi_i(y + \langle \mathbf{c}, l_y \rangle) \\ &= \sum_{s_1, \dots, s_{d_i}} \left( \prod_{j=1}^{d_i} (\delta_y(s_j) + \langle \mathbf{c}, \omega_y(s_j) \rangle) \right) f_i(s_1, \dots, s_{d_i})\end{aligned}$$

where  $(s_i)_i$  is the list of the jump-times of  $y \in \mathbb{D}$ . We obtain a polynomial in  $c_1, \dots, c_p$ ; we can develop it like in [5, Sec. 4.2] and finally, we have:

$$\varphi_{i,y}(\mathbf{c}) = \sum_{\substack{\mathbf{a}^i = (a_0^i, \dots, a_p^i) \\ |\mathbf{a}^i| = d_i}} B(\mathbf{a}^i, y) \mathbf{c}^{\mathbf{a}^i} \quad (6.25)$$

with, in order to simplify notations,

- $\mathbf{c}^{\mathbf{a}^i} = 1^{a_0^i} c_1^{a_1^i} \dots c_p^{a_p^i}$  for  $\mathbf{a}^i = (a_0^i, a_1^i, \dots, a_p^i)$ ;
- $B(\mathbf{a}^i, y) =$

$$\sum_{\substack{\{I_k\} \text{ partition of} \\ \{1, \dots, d_i\}, \text{ card } I_k = a_k^i}} \sum_{s_1, \dots, s_{d_i}} \left( \prod_{j \in I_0} \delta_y(s_j) \right) \left( \prod_{j \in I_1} w_y^1(s_j) \right) \dots \left( \prod_{j \in I_p} w_y^p(s_j) \right) f_i(s_1, \dots, s_{d_i}) \quad (6.26)$$

where  $(s_j)_j$  is the list of jump-time of  $y$ .

Using the polynomial expression of  $\varphi_y^i$ , the following key point is shown in [5, Sec. 4.2–4.3]. It shall be used later to apply Proposition 1 to the measure images  $\lambda_{[0,\epsilon]^p} \varphi_{n,y}^{-1}$ .

**Lemma 3** *Under hypothesis (H), for  $P$ -almost all  $y \in V(x)$ , the Jacobian  $J_y(\mathbf{c}) := \det \left( \frac{\partial \varphi_y^i}{\partial c_j}(\mathbf{c}) \right)_{1 \leq i, j \leq p}$  is non-zero for almost all  $\mathbf{c} \in (\mathbb{R}_+)^p$ .*

Since all functionals  $\varphi_{n,y}$  can be developed in the same way, we introduce also the coefficients  $B(\mathbf{a}^i, y, n)$  defined like in (6.26) with  $f_i^n$  in place of  $f_i$ , and we study the convergence of  $B(\mathbf{a}^i, y, n)$  to  $B(\mathbf{a}^i, y)$  when  $n \rightarrow +\infty$ .

In order to simplify the study of  $B(\mathbf{a}^i, y, n)$ , we begin with the preliminary simpler case of coefficients  $B(\mathbf{a}^i, x, n)$  relative to  $x \in \mathbb{D}(\varepsilon)$ .

### Study of the coefficients $B(\mathbf{a}^i, x, n)$ .

**Lemma 4** In  $\mathbb{D}(\varepsilon)$ , we have  $B(\mathbf{a}^i, x, n) \xrightarrow{P} B(\mathbf{a}^i, x)$  when  $n \rightarrow +\infty$ .

**Proof:** Note that  $\prod_{j \in I_1} w_x^1(t_j) \neq 0$  if for all  $j \in I_1$ ,  $t_j \in \Delta_k^1$  for some  $1 \leq k \leq b_1$ . Then  $w_x^1(t_j) = \pm\tau$  with the same sign as that of the jump  $\delta_x(t_j)$ . Thus  $\prod_{j \in I_1} w_x^1(t_j) = \pm\tau^{a_1^i}$  for  $A_{b_1}^{a_1^i} = b_1!/(b_1 - a_1^i)!$  choices of  $t_j$ ,  $j \in I_1$ . The same holds true for all inner products  $\prod_{j \in I_k} w_x^k(t_j) = \pm\tau^{a_k^i}$  for  $A_{b_k}^{a_k^i}$  choices of  $t_j$ ,  $j \in I_k$ , for  $1 \leq k \leq p$ . Finally, we have:

$$\left( \prod_{j \in I_1} w_x^1(t_j) \right) \cdots \left( \prod_{j \in I_p} w_x^p(t_j) \right) = \pm\tau^{a_1^i + \cdots + a_p^i}$$

for  $A(\mathbf{a}^i) := A_{b_1}^{a_1^i} \times \cdots \times A_{b_p}^{a_p^i}$  choices of index  $j$ , else the product is zero. Using the symmetry of the kernels  $f_i$  and the nullity of  $f_i$  on the diagonals, we can thus rewrite

$$\begin{aligned} & B(\mathbf{a}^i, x) \\ &= \sum_{\substack{\{I_k\} \text{ partition of} \\ \{1, \dots, d_i\}, \text{ card } I_k = a_k^i}} \sum_{\substack{A(\mathbf{a}^i) \text{ choices of} \\ t_{a_0^i+1}, \dots, t_{d_i}}} \pm\tau^{a_1^i + \cdots + a_p^i} \sum_{t_1, \dots, t_{a_0^i}} \left( \prod_{j \in I_0} \delta_x(t_j) \right) f_i(t_1, \dots, t_{d_i}). \end{aligned} \quad (6.27)$$

Observe that the outer sums in (6.27)

$$\sum_{\substack{\{I_k\} \text{ partition of} \\ \{1, \dots, d_i\}, \text{ card } I_k = a_k^i}} \quad \text{and} \quad \sum_{\substack{A(\mathbf{a}^i) \text{ choices of} \\ t_{a_0^i+1}, \dots, t_{d_i}}}$$

are both finite. Moreover, the same computations hold true for the coefficients  $B(\mathbf{a}^i, x, n)$  with  $f_i^n$  in place of  $f_i$ .

In order to study the convergence of  $B(\mathbf{a}^i, x, n)$  (with respect to  $n$ ), we first deal with the convergence of the inner sum  $\sum_{t_1, \dots, t_{a_0^i}} \left( \prod_{j \in I_0} \delta_x(t_j) \right) f_i^n(t_1, \dots, t_{d_i})$  when  $n \rightarrow +\infty$  and where  $t_{a_0^i+1}, \dots, t_{d_i}$  in  $f_i^n$  appear here as parameters.

When  $a_0^i \neq 0$ , this sum can be seen as a MSI like in (3.6). First, since for all  $1 \leq i \leq p$ ,  $f_i^n \rightarrow f_i$  in  $L^\alpha(\log_+)^{d_i-1}([0, 1]^{d_i})$ , taking some subsequence  $(n') \subset (n)$ , the convergence

$$f_i^{n'}(\cdot, t_{a_0^i+1}, \dots, t_{d_i}) \longrightarrow f(\cdot, t_{a_0^i+1}, \dots, t_{d_i}), \quad n' \rightarrow +\infty$$

holds in  $L^\alpha(\log_+)^{d_i-1}([0, 1]^{a_0^i})$  and thus also in  $L^\alpha(\log_+)^{a_0^i-1}([0, 1]^{a_0^i})$  for almost all  $t_{a_0^i+1}, \dots, t_{d_i}$ . Therefore, from Proposition 2, when  $a_0^i \neq 0$ , we have

$$I_{a_0^i}(f_i^{n'}(\cdot, t_{a_0^i+1}, \dots, t_{d_i})) \xrightarrow{\mathbb{P}} I_{a_0^i}(f(\cdot, t_{a_0^i+1}, \dots, t_{d_i})), \quad n' \rightarrow +\infty.$$

Arguing like in Proposition 3, we have when  $n' \rightarrow +\infty$ , for almost all  $t_{a_0^i+1}, \dots, t_{d_i}$

$$\sum_{t_1, \dots, t_{a_0^i}} \left( \prod_{j \in I_0} \delta_x(t_j) \right) f_i^{n'}(t_1, \dots, t_{d_i}) \xrightarrow{P} \sum_{t_1, \dots, t_{a_0^i}} \left( \prod_{j \in I_0} \delta_x(t_j) \right) f_i(t_1, \dots, t_{d_i}) \quad (6.28)$$

where we recall that  $P$  still stands for the law of the stable process  $\eta$ . Since from Lemma 1, the  $t_i$ 's can be seen as uniform and independent random variables on the  $\Delta_i$ 's, the following elementary lemma applied with  $X = t_1, \dots, t_{a_0^i}$  and  $Y = t_{a_0^i+1}, \dots, t_{d_i}$  yields the same convergence in probability than in (6.28) but with a convergence in probability involving now all the jump-times  $t_1, \dots, t_{a_0^i}, t_{a_0^i+1}, \dots, t_{d_i}$ .

**Lemma 5** *Let  $X$  and  $Y$  be independent random variables and  $f_n, f$  be some measurable functions from  $\mathbb{R}^2$  to  $\mathbb{R}$ . Suppose that for  $\mathbb{P}_Y$ -almost all  $y$ ,  $f_n(X, y) \xrightarrow{\mathbb{P}} f(X, y)$  when  $n \rightarrow +\infty$ . Then  $f_n(X, Y) \xrightarrow{\mathbb{P}} f(X, Y)$  when  $n \rightarrow +\infty$ .*

Next since the outer sums in (6.27) are both finite, we derive  $B(\mathbf{a}^i, x, n') \xrightarrow{P} B(\mathbf{a}^i, x)$  when  $n' \rightarrow +\infty$ . Thus when  $a_0^i \neq 0$ , for any subsequence  $(n') \subset (n)$ , there is some further subsequence  $(n'') \subset (n')$  such that  $B(\mathbf{a}^i, x, n'') \rightarrow B(\mathbf{a}^i, x)$  for  $P$ -almost all  $x$ .

If  $a_0^i = 0$ , the inner sum in (6.27) is empty and reduces to  $f_i^n(t_1, \dots, t_{d_i})$ . But taking eventually a subsequence, for almost all  $t_1, \dots, t_{d_i}$ , we have  $f_i^n(t_1, \dots, t_{d_i}) \rightarrow f_i(t_1, \dots, t_{d_i})$ . Since the outer sums in (6.27) are still both finite, we derive once more that for any subsequence  $(n') \subset (n)$  there is some further  $(n'') \subset (n')$  with  $B(\mathbf{a}^i, x, n'') \rightarrow B(\mathbf{a}^i, x)$ ,  $n'' \rightarrow +\infty$ , for  $P$ -almost all  $x$ .

In both cases ( $a_0^i$  is zero or not), the convergence in probability is proved.  $\square$

### Study of the coefficients $B(\mathbf{a}^i, y, n)$ .

We deal now with  $B(\mathbf{a}^i, y, n)$  for  $y \in V(x)$  and to this end, we adapt the study of the coefficients  $B(\mathbf{a}^i, x, n)$  given in the proof of Lemma 4. First, we have to study the jumps and the jump-times of  $y \in V(x)$ . Note that the (technical) choice of the parameters of local fields  $l^1, \dots, l^p$  (see around (4.12)) are required specifically for this study. This preliminary work has yet be done in [5, p. 66–67] for which we will refer for a more precise justification.

**Lemma 6 (Jumps of  $y \in V(x)$ )** *Let  $y \in V(x)$ , the neighbourhood of  $x$  defined in (4.15). The list of the jump-times of  $x$  is denoted  $(t_i)_i$  and that of  $y$  is  $(s_i)_i$ . We have*

$$T_\varepsilon(y) = (\rho^{-1}(t_1), \dots, \rho^{-1}(t_N))$$

for some increasing continuous bijection  $\rho$  of  $[0, 1]$  realizing the Skorohod distance between  $x$  and  $y$ . Moreover, the jump-times  $s_i$ ,  $i > 0$ , of  $y$  satisfy:

- $\omega_y^i(s_k) = 0$  if  $s_k \notin \cup_{j=1}^{b_i} \Delta_j^i$ ;
- $\omega_y^i(s_k) \neq 0$  if  $s_k \in \cup_{j=1}^{b_i} \Delta_j^i$  and  $s_k = \rho^{-1}(t_j^i)$ .

**Proof:** From the definition of Skorohod's topology (see [2]), let  $\rho \in \Lambda([0, 1])$ , the set of increasing continuous bijections of  $[0, 1]$ , with

$$\sup_{t \in [0, 1]} |x(\rho(t)) - y(t)| < \delta_2 \quad \text{and} \quad \sup_{t \in [0, 1]} |\rho(t) - t| < \delta_2$$

where  $\delta_2$  is given in (4.12). We have

$$\begin{aligned} \delta_x(\rho(t)) - 2\delta_2 &< \delta_y(t) < \delta_x(\rho(t)) + 2\delta_2 \\ |\delta_x(\rho(t))| - 2\delta_2 &< |\delta_y(t)| < |\delta_x(\rho(t))| + 2\delta_2. \end{aligned}$$

First  $\rho^{-1}(t_i) \in \Delta_i = (t_i - \beta, t_i + \beta)$  because  $|\rho(t_i) - t_i| < \delta_2$  and  $\delta_2 < \beta$ .  
Moreover

$$|\delta_y(\rho^{-1}(t_i))| > |\delta_x(t_i)| - 2\delta_2 \geq 2\varepsilon_0 - \frac{1}{2}\varepsilon_0 = \frac{3}{2}\varepsilon_0 > \varepsilon_0 > \varepsilon_i.$$

If  $t \in \Delta_i \setminus \{\rho^{-1}(t_i)\}$ , we have also  $\rho(t) \in \Delta'_i = (t_i - \delta_1, t_i + \delta_1)$  because  $|\rho(t) - t| < \delta_2$  and  $\beta = \delta_1 - \delta_2$ . Whence, since

- $|\delta_x(\rho(t))| \leq \varepsilon_0/2$  because  $\rho(t) \neq t_i$  and  $t_i$  is the unique time in  $\Delta'_i$  when occurs a jumps of  $x$  larger than  $\varepsilon_0/2$ ,
- $2\delta_2 < \varepsilon_1 - \varepsilon_0/2$  by choice of  $\delta_2$  in (4.12),

we have

$$|\delta_y(t)| < |\delta_x(\rho(t))| + 2\delta_2 \leq \frac{\varepsilon_0}{2} + 2\delta_2 < \varepsilon_1 \leq \varepsilon_i.$$

For  $t \in \Delta_i$ ,

- if  $t = \rho^{-1}(t_i)$  then  $t \in \Delta_i$ ,  $|\delta_y(t)| > \varepsilon_i$ ,  $\delta_y(t)$  has the same sign as  $\delta_x(t_i)$ ,
- if  $t \neq \rho^{-1}(t_i)$  then  $|\delta_y(t)| < \varepsilon_i$ .

Observe moreover that for  $t \in U_i^\varepsilon$ ,  $t \neq \rho^{-1}(t_i)$ :

$$|\delta_y(t)| \leq |\delta_x(\rho^{-1}(t))| + 2\delta_2 < |\delta_x(t'_i)| + 2\delta_2 \quad (6.29)$$

because  $\rho^{-1}(t) \neq t_i$  implies  $|\delta_x(\rho^{-1}(t))| < |\delta_x(t'_i)|$  and

$$|\delta_y(\rho^{-1}(t_i))| > |\delta_x(t_i)| - 2\delta_2. \quad (6.30)$$

From (4.12):  $\delta_2 < \frac{1}{4} \min_{i=1, \dots, N} \{|\delta_x(t_i)| - |\delta_x(t'_i)|\}$ , and from (6.29) and (6.30), we deduce:  $|\delta_y(t)| < |\delta_y(\rho^{-1}(t_i))|$ . Then we have  $\rho^{-1}(t_i) = T_{U_i^\varepsilon}(y)$  and

$$(\rho^{-1}(t_1), \dots, \rho^{-1}(t_N)) = T_\varepsilon(y).$$

This argument justifies also the second part of Lemma 6. □

**Lemma 7** *In  $V(x)$ , we have  $B(\mathbf{a}^i, y, n) \xrightarrow{P} B(\mathbf{a}^i, y)$  when  $n \rightarrow +\infty$ .*

**Proof:** We study the coefficients  $B(\mathbf{a}^i, y, n)$  just like we did for  $B(\mathbf{a}^i, x, n)$  but using moreover the study of the jump-times of  $y$  in Lemma 6. Here, from Lemma 6,  $\prod_{j \in I_1} w_y^1(s_j) = \pm \tau^{a_1^i}$  for  $A_{b_1^i}^{a_1^i}$  choices of  $s_j$ ,  $j \in I_1$ , else it is 0. Doing the same for the other products, we have

$$\left( \prod_{j \in I_1} w_x^1(s_j) \right) \cdots \left( \prod_{j \in I_p} w_x^p(s_j) \right) = \pm \tau^{a_1^i + \dots + a_p^i}$$



for  $A(\mathbf{a}^i) := A_{b_1}^{a_1^i} \times \cdots \times A_{b_p}^{a_p^i}$  choices of index  $j$ , else the product is zero. Using the symmetry of the kernels  $f_i$  and the nullity of  $f_i$  on the diagonals, we can thus rewrite

$$\begin{aligned} & B(\mathbf{a}^i, y) \\ &= \sum_{\substack{\{I_k\} \text{ partition of} \\ \{1, \dots, d_i\}, \text{ card } I_k = a_k^i}} \sum_{\substack{A(\mathbf{a}^i) \text{ choices of} \\ s_{a_0^i+1}, \dots, s_{d_i}}} \pm \tau^{a_1^i + \dots + a_p^i} \sum_{s_1, \dots, s_{a_0^i}} \left( \prod_{j \in I_0} \delta_x(s_j) \right) f_i(s_1, \dots, s_{d_i}). \end{aligned} \quad (6.31)$$

Observe again that the outer sums in (6.31)

$$\sum_{\substack{\{I_k\} \text{ partition of} \\ \{1, \dots, d_i\}, \text{ card } I_k = a_k^i}} \quad \text{and} \quad \sum_{\substack{A(\mathbf{a}^i) \text{ choices of} \\ s_{a_0^i+1}, \dots, s_{d_i}}}$$

are both finite. The same computations hold true for the coefficients  $B(\mathbf{a}^i, y, n)$  with  $f_i^n$  in place of  $f_i$ .

In order to study the convergence of the coefficients  $B(\mathbf{a}^i, y, n)$ , we first deal with the convergence of the inner sum  $\sum_{s_1, \dots, s_{a_0^i}} \left( \prod_{j \in I_0} \delta_y(s_j) \right) f_i^n(s_1, \dots, s_{d_i})$  when  $n \rightarrow +\infty$ , where  $s_{a_0^i+1}, \dots, s_{d_i}$  in  $f_i^n$  appear here as parameters.

Like in the proof of Lemma 4 for the case with  $x$ , when  $a_0^i \neq 0$ , this sum can be seen from (3.6) as a MSI and since from  $f_i^n \rightarrow f_i$  in  $L^\alpha(\log_+)^{d_i-1}([0, 1]^{d_i})$ , eventually taking some subsequence, we derive the convergence

$$f_i^n(\cdot, s_{a_0^i+1}, \dots, s_{d_i}) \longrightarrow f(\cdot, s_{a_0^i+1}, \dots, s_{d_i})$$

in  $L^\alpha(\log_+)^{d_i-1}([0, 1]^{a_0^i})$  and thus also in  $L^\alpha(\log_+)^{a_0^i-1}([0, 1]^{a_0^i})$  for almost all  $s_{a_0^i+1}, \dots, s_{d_i}$ . Thus, from Proposition 2 when  $a_0^i \neq 0$ , we have

$$I_{a_0^i}(f_i^n(\cdot, t_{a_0^i+1}, \dots, t_{d_i})) \xrightarrow{\mathbb{P}} I_{a_0^i}(f(\cdot, t_{a_0^i+1}, \dots, t_{d_i})).$$

Arguing like in Proposition 3, we rewrite when  $n \rightarrow +\infty$

$$\sum_{t_1, \dots, t_{a_0^i}} \left( \prod_{j \in I_0} \delta_x(t_j) \right) f_i^n(t_1, \dots, t_{d_i}) \xrightarrow{P} \sum_{t_1, \dots, t_{a_0^i}} \left( \prod_{j \in I_0} \delta_y(t_j) \right) f_i(t_1, \dots, t_{d_i})$$

for almost all  $s_{a_0^i+1}, \dots, s_{d_i}$ . Applying first Lemma 5 and using next the finiteness of the outer sums in (6.31), we have  $B(\mathbf{a}^i, y, n) \xrightarrow{P} B(\mathbf{a}^i, y)$  when  $n \rightarrow +\infty$  and thus for any subsequence  $(n') \subset (n)$ , there is some further subsequence  $(n'') \subset (n')$  such that  $B(\mathbf{a}^i, y, n'') \rightarrow B(\mathbf{a}^i, y)$  for  $P$ -almost all  $y \in V(x)$  in the case where  $a_0^i \neq 0$ .

If  $a_0^i = 0$ , like in the case for  $x$  in the proof of Lemma 6, the inner sum in (6.31) is empty and reduces to  $f_i^n(s_1, \dots, s_{d_i})$ . But taking eventually a subsequence, for almost all  $s_1, \dots, s_{d_i}$ , we have  $f_i^n(s_1, \dots, s_{d_i}) \rightarrow f_i(s_1, \dots, s_{d_i})$ . Since the outer sums in (6.27) are still both finite, we derive once more that for any subsequence  $(n') \subset (n)$  there is some further  $(n'') \subset (n')$  with  $B(\mathbf{a}^i, y, n'') \rightarrow B(\mathbf{a}^i, y)$  for  $P$ -almost all  $y$ .

We thus have for  $P$ -almost all  $y \in V(x)$  the convergence of the coefficients  $B(\mathbf{a}^i, y, n'')$  of  $\varphi_{n'',y}^i$  to the coefficient  $B(\mathbf{a}^i, y)$  of  $\varphi_y^i$  and the convergence in probability follows.  $\square$

Finally, we conclude now this section with the proof of (5.24). From the expression in (6.25) and from Lemma 7, we derive for any subsequence  $(n') \subset (n)$ , there is some further  $(n'') \subset (n')$  such that for  $P$ -almost all  $y \in V(x)$  we have the convergence of  $\varphi_{n'',y}$  to  $\varphi_y$  in the local Sobolev space  $W_{loc}^{p,1}(\mathbb{R}^p, \mathbb{R}^p)$ . Moreover from Lemma 3, under **(H)**, for  $P$ -almost all  $y \in V(x)$ , we have  $J_y(\mathbf{c}) \neq 0$  for almost all  $\mathbf{c}$ . We can thus apply Proposition 1 (Corollary 4 in [1]) to derive the convergence (5.24) when  $n'' \rightarrow +\infty$  that is

$$\lambda_{[0,\epsilon]^p} \varphi_{n'',y}^{-1} \xrightarrow{var} \lambda_{[0,\epsilon]^p} \varphi_y^{-1}.$$

## 7 Conclusion

For  $P$ -almost all  $y \in V(x)$ , the convergence in (5.24) has been derived for some subsequence  $(n'')$  taken from any subsequence  $(n') \subset (n)$ . Finally, returning to the term in the middle of (5.23), we derive for all  $\epsilon > 0$  and for a further subsequence  $(n'') \subset (n')$

$$\lim_{n'' \rightarrow +\infty} \|Q_\epsilon F_\epsilon^{-1} - Q_\epsilon (F_\epsilon^{n''})^{-1}\| = 0.$$

We thus have from (5.23),

$$\overline{\lim}_{n'' \rightarrow +\infty} \|P^x F^{-1} - P^x (F^{n''})^{-1}\| \leq 3\epsilon.$$

Since  $\epsilon > 0$  is arbitrary, we have

$$P^x (F^n)^{-1} \xrightarrow{var} P^x F^{-1}, \quad n \rightarrow +\infty.$$

Finally, gathering together all the steps, we have: First,

$$P^x (F^n)^{-1} \xrightarrow{var} P^x F^{-1}, \quad n \rightarrow +\infty.$$

Next by localization

$$P_{\mathbb{D}(\epsilon)} (F^n)^{-1} \xrightarrow{var} P_{\mathbb{D}(\epsilon)} F^{-1}, \quad n \rightarrow +\infty.$$

And finally by approximation

$$P (F^n)^{-1} \xrightarrow{var} P F^{-1}, \quad n \rightarrow +\infty.$$

We have proved the convergence in variation of  $\mathcal{L}(S_{d_1}(f_1^n), \dots, S_{d_p}(f_p^n))$  to  $\mathcal{L}(S_{d_1}(f_1), \dots, S_{d_p}(f_p))$ . By the Representation theorem the same holds true for the law of  $(I_{d_1}(f_1^n), \dots, I_{d_p}(f_p^n))$  to those of  $(I_{d_1}(f_1), \dots, I_{d_p}(f_p))$ . This ends the proof of Theorem 1.  $\square$

## References

- [1] D. ALEXANDROVA, V. BOGACHEV, A. PILIPENKO, *On the convergence in variation for the images of measures under differentiable mappings*, C. R. Acad. Sci. Paris, Sér. I Math., vol 328, (1999) 1055–1060.
- [2] P. BILLINGSLEY, *Convergence of Probability Measures*, Wiley, 1968.
- [3] J.-C. BRETON, *Multiple Stable Stochastic Integrals: Series Representation and Absolute Continuity of their Law*, J. Theoret. Probab., vol. 15, no. 4, (2002) 877–901.
- [4] J.-C. BRETON, *Convergence in variation of the law of stable multiple integrals*, C. R. Acad. Sci. Paris, Sér. I Math., vol. 338, (2004) 239–244.
- [5] J.-C. BRETON, *Absolute continuity of joint laws of multiple stable stochastic integrals*, J. Theoret. Probab., vol. 18, no. 1, (2005) 43–77.
- [6] J.-C. BRETON, *Convergence in variation of the laws of joint multiple Wiener-Itô integrals*. Statistic and Probability Letters. *To appear in* 2006.
- [7] Y. A. DAVYDOV, M. A. LIFSHITS, *Stratification Method in Some Probability Problems*, J. Soviet. Math. 31, no. 2, (1985) 2796–2858.
- [8] Y. A. DAVYDOV, *On Distributions of Multiple Wiener-Itô Integrals*, Theory Probab. Appl., vol. 35, no. 1, (1991) 27–37.
- [9] Y. A. DAVYDOV, M. A. LIFSHITS, N. V. SMORODINA, *Local Properties of Distributions of Stochastic Functionals*, American Mathematical Society, 1998.
- [10] W. KRAKOWIAK, J. SZULGA, *Random multilinear forms*, ANN. PROBAB., VOL. 14, NO. 3, (1986) 957–973.
- [11] J. ROSIŃSKI, G. SAMORODNITSKY, M. S. TAQQU, *Sample path properties of stochastic processes represented as multiple stable integrals*, J. MULTIVARIATE ANAL., VOL. 37, (1991) 115–134.
- [12] J. ROSIŃSKI, G. SAMORODNITSKY, *Product formula, tails and independence of multiple stable integrals*, ADVANCE IN STOCH. INEQ. (ATLANTA GA 1997), (1999) 169–194.
- [13] G. SAMORODNITSKY, J. SZULGA, *An asymptotic evaluation of the tail of a multiple symmetric  $\alpha$ -stable integral*, ANN. PROBAB., VOL. 17, (1989) 1503–1520.
- [14] G. SAMORODNITSKY, M. S. TAQQU, *Stable non-Gaussian random processes*, CHAPMAN AND HALL, 1994.
- [15] A. S. ÜSTUNEL, M. ZAKAÏ, *On independence and conditionning on Wiener space*, ANN. PROBAB., VOL 17, NO 4, (1989) 1441–1453.

## Liste des prépublications

- 99-1 Monique Jeanblanc et Nicolas Privault. A complete market model with Poisson and Brownian components. A paraître dans *Proceedings of the Seminar on Stochastic Analysis, Random Fields and Applications*, Ascona, 1999.
- 99-2 Laurence Cherfilis et Alain Miranville. Generalized Cahn-Hilliard equations with a logarithmic free energy. A paraître dans *Revista de la Real Academia de Ciencias*.
- 99-3 Jean-Jacques Prat et Nicolas Privault. Explicit stochastic analysis of Brownian motion and point measures on Riemannian manifolds. *Journal of Functional Analysis* **167** (1999) 201-242.
- 99-4 Changgui Zhang. Sur la fonction  $q$ -Gamma de Jackson. A paraître dans *Aequationes Math.*
- 99-5 Nicolas Privault. A characterization of grand canonical Gibbs measures by duality. A paraître dans *Potential Analysis*.
- 99-6 Guy Wallet. La variété des équations surstables. A paraître dans *Bulletin de la Société Mathématique de France*.
- 99-7 Nicolas Privault et Jiang-Lun Wu. Poisson stochastic integration in Hilbert spaces. *Annales Mathématiques Blaise Pascal*, **6** (1999) 41-61.
- 99-8 Augustin Fruchard et Reinhard Schäfke. Sursabilité et résonance.
- 99-9 Nicolas Privault. Connections and curvature in the Riemannian geometry of configuration spaces. *C. R. Acad. Sci. Paris, Série I* **330** (2000) 899-904.
- 99-10 Fabienne Marotte et Changgui Zhang. Multisommabilité des séries entières solutions formelles d'une équation aux  $q$ -différences linéaire analytique. A paraître dans *Annales de l'Institut Fourier*, 2000.
- 99-11 Knut Aase, Bernt Øksendal, Nicolas Privault et Jan Ubøe. White noise generalizations of the Clark-Hausmann-Ocone theorem with application to mathematical finance. *Finance and Stochastics*, **4** (2000) 465-496.
- 00-01 Eric Benoît. Canards en un point pseudo-singulier nœud. A paraître dans *Bulletin de la Société Mathématique de France*.
- 00-02 Nicolas Privault. Hypothesis testing and Skorokhod stochastic integration. *Journal of Applied Probability*, **37** (2000) 560-574.
- 00-03 Changgui Zhang. La fonction  $\theta$  de Jacobi et la sommabilité des séries entières  $q$ -Gevrey, I. *C. R. Acad. Sci. Paris, Série I* **331** (2000) 31-34.
- 00-04 Guy Wallet. Déformation topologique par changement d'échelle.
- 00-05 Nicolas Privault. Quantum stochastic calculus for the uniform measure and Boolean convolution. A paraître dans *Séminaire de Probabilités XXXV*.
- 00-06 Changgui Zhang. Sur les fonctions  $q$ -Bessel de Jackson.
- 00-07 Laure Coutin, David Nualart et Ciprian A. Tudor. Tanaka formula for the fractional Brownian motion. A paraître dans *Stochastic Processes and their Applications*.
- 00-08 Nicolas Privault. On logarithmic Sobolev inequalities for normal martingales. *Annales de la Faculté des Sciences de Toulouse* **9** (2000) 509-518.
- 01-01 Emanuelle Augeraud-Veron et Laurent Augier. Stabilizing endogenous fluctuations by fiscal policies; Global analysis on piecewise continuous dynamical systems. A paraître dans *Studies in Nonlinear Dynamics and Econometrics*
- 01-02 Delphine Boucher. About the polynomial solutions of homogeneous linear differential equations depending on parameters. A paraître dans *Proceedings of the 1999 International Symposium on Symbolic and Algebraic Computation: ISSAC 99, Sam Dooley Ed., ACM, New York 1999*.
- 01-03 Nicolas Privault. Quasi-invariance for Lévy processes under anticipating shifts.
- 01-04 Nicolas Privault. Distribution-valued iterated gradient and chaotic decompositions of Poisson jump times functionals.
- 01-05 Christian Houdré et Nicolas Privault. Deviation inequalities: an approach via covariance representations.
- 01-06 Abdallah El Hamidi. Remarques sur les sentinelles pour les systèmes distribués

- 02-01 Eric Benoît, Abdallah El Hamidi et Augustin Fruchard. On combined asymptotic expansions in singular perturbation.
- 02-02 Rachid Bebbouchi et Eric Benoît. Equations différentielles et familles bien nées de courbes planes.
- 02-03 Abdallah El Hamidi et Gennady G. Laptev. Nonexistence of solutions to systems of higher-order semilinear inequalities in cone-like domains.
- 02-04 Hassan Lakhel, Youssef Ouknine, et Ciprian A. Tudor. Besov regularity for the indefinite Skorohod integral with respect to the fractional Brownian motion: the singular case.
- 02-05 Nicolas Privault et Jean-Claude Zambrini. Markovian bridges and reversible diffusions with jumps.
- 02-06 Abdallah El Hamidi et Gennady G. Laptev. Existence and Nonexistence Results for Reaction-Diffusion Equations in Product of Cones.
- 02-07 Guy Wallet. Nonstandard generic points.
- 02-08 Gilles Bailly-Maitre. On the monodromy representation of polynomials.
- 02-09 Abdallah El Hamidi. Necessary conditions for local and global solvability of nondiagonal degenerate systems.
- 02-10 Abdallah El Hamidi et Amira Obeid. Systems of Semilinear higher order evolution inequalities on the Heisenberg group.
- 03-01 Abdallah El Hamidi et Gennady G. Laptev. Non existence de solutions d'inéquations semilinéaires dans des domaines coniques.
- 03-02 Eric Benoît et Marie-Joëlle Rochet. A continuous model of biomass size spectra governed by predation and the effects of fishing on them.
- 03-03 Catherine Stenger: On a conjecture of Wolfgang Wasow concerning the nature of turning points.
- 03-04 Christian Houdré et Nicolas Privault. Surface measures and related functional inequalities on configuration spaces.
- 03-05 Abdallah El Hamidi et Mokhtar Kirane. Nonexistence results of solutions to systems of semilinear differential inequalities on the Heisenberg group.
- 03-06 Uwe Franz, Nicolas Privault et René Schott. Non-Gaussian Malliavin calculus on real Lie algebras.
- 04-01 Abdallah El Hamidi. Multiple solutions to a nonlinear elliptic equation involving Paneitz type operators.
- 04-02 Mohamed Amara, Amira Obeid et Guy Vallet. Relaxed formulation and existence result of the degenerated elliptic small disturbance model.
- 04-03 Hippolyte d'Albis et Emmanuelle Augeraud-Veron. Competitive Growth in a Life-cycle Model: Existence and Dynamics
- 04-04 Sadjia Aït-Mokhtar: Third order differential equations with fixed critical points.
- 04-05 Mokhtar Kirane et Nasser-eddine Tatar. Asymptotic Behavior for a Reaction Diffusion System with Unbounded Coefficients.
- 04-06 Mokhtar Kirane, Eric Nabana et Stanislav I. Pohozaev. Nonexistence of Global Solutions to an Elliptic Equation with a Dynamical Boundary Condition.
- 04-07 Khaled M. Furati, Nasser-eddine Tatar and Mokhtar Kirane. Existence and asymptotic behavior for a convection Problem.
- 04-08 José Alfredo López-Mimbela et Nicolas Privault. Blow-up and stability of semilinear PDE's with gamma generator.
- 04-09 Abdallah El Hamidi. Multiple solutions with changing sign energy to a nonlinear elliptic equation.
- 04-10 Sadjia Aït-Mokhtar: A singularly perturbed Riccati equation.
- 04-11 Mohamed Amara, Amira Obeid et Guy Vallet. Weighted Sobolev spaces for a degenerated nonlinear elliptic equation.
- 04-12 Abdallah El Hamidi. Existence results to elliptic systems with nonstandard growth conditions.

- 04-13 Eric Edo et Jean-Philippe Furter: Some families of polynomial automorphisms.
- 04-14 Laurence Cherfils et Yavdat Il'yasov. On the stationary solutions of generalized reaction diffusion equations with  $p$  &  $q$ - Laplacian.
- 04-15 Jean-Christophe Breton et Youri Davydov. Local limit theorem for supremum of an empirical processes for i.i.d. random variables.
- 04-16 Jean-Christophe Breton, Christian Houdré et Nicolas Privault. Dimension free and infinite variance tail estimates on Poisson space.
- 04-17 Abdallah El Hamidi et Gennady G. Laptev. Existence and nonexistence results for higher-order semilinear evolution inequalities with critical potential.
- 05-01 Mokhtar Kirane et Nasser-eddine Tatar. Nonexistence of Solutions to a Hyperbolic Equation with a Time Fractional Damping.
- 05-02 Mokhtar Kirane et Yamina Laskri. Nonexistence of Global Solutions to a Hyperbolic Equation with a Time Fractional Damping.
- 05-03 Mokhtar Kirane, Yamina Laskri et Nasser-eddine Tatar. Critical Exponents of Fujita Type for Certain Evolution Equations and Systems with Spatio-Temporal Fractional Derivatives.
- 05-04 Abdallah El Hamidi et Jean-Michel Rakotoson. Compactness and quasilinear problems with critical exponents
- 05-05 Claudianor O. Alves et Abdallah El Hamidi. Nehari manifold and existence of positive solutions to a class of quasilinear problems.
- 05-06 Khalid Adriouch et Abdallah El Hamidi. The Nehari manifold for systems of nonlinear elliptic equations.
- 05-07 Eric Benoît. Equation fonctionnelle: Transport et convolution.
- 05-08 Jean-Philippe Furter et Stefan Maubach. Locally Finite Polynomial Endomorphisms and an extension of the Cayley-Hamilton Theorem.
- 05-09 Thomas Forget. Solutions canards en des points tournants dégénérés.
- 05-10 José Alfredo López-Mimbela et Nicolas Privault. Critical Exponents for Semilinear PDEs with Bounded Potentials.
- 06-01 Aldéric Joulin. On maximal inequalities for stable stochastic integrals.
- 06-02 Aldéric Joulin. On local Poisson-type deviation inequalities for curved continuous time Markov chains, with applications to birth-death processes.
- 06-03 Abdallah El Hamidi et Jean-Michel Rakotoson. On a pertubed anisotropic equation. A paraître dans *Ricerche di Matematica*.
- 06-04 Khalid Adriouch et Abdallah El Hamidi. On local compactness in quasilinear elliptic problems. A paraître dans *Diff. Integ. Equ.*
- 06-05 Jean-Christophe Breton. Convergence in variation of the joint laws of multiple stable stochastic integrals.