



Prépublications du Département de Mathématiques

Université de La Rochelle
Avenue Michel Crépeau
17042 La Rochelle Cedex 1
<http://www.univ-lr.fr/labo/lmca>

Fonctions minimales pour des inégalités de Sobolev anisotropiques.

A. El Hamidi et J-M Rakotoson

Septembre 2006

Classification: 35J70; 46E35; 35B65.

Mots clés: Quasilinear problems, concentration-compactness, anisotropic Sobolev inequalities.

Extremal functions for the anisotropic Sobolev inequalities.

Fonctions minimales pour des inégalités de Sobolev anisotropiques.

A. EL Hamidi(1)

(1) Laboratoire de Mathématiques, Université de La Rochelle
Av. Michel Crépeau, 17042 LA ROCHELLE Cedex 09 - France

J.M. Rakotoson(2)

(2) Laboratoire de Mathématiques - U.M.R. 6086 - Université de Poitiers - SP2MI -
Boulevard Marie et Pierre Curie, Téléport 2

BP30179 86962 FUTUROSCOPE CHASSENEUIL Cedex - France.

Keywords : Quasilinear problems, concentration-compactness, anisotropic Sobolev inequalities.

Abstract

The existence of multiple nonnegative solutions to the anisotropic critical problem

$$-\sum_{i=1}^N \frac{\partial}{\partial x_i} \left(\left| \frac{\partial u}{\partial x_i} \right|^{p_i-2} \frac{\partial u}{\partial x_i} \right) = |u|^{p^*-2} u \text{ in } \mathbb{R}^N$$

is proved in suitable anisotropic Sobolev spaces. The solutions correspond to extremal functions of a certain best Sobolev constant. The main tool in our study is an adaptation of the well-known concentration-compactness lemma of P.-L. Lions to anisotropic operators. Furthermore, we show that the set of nontrivial solutions $\mathcal{S}$ is included in $L^\infty(\mathbb{R}^N)$ and is located outside of a ball of radius $\tau > 0$ in $L^{p^*}(\mathbb{R}^N)$.

Résumé

Nous montrons l'existence d'une infinité de solutions positives pour le problème anisotrope avec exposant critique. La méthode consiste à regarder la meilleure constante d'une inégalité du type Poincaré-Sobolev et à adapter le fameux principe de concentration-compactité de P.L. Lions. De plus, on montre que l'ensemble des solutions $\mathcal{S}$ est contenu dans $L^\infty(\mathbb{R}^N)$ et est localisé en dehors d'une boule de rayon $\tau > 0$ dans $L^{p^*}(\mathbb{R}^N)$.

1 Introduction.

In this paper, the existence of nontrivial nonnegative solutions to the anisotropic critical problem

$$-\sum_{i=1}^N \frac{\partial}{\partial x_i} \left(\left| \frac{\partial u}{\partial x_i} \right|^{p_i-2} \frac{\partial u}{\partial x_i} \right) = |u|^{p^*-2} u \text{ in } \mathbb{R}^N \quad (1)$$

is studied, where the exponents p_i and p^* satisfy the following conditions

$$p_i > 1, \quad \sum_{i=1}^N \frac{1}{p_i} > 1,$$

and the critical exponent p^* is defined by

$$p^* := \frac{N}{\sum_{i=1}^N \frac{1}{p_i} - 1}.$$

In the best of our knowledge, anisotropic equations with different orders of derivation in different directions, involving critical exponents were never studied before. In the subcritical case, we can refer the reader to the recent paper by I. Fragala *et al* [4].

In the special case $p_i = 2$, $i \in \{1, 2, \dots, N\}$, Problem (1) is reduced to the limiting equation arising in the famous Yamabe problem [13]:

$$-\Delta u = u^{2^*-1}, \quad u > 0 \text{ in } \mathbb{R}^N. \quad (2)$$

Indeed, let (M, g) be a N -dimensional Riemannian manifold and S_g be the scalar curvature of the metric g . Consider a conformal metric $\tilde{g}$ on M defined by $\tilde{g} := u^{\frac{4}{N-2}} g$ whose scalar curvature (which is assumed to be constant) is denoted by $S_{\tilde{g}}$, where u is a positive function in $C^\infty(M, \mathbb{R})$. The unknown function u satisfies then

$$-\Delta_g u + \frac{N-2}{4(N-1)} S_g u = \frac{N-2}{4(N-1)} S_{\tilde{g}} u^{2^*-1}, \quad u > 0 \text{ in } M, \quad (3)$$

where Δ_g denotes the Laplace-Beltrami operator. It is clear that, up to a scaling, the limiting problem of (3) (Equation (3) without the subcritical term $\frac{N-2}{4(N-1)} S_g u$) is exactly (2). The question of existence of minimizing solutions to (2) was completely solved by Aubin [1] and G. Talenti [9]. Their proofs are based on symmetrisation theory. Notice that this theory is not relevant in our context since the radial symmetry of solutions can not hold true because of the anisotropy of the operator.

In [5], P.-L. Lions introduced the famous concentration-compactness lemma which constitutes a powerful tool for the study of critical nonlinear elliptic equations. The concentration-compactness lemma allows an elegant and simple proof of the existence of solutions to (2) by minimization arguments. In the present work, we will adapt the concentration-compactness lemma to the anisotropic case and show that the infimum

$$\inf_{|u|_{L^{p^*}(\mathbb{R}^N)}=1} \left\{ \sum_{i=1}^N \frac{1}{p_i} \left\| \frac{\partial u}{\partial x_i} \right\|_{p_i}^{p_i} \right\}$$

is achieved, of course, the functional space has to be specified.

The motivation of the present work is to give a new result which can provide extremal functions associated to the critical level corresponding to anisotropic problems involving critical exponents. Notice that the genuine extremal functions are obtained by minimization on the Nehari manifold associated to the problem and the critical level is nothing than the energy of these extremal functions.

The natural functional framework of Problem (1) is the anisotropic Sobolev spaces theory developed by [6, 11, 7, 8, 10]. Then, let $\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$ be the completion of the space $\mathcal{D}(\mathbb{R}^N)$ with respect to the norm

$$\|u\|_{1,\vec{p}} := \sum_{i=1}^N \left\| \frac{\partial u}{\partial x_i} \right\|_{p_i}.$$

It is well known that $(\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N), \|\cdot\|_{1,\vec{p}})$ is a reflexive Banach space which is continuously embedded in $L^{p^*}(\mathbb{R}^N)$.

In what follows, we will assume that

$$p_+ = \max\{p_1, p_2, \dots, p_N\} < p^*,$$

then p^* is the critical exponent associated to the operator:

$$\sum_{i=1}^N \frac{\partial}{\partial x_i} \left(\left| \frac{\partial}{\partial x_i} \right|^{p_i-2} \frac{\partial}{\partial x_i} \right).$$

The space $\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$ can also be seen as

$$\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N) = \left\{ u \in L^{p^*}(\mathbb{R}^N) : \left| \frac{\partial u}{\partial x_i} \right| \in L^{p_i}(\mathbb{R}^N) \right\}.$$

In the sequel, we will set $p_- = \min\{p_1, p_2, \dots, p_N\}$, $p_+ = \max\{p_1, p_2, \dots, p_N\}$ and $\vec{p} = (p_1, p_2, \dots, p_N)$. Also, the integral symbol $\int$ will denote $\int_{\mathbb{R}^N}$ and $\|\cdot\|_{p_i}$ will denote the usual Lebesgue norm in $L^{p_i}(\mathbb{R}^N)$. We denote by $\mathfrak{M}(\mathbb{R}^N)$ (resp. $\mathfrak{M}^+(\mathbb{R}^N)$) the space of finite measures (resp. positive finite measures) on $\mathbb{R}^N$, and by $\|\cdot\|$ its usual norm.

2 Existence of extremal functions for a Sobolev type inequality

In this paragraph, we shall prove that a certain best Sobolev constant is achieved.

Theorem 1 *Under the above assumptions on p_i , $i = 1, \dots, N$, $N \geq 2$, there exists at least one function $u \in \mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$, $u \geq 0$, $u \neq 0$:*

$$-\sum_{i=1}^N \frac{\partial}{\partial x_i} \left(\left| \frac{\partial u}{\partial x_i} \right|^{p_i-2} \frac{\partial u}{\partial x_i} \right) = u^{p^*-1} \text{ in } \mathcal{D}'(\mathbb{R}^N).$$

The proof will need two fundamental lemmas, the first one is a result due to M. Troisi [10]:

Lemma 1 (Troisi [10])

There is a constant $T_0 > 0$ depending only on $\vec{p}$ and N such that :

$$T_0 \|u\|_{p^*} \leq \prod_{j=1}^N \left\| \frac{\partial u}{\partial x_j} \right\|_{p_j}^{\frac{1}{N}} \text{ and } \|u\|_{p^*} \leq \frac{1}{NT_0} \sum_{i=1}^N \left\| \frac{\partial u}{\partial x_i} \right\|_{p_i},$$

for all $u \in \mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$.

The second lemma is a rescaling type result ensuring the conservation of suitable norms:

Lemma 2

Let $\alpha_i = \frac{p_i^*}{p_i} - 1$, $i = 1, \dots, N$. For every $y \in \mathbb{R}^N$, $u \in \mathcal{D}^{1, \vec{p}}(\mathbb{R}^N)$, and $\lambda > 0$, if we write $x = (x_1, \dots, x_N)$, $y = (y_1, \dots, y_N)$, $v(x) \doteq u^{\lambda, y}(x) = \lambda u(\lambda^{\alpha_1} x_1 + y_1, \dots, \lambda^{\alpha_N} x_N + y_N)$, we get

$$\|u\|_{p^*} = \|v\|_{p^*},$$

$$\left\| \frac{\partial u}{\partial x_i} \right\|_{p_i} = \left\| \frac{\partial v}{\partial x_i} \right\|_{p_i}, \text{ for } i = 1, \dots, N,$$

$$\text{thus, } \|u\|_{1, \vec{p}} = \|u^{\lambda, y}\|_{1, \vec{p}}.$$

Proof.

Noticing that $\sum_{i=1}^N \alpha_i = p^*$, a straightforward computation with adequate changes of variables gives the result.

Lemma 3

Let $S = \inf_{u \in \mathcal{D}^{1, \vec{p}}(\mathbb{R}^N), \|u\|_{p^*}=1} \left\{ \sum_{i=1}^N \frac{1}{p_i} \left\| \frac{\partial u}{\partial x_i} \right\|_{p_i}^{p_i} \right\}$. Then $S > 0$.

Proof.

From Lemma 1, we obtain that if $\|u\|_{p^*} = 1$, then

$$\sum_{i=1}^N \left\| \frac{\partial u}{\partial x_i} \right\|_{p_i} \geq NT_0 > 0. \quad (4)$$

Using standard argument, the infimum

$$\inf \left\{ \sum_{i=1}^N \frac{1}{p_i} a_i^{p_i}, (a_1, \dots, a_N) \in \mathbb{R}^N, \sum_{i=1}^N a_i \geq NT_0, a_i \geq 0 \right\} \doteq S_1$$

is achieved and thus this minimum is positive. By relation (4), one concludes that $S \geq S_1 > 0$. $\diamond$

Corollary 1 of Lemma 3 (Sobolev type inequality)

Let $p_- = \min(p_1, \dots, p_N)$, $p_+ = \max(p_1, \dots, p_N)$ and F be the real valued function defined

$$\text{by } F(\sigma) = \begin{cases} \sigma^{p_+} & \text{if } \sigma \leq 1, \\ \sigma^{p_-} & \text{if } \sigma \geq 1. \end{cases}$$

Then for every $u \in \mathcal{D}^{1, \vec{p}}(\mathbb{R}^N)$, one has

$$SF(\|u\|_{p^*}) \leq \sum_{i=1}^N \frac{1}{p_i} \left\| \frac{\partial u}{\partial x_i} \right\|_{p_i}^{p_i} \doteq P(\nabla u).$$

Proof.

Let u be in $\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$. If $u = 0$ the inequality is true. If $u \neq 0$, set $w = \frac{u}{\|u\|_{p^*}}$, then from the definition of S one has :

$$\sum_{i=1}^N \frac{1}{p_i} \left\| \frac{\partial w}{\partial x_i} \right\|_{p_i}^{p_i} \geq S. \quad (5)$$

Since $t^{p_i} \leq t^{p^+}$ if $t > 1$ and $t^{p_i} \leq t^{p^-}$ otherwise, the result follows from relation (5) and the definition of F . $\diamond$

Remark 1 *Along this paragraph, we only need the inequality :*

$$S \|u\|_{p^*}^{p^+} \leq P(\nabla u) \text{ whenever } \|u\|_{p^*} \leq 1.$$

We shall call $(\mathcal{P})$ the minimization problem

$$(\mathcal{P}) \quad \inf_{\|u\|_{p^*}=1} \left\{ \sum_{i=1}^N \frac{1}{p_i} \left\| \frac{\partial u}{\partial x_i} \right\|_{p_i}^{p_i} \right\} = \inf_{\|u\|_{p^*}=1} \{P(\nabla u)\}.$$

Let $(u_n) \subset \mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$ be a minimizing sequence for the problem $(\mathcal{P})$. As in [5] and Willem [12], we define the Levy concentration function:

$$Q_n(\lambda) = \sup_{y \in \mathbb{R}^N} \int_{E(y, \lambda^{\alpha_1}, \dots, \lambda^{\alpha_N})} |u_n|^{p^*} dx, \quad \lambda > 0.$$

Here $E(y, \lambda^{\alpha_1}, \dots, \lambda^{\alpha_N})$ is the ellipse defined by

$$\left\{ z = (z_1, \dots, z_N) \in \mathbb{R}^N, \sum_{i=1}^N \frac{(z_i - y_i)^2}{\lambda^{2\alpha_i}} \leq 1 \right\}$$

with $y = (y_1, \dots, y_N)$ and $\alpha_i > 0$ as in Lemma 2. Since for every n , $\lim_{\lambda \rightarrow 0} Q_n(\lambda) = 0$ and $Q_n(\lambda) \xrightarrow{\lambda \rightarrow +\infty} 1$. There exists $\lambda_n > 0$ such that $Q_n(\lambda_n) = \frac{1}{2}$. Moreover there exists $y_n \in \mathbb{R}^N$ such that

$$\int_{E(y_n, \lambda_n^{\alpha_1}, \dots, \lambda_n^{\alpha_N})} |u_n|^{p^*} dx = \frac{1}{2}.$$

Thus by a change of variables one has for $v_n = u_n^{\lambda_n, y_n}$:

$$\int_{B(0,1)} |v_n|^{p^*} dx = \frac{1}{2} = \sup_{y \in \mathbb{R}^N} \int_{B(y,1)} |v_n|^{p^*} dx.$$

Since $\|v_n\|_{p^*} = \|u_n\|_{p^*}$, $\left\| \frac{\partial v_n}{\partial x_i} \right\|_{p_i} = \left\| \frac{\partial u_n}{\partial x_i} \right\|_{p_i}$, $P(\nabla u_n) = P(\nabla v_n)$ we deduce that (v_n) is bounded in $\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$ and is also a minimizing sequence for $(\mathcal{P})$. We may then assume that :

- $v_n \rightharpoonup v$ in $\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$,

- $\left| \frac{\partial}{\partial x_i} (v_n - v) \right|^{p_i} \rightharpoonup \mu_i$ in $\mathfrak{M}^+(\mathbb{R}^N)$,
- $|v_n - v|^{p^*} \rightharpoonup \nu$ in $\mathfrak{M}^+(\mathbb{R}^N)$,
- $v_n \rightarrow v$ a.e in $\mathbb{R}^N$.

We define :

$$\begin{aligned} \mu &= \sum_{i=1}^N \frac{1}{p_i} \mu_i, \\ \mu_\infty &= \lim_{R \rightarrow +\infty} \overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int_{|x|>R} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx, \end{aligned} \quad (6)$$

$$\nu_\infty = \lim_{R \rightarrow +\infty} \overline{\lim}_n \int_{|x|>R} |v_n|^{p^*} dx. \quad (7)$$

We start with some general lemmas. First by the Brezis-Lieb's Lemma [2], direct computations give the following

Lemma 4

$$|v_n|^{p^*} \rightharpoonup |v|^{p^*} + \nu \text{ in } \mathfrak{M}^+(\mathbb{R}^N).$$

The lemma which follows gives some reverse Hölder type inequalities connecting the measures ν , μ and μ_i , $1 \leq i \leq N$.

Lemma 5

Under the above statement, one has for all $\varphi \in C_c^\infty(\mathbb{R}^N)$

$$\begin{aligned} \left(\int |\varphi|^{p^*} d\nu \right)^{\frac{1}{p^*}} &\leq \frac{1}{T_0} \prod_{i=1}^N \left(\int |\varphi|^{p_i} d\mu_i \right)^{\frac{1}{N p_i}}, \\ \left(\int |\varphi|^{p^*} d\nu \right)^{\frac{1}{p^*}} &\leq p_+^{\frac{1}{N} + \frac{1}{p^*}} \|\mu\|^{\frac{1}{N} + \frac{1}{p^*} - \frac{1}{p_+}} \cdot \frac{1}{T_0} \left(\int |\varphi|^{p_+} d\mu \right)^{\frac{1}{p_+}}. \end{aligned}$$

Proof.

Let $\varphi \in C_c^\infty(\mathbb{R}^N)$ and set $w_n = v_n - v$. Since $\int |\varphi_{x_i}|^{p_i} |w_n|^{p_i} dx \xrightarrow{n \rightarrow +\infty} 0$, we then have :

$$\lim_n \int \left| \frac{\partial}{\partial x_i} (\varphi w_n) \right|^{p_i} dx = \lim_n \int |\varphi|^{p_i} \left| \frac{\partial w_n}{\partial x_i} \right|^{p_i} dx = \int |\varphi|^{p_i} d\mu_i. \quad (8)$$

Thus from Lemma 1, it follows that

$$\left(\int |\varphi|^{p^*} d\nu \right)^{\frac{1}{p^*}} = \lim_n \left(\int |\varphi w_n|^{p^*} dx \right)^{\frac{1}{p^*}} \leq \frac{1}{T_0} \prod_{i=1}^N \left(\int |\varphi|^{p_i} d\mu_i \right)^{\frac{1}{N p_i}}. \quad (9)$$

On the other hand, since

$$\int |\varphi|^{p_i} d\mu_i \leq p_+ \int |\varphi|^{p_i} d\mu \leq p_+ \|\mu\|^{1 - \frac{p_i}{p_+}} \left(\int |\varphi|^{p_+} d\mu \right)^{\frac{p_i}{p_+}} \quad (10)$$

applying the estimates (9) and (10) and knowing that $\sum_{i=1}^N \frac{1}{p_i} = 1 + \frac{N}{p^*}$, we deduce

$$\left(\int |\varphi|^{p^*} d\nu \right)^{\frac{1}{p^*}} \leq p_+^{\frac{1}{N} + \frac{1}{p^*}} \|\mu\|^{\frac{1}{N} + \frac{1}{p^*} - \frac{1}{p_+}} \cdot \frac{1}{T_0} \left(\int |\varphi|^{p_+} d\mu \right)^{\frac{1}{p_+}}.$$

This ends the proof. $\diamond$

We then have $\|v\|_{p^*} \leq 1$. So if $\|v\|_{p^*} = 1$ then v is an extremal function since $P(\nabla v) \leq \liminf_n P(\nabla v_n) = S$ and $S \leq P(\nabla v)$. Thus, we want to show that fact, by proving that if it is not true then we have a concentration of ν at a single point and therefore $v = 0$.

Main Lemma

$$\|v\|_{p^*} = 1.$$

The remainder of this section is devoted to the proof of the main Lemma

Lemma 6

If $v \neq 0$ then

$$\lim_n \|v_n - v\|_{p^*}^{p^*} = 1 - \|v\|_{p^*}^{p^*} < 1.$$

Proof.

From Brezis-Lieb's Lemma we have :

$$\lim_n \left(\|v_n\|_{p^*}^{p^*} - \|v_n - v\|_{p^*}^{p^*} \right) = \|v\|_{p^*}^{p^*},$$

Since $\|v_n\|_{p^*} = 1$, we derive the result. $\diamond$

Lemma 7

$$S \|\nu\|^{\frac{p_+}{p^*}} \leq \|\mu\|.$$

Proof.

For large n , according to Lemma 6, we have :

$$\int |v_n - v|^{p^*} dx \leq 1.$$

Thus for all $\varphi \in C_c^\infty(\mathbb{R}^N)$, $|\varphi|_\infty \leq 1$, it holds:

$$S \left(\int |\varphi|^{p^*} |v_n - v|^{p^*} \right)^{\frac{p_+}{p^*}} \leq \sum_{i=1}^N \frac{1}{p_i} \int |\varphi|^{p_i} \left| \frac{\partial(v_n - v)}{\partial x_i} \right|^{p_i} dx + o_n(1).$$

Letting $n \rightarrow +\infty$, one gets :

$$S \left(\int |\varphi|^{p^*} d\nu \right)^{\frac{p_+}{p^*}} \leq \sum_{i=1}^N \frac{1}{p_i} \int |\varphi|^{p_i} d\mu_i \leq \|\mu\|. \quad (11)$$

Using the density of $C_c^\infty(\mathbb{R}^N)$ in $C_c(\mathbb{R}^N)$, we get then

$$S \left(\sup_{\varphi \in C_c(\mathbb{R}^N), |\varphi|_\infty = 1} \int |\varphi|^{p^*} d\nu \right)^{\frac{p_+}{p^*}} \leq \|\mu\|,$$

that is the desired result. $\diamond$

Lemma 8 *Let ψ_R be in $C^1(\mathbb{R})$, $0 \leq \psi_R \leq 1$, $\psi_R = 1$ if $|x| > R + 1$, $\psi_R(x) = 0$ if $|x| < R$. Then for any $\gamma_i > 0$, $i = 0, \dots, N$, the two equalities*

$$\begin{aligned}\nu_\infty &= \lim_{R \rightarrow +\infty} \overline{\lim}_n \int |v_n|^{p^*} \psi_R^{\gamma_0} dx, \\ \mu_\infty &= \lim_{R \rightarrow +\infty} \overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} \psi_R^{\gamma_i} dx.\end{aligned}$$

hold true, where ν_∞ and μ_∞ are defined by (6), (7).

Proof.

As in Willem [12], one has :

$$\begin{aligned}\int_{|x| > R+1} |v_n|^{p^*} dx &\leq \int |v_n|^{p^*} \psi_R^{\gamma_0} dx \leq \int_{|x| > R} |v_n|^{p^*} dx, \\ \int_{|x| > R+1} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx &\leq \int \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} \psi_R^{\gamma_i} \leq \int_{|x| > R} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx.\end{aligned}$$

We conclude with the definition of ν_∞ and μ_∞ . ◇

Lemma 9

Let $w_n = v_n - v$. Then, for any $\gamma_i > 0$, $i = 0, \dots, N$, we get

$$\nu_\infty = \lim_{R \rightarrow \infty} \overline{\lim}_n \int |w_n|^{p^*} \psi_R^{\gamma_0} dx,$$

and

$$\mu_\infty = \lim_{R \rightarrow \infty} \overline{\lim}_n \int \left| \frac{\partial w_n}{\partial x_i} \right|^{p_i} \psi_R^{\gamma_i} dx.$$

Proof.

Since

$$\lim_{R \rightarrow +\infty} \int |v|^{p^*} \psi_R^{\gamma_0} = \lim_{R \rightarrow +\infty} \int \left| \frac{\partial v}{\partial x_i} \right|^{p_i} \psi_R^{\gamma_i} dx = 0.$$

Thus

$$\lim_{R \rightarrow \infty} \overline{\lim}_n \int |w_n|^{p^*} \psi_R^{\gamma_0} dx = \lim_{R \rightarrow \infty} \overline{\lim}_n \int |v_n|^{p^*} \psi_R^{\gamma_0} dx = \nu_\infty$$

and

$$\lim_{R \rightarrow \infty} \overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial w_n}{\partial x_i} \right|^{p_i} \psi_R^{\gamma_i} dx = \lim_{R \rightarrow \infty} \overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} \psi_R^{\gamma_i} dx.$$

◇

Lemma 10

$$S\nu_\infty^{\frac{p_+}{p^*}} \leq \mu_\infty.$$

Proof.

From Lemma 6, we know that for n large enough, we have

$$\int \psi_R^{p^*} |w_n|^{p^*} \leq \int |w_n|^{p^*} dx \leq 1.$$

Thus by Sobolev inequality (Corollary 1 of Lemma 3), it follows

$$\begin{aligned} S \left(\int |\psi_R w_n|^{p^*} dx \right)^{\frac{p_+}{p^*}} &\leq \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial}{\partial x_i} (\psi_R w_n) \right|^{p_i}, \\ S \left(\lim_{R \rightarrow +\infty} \overline{\lim}_n \int |\psi_R w_n|^{p^*} dx \right)^{\frac{p_+}{p^*}} &\leq \lim_{R \rightarrow +\infty} \overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial}{\partial x_i} (\psi_R w_n) \right|^{p_i}. \end{aligned} \quad (12)$$

Since

$$\lim_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial \psi_R}{\partial x_i} \right|^{p_i} |w_n|^{p_i} = 0,$$

then

$$\lim_{R \rightarrow +\infty} \overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial}{\partial x_i} (\psi_R w_n) \right|^{p_i} = \lim_{R \rightarrow +\infty} \overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial w_n}{\partial x_i} \right|^{p_i} \psi_R^{p_i} = \mu_\infty.$$

relation (12) and Lemma 9 give :

$$S \nu_\infty^{\frac{p_+}{p^*}} \leq \mu_\infty.$$

◇

Following again the arguments used in [12] we claim that:

Lemma 11

$$1 = \lim_n \|v_n\|_{p^*}^{p^*} = \|v\|_{p^*}^{p^*} + \|\nu\| + \nu_\infty.$$

Proof.

From Lemma 4, we have :

$$|v_n|^{p^*} \rightharpoonup |v|^{p^*} + \nu.$$

Thus

$$\lim_{R \rightarrow +\infty} \lim_n \int (1 - \psi_R^{p^*}) |v_n|^{p^*} dx = \int |v|^{p^*} dx + \int d\nu.$$

Rewriting $\|v_n\|_{p^*}^{p^*}$ as

$$\|v_n\|_{p^*}^{p^*} = \int (1 - \psi_R^{p^*}) |v_n|^{p^*} + \int \psi_R^{p^*} |v_n|^{p^*},$$

we obtain

$$\begin{aligned} \lim_n \|v_n\|_{p^*}^{p^*} &= \lim_{R \rightarrow +\infty} \lim_n \int (1 - \psi_R^{p^*}) |v_n|^{p^*} + \lim_{R \rightarrow +\infty} \overline{\lim}_n \int \psi_R^{p^*} |v_n|^{p^*} \\ &= \|v\|_{p^*}^{p^*} + \|\nu\| + \nu_\infty \end{aligned}$$

◇

Next, we shall prove the following corollary:

Corollary 1 (of Lemma 5)

There exists an at most countable index set J of distinct points $\{x_j\}_{j \in J} \subset \mathbb{R}^N$ and non-negative weights a_j and b_j , $j \in J$ such that :

$$1. \quad \nu = \sum_{j \in J} a_j \delta_{x_j}.$$

$$2. \quad \mu \geq \sum_{j \in J} b_j \delta_{x_j}.$$

$$3. \quad S a_j^{\frac{p_+}{p^*}} \leq b_j, \quad \forall j \in J.$$

Proof.

The proof follows essentially the concentration compactness principle of P.L. Lions [5] because we have the reverse Hölder type inequalities of Lemma 5.

Indeed, the second statement of this lemma implies that for all borelian sets $E \subset \mathbb{R}^N$, one has:

$$\nu(E) \leq c_\mu \mu(E)^{\frac{p^*}{p_+}}. \quad (13)$$

Since the set $D = \{x \in \mathbb{R}^N : \mu(\{x\}) > 0\}$ is at most countable because $\mu \in \mathfrak{M}(\mathbb{R}^N)$, therefore $D = \{x_j, j \in J\}$ and $b_j \doteq \mu(\{x_j\})$ satisfies $\mu \geq \sum_{j \in J} b_j \delta_{x_j}$.

Relation (13) implies that ν is absolutely continuous with respect to μ , i.e., $\nu \ll \mu$ and

$$\frac{\nu(B(x, r))}{\mu(B(x, r))} \leq c_\mu \mu(B(x, r))^{\frac{p^*}{p_+} - 1},$$

provided that $\mu(B(x, r)) \neq 0$ (remember that $p^* > p_+$). Thus, we have :

$$\nu(E) = \int_E \lim_{r \rightarrow 0} \frac{\nu(B(x, r))}{\mu(B(x, r))} d\mu(x),$$

and

$$D_\mu \nu(x) = \lim_{r \rightarrow 0} \frac{\nu(B(x, r))}{\mu(B(x, r))} = 0, \quad \mu \text{ a.e. on } \mathbb{R}^N \setminus D.$$

Setting $a_j = D_\mu \nu(x_j) b_j$, relation (13) implies that ν has only atoms that are given by $\{x_j\}$, that we have already get.

Let $\varphi \in C_c^\infty(\mathbb{R}^N)$, $\varphi(x_j) = 1$, $\|\varphi\|_\infty = 1$. Then, using statement 1. of this corollary and relation (11), we have

$$S a_j^{\frac{p_+}{p^*}} \leq S \left(\int |\varphi|^{p^*} d\nu \right)^{\frac{p_+}{p^*}} \leq \sum_{i=1}^N \frac{1}{p_i} \int |\varphi|^{p_i} d\mu_i. \quad (14)$$

We shall consider $\phi \in C_c^\infty(\mathbb{R}^N)$, $0 \leq \phi \leq 1$, $\text{support}(\phi) \subset B(0, 1)$, $\phi(0) = 1$. We fix $j \in J$ and set $x_j = (x_{j,1}, \dots, x_{j,N})$, $q_i = \frac{p_i p^*}{p^* - p_i}$, $i = 1, \dots, N$. Then $\alpha_i \doteq \frac{1}{q_i}$ satisfy

$\sum_{k=1}^N \alpha_k - \alpha_i q_i = 0$. For $\varepsilon > 0$, we define, for every $z \in \mathbb{R}^N$, $z = (z_1, \dots, z_N)$:

$$\phi_\varepsilon(z) = \phi \left(\frac{z_1 - x_{j,1}}{\varepsilon^{\alpha_1}}, \dots, \frac{z_N - x_{j,N}}{\varepsilon^{\alpha_N}} \right). \quad (15)$$

Thus we have :

$$\int \left| \frac{\partial \phi_\varepsilon}{\partial x_i} \right|^{q_i} = \int \left| \frac{\partial \phi}{\partial x_i} \right|^{q_i} (z) dz \quad (16)$$

and then

$$\int \left| \frac{\partial \phi_\varepsilon}{\partial x_i} \right|^{p_i} |v|^{p_i} \leq \left(\int \left| \frac{\partial \phi}{\partial x_i} \right|^{q_i} dz \right)^{1 - \frac{p_i}{p^*}} \left(\int_{B(x_j, \max_i \varepsilon^{\frac{1}{q_i}})} |v|^{p^*} dz \right)^{\frac{p_i}{p^*}} \xrightarrow{\varepsilon \rightarrow 0} 0. \quad (17)$$

Lemma 12 *Let $x_j \in D$ and ϕ_ε be the function defined above associated to x_j . Then :*

$$Sa_j^{\frac{p_+}{p^*}} \leq \lim_{\varepsilon \rightarrow 0} \lim_n \sum_{i=1}^N \frac{1}{p_i} \int \phi_\varepsilon^{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx.$$

Proof.

Since $0 \leq \phi_\varepsilon \leq 1$ then $\int \phi_\varepsilon^{p^*} |v_n|^{p^*} dx \leq 1$. From Corollary 1 of Lemma 3, it follows

$$S \left(\int \phi_\varepsilon^{p^*} |v_n|^{p^*} dx \right)^{\frac{p_+}{p^*}} \leq \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial}{\partial x_i} (\phi_\varepsilon v_n) \right|^{p_i}. \quad (18)$$

From relation (17), we have

$$\lim_{\varepsilon \rightarrow 0} \int \left| \frac{\partial \phi_\varepsilon}{\partial x_i} \right|^{p_i} |v|^{p_i} dx = 0. \quad (19)$$

Since

$$\lim_{n \rightarrow +\infty} \int \left| \frac{\partial \phi_\varepsilon}{\partial x_i} \right|^{p_i} |v_n - v|^{p_i} dx = 0, \quad (20)$$

then one has :

$$\lim_{\varepsilon \rightarrow 0} \lim_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial}{\partial x_i} (\phi_\varepsilon v_n) \right|^{p_i} dx = \lim_{\varepsilon \rightarrow 0} \lim_n \sum_{i=1}^N \frac{1}{p_i} \int \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} \phi_\varepsilon^{p_i} dx \quad (21)$$

From relations (18) and (21), knowing that $|v_n|^{p^*} \rightharpoonup |v|^{p^*} + \nu$ (see Lemma 4), we obtain

$$Sa_j^{\frac{p_+}{p^*}} \leq \lim_{\varepsilon \rightarrow 0} \lim_n \sum_{i=1}^N \frac{1}{p_i} \int \phi_\varepsilon^{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx.$$

◇

Lemma 13

Assume that $\sum_{i=1}^N \frac{1}{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} \rightharpoonup \tilde{\mu}$ in $\mathfrak{M}^+(\mathbb{R}^N)$. Then

1. For all $j \in J$, $Sa_j^{\frac{p_+}{p^*}} \leq \lim_{\varepsilon \rightarrow 0} \tilde{\mu}(\text{support } \phi_\varepsilon)$
(one has $\text{support } \phi_\varepsilon \subset B(x_j, \max_i \varepsilon^{\frac{1}{q_i}})$).
2. $\|\tilde{\mu}\| \geq S \|\nu\|^{\frac{p_+}{p^*}} + P(\nabla v)$.
3. $S = \lim_{n \rightarrow +\infty} P(\nabla v_n) \doteq \|\tilde{\mu}\| + \mu_\infty \geq P(\nabla v) + S \|\nu\|^{\frac{p_+}{p^*}} + \mu_\infty$.

Proof.

From Lemma 12, since $\phi_\varepsilon^{p_i} \leq \phi_\varepsilon$ and

$$\overline{\lim}_n \sum_{i=1}^N \frac{1}{p_i} \int \phi_\varepsilon^{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx \leq \int \phi_\varepsilon d\tilde{\mu},$$

one obtains

$$Sa_j^{\frac{p_+}{p^*}} \leq \overline{\lim}_{\varepsilon \rightarrow 0} \int \phi_\varepsilon d\tilde{\mu} \leq \lim_{\varepsilon \rightarrow 0} \tilde{\mu} \left(B(x_j; \max_{1 \leq i \leq N} \varepsilon^{\frac{1}{q_i}}) \right). \quad (22)$$

This shows that $\{x_j\}_{j \in J}$ are all atomic points of $\tilde{\mu}$ and since $\sum_{i=1}^N \frac{1}{p_i} \left| \frac{\partial v}{\partial x_i} \right|^{p_i}$ is orthogonal to the atomic part of $\tilde{\mu}$, one deduces from relation (22) that

$$\tilde{\mu} \geq S \sum_{j \in J} a_j^{\frac{p_+}{p^*}} \delta_{x_j} + \sum_{i=1}^N \frac{1}{p_i} \left| \frac{\partial v}{\partial x_i} \right|^{p_i}. \quad (23)$$

This implies in particular that :

$$\|\tilde{\mu}\| \geq S \sum_{j \in J} a_j^{\frac{p_+}{p^*}} + P(\nabla v). \quad (24)$$

Since $\frac{p_+}{p^*} < 1$ one has

$$\left(\sum_{j \in J} a_j \right)^{\frac{p_+}{p^*}} \leq \sum_{j \in J} a_j^{\frac{p_+}{p^*}}. \quad (25)$$

As $\nu = \sum_{j \in J} a_j \delta_{x_j}$, it holds

$$\|\nu\| = \sum_{j \in J} a_j, \quad (26)$$

which means, combining relations (24) to (26), that :

$$\|\tilde{\mu}\| \geq S \|\nu\|^{\frac{p_+}{p^*}} + P(\nabla v).$$

For the last statement, we argue as before:

$$\begin{aligned}
S &= \lim_n P(\nabla v_n) \\
&= \lim_{R \rightarrow +\infty} \lim_n \int_{\mathbb{R}^N} (1 - \psi_R) \sum_{i=1}^N \frac{1}{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx \\
&\quad + \lim_{R \rightarrow +\infty} \overline{\lim}_n \int \psi_R \sum_{i=1}^N \frac{1}{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx,
\end{aligned}$$

where $\psi_R = 1$ on $|x| > R + 1$, $0 \leq \psi_R \leq 1$, $\psi_R = 0$ if $|x| < R$, $\psi_R \in C(\mathbb{R})$. By the definition of $\tilde{\mu}$, one has :

$$\lim_{R \rightarrow +\infty} \lim_n \int (1 - \psi_R) \sum_{i=1}^N \frac{1}{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx = \lim_R \int (1 - \psi_R) d\tilde{\mu} = \|\tilde{\mu}\|,$$

and (see Lemma 8):

$$\lim_{R \rightarrow +\infty} \overline{\lim}_n \int \psi_R \sum_{i=1}^N \frac{1}{p_i} \left| \frac{\partial v_n}{\partial x_i} \right|^{p_i} dx = \mu_\infty,$$

thus, by the preceding statements:

$$S = \|\tilde{\mu}\| + \mu_\infty \geq P(\nabla v) + S \|\nu\|^{\frac{p_+}{p^*}} + \mu_\infty.$$

◇

Lemma 14

If $\|\nu\|_{p^*} < 1$ then $\|\nu\| = 1$, $\nu_\infty = 0$ and $v = 0$.

Proof.

From Lemma 10, we know that

$$S \nu_\infty^{\frac{p_+}{p^*}} \leq \mu_\infty.$$

And by Corollary 1 of Lemma 3, we have

$$S \|\nu\|_{p^*}^{p_+} \leq P(\nabla v).$$

From the last statement of Lemma 13 and the above inequalities we deduce that :

$$S \geq S \left((\|\nu\|_{p^*}^{p_+})^{\frac{p_+}{p^*}} + \|\nu\|^{\frac{p_+}{p^*}} + \nu_\infty^{\frac{p_+}{p^*}} \right).$$

Thus we obtain, due to Lemma 11, that

$$\left((\|\nu\|_{p^*}^{p_+})^{\frac{p_+}{p^*}} + \|\nu\|^{\frac{p_+}{p^*}} + \nu_\infty^{\frac{p_+}{p^*}} \right) \leq 1 = \left(\|\nu\|_{p^*}^{p_+} + \|\nu\| + \nu_\infty \right)^{\frac{p_+}{p^*}}.$$

Using the inequality

$$\left(\|\nu\|_{p^*}^{p_+} + \|\nu\| + \nu_\infty \right)^{\frac{p_+}{p^*}} \leq \|\nu\|_{p^*}^{\frac{p_+}{p^*}} + \|\nu\|^{\frac{p_+}{p^*}} + \nu_\infty^{\frac{p_+}{p^*}},$$

we get

$$\|v\|_{p^*}^{\frac{p_+}{p^*}} + \|\nu\|_{p^*}^{\frac{p_+}{p^*}} + \nu_\infty^{\frac{p_+}{p^*}} = \left(\|v\|_{p^*}^{p^*} + \|\nu\| + \nu_\infty \right)^{\frac{p_+}{p^*}}.$$

It follows that $\|v\|_{p^*}^{p^*}$, $\|\nu\|$ and ν_∞ are equal either to 0 or to 1. But using the fact that $\nu_\infty \leq \frac{1}{2}$, since $\int_{B(0,1)} |v_n|^{p^*} dx = \frac{1}{2}$, we conclude that $\nu_\infty = 0$, $\|v\|_{p^*}^{p^*} < 1$ (by our assumption) so that $v = 0$ and thus $\|\nu\| = 1$. $\diamond$

Lemma 15

If $\|v\|_{p^} < 1$ then the measure ν is concentrated at a single point $z = x_{i_0}$.*

Proof.

Since

$$S = \|\tilde{\mu}\| + \mu_\infty \geq S \sum_{j \in J} a_j^{\frac{p_+}{p^*}},$$

(see relation(24)) and $1 = \|\nu\| = \sum_{j \in J} a_j$, we then have :

$$\left(\sum_{j \in J} a_j \right)^{\frac{p_+}{p^*}} \geq \sum_{j \in J} a_j^{\frac{p_+}{p^*}} \geq \left(\sum_{j \in J} a_j \right)^{\frac{p_+}{p^*}}.$$

Thus the a_j are equal either to zero or to 1 that is, there is only one index i_0 such that $a_{i_0} = 1$ and $a_j = 0$ for $j \neq i_0$: $\nu = a_{i_0} \delta_{x_{i_0}}$. $\diamond$

End of the proof of the main Lemma :

If $\|v\|_{p^*} < 1$ thus ν concentrates at x_{i_0} and $\|\nu\| = 1$. On the other hand we have $\frac{1}{2} = \sup_{y \in \mathbb{R}^N} \int_{B(y,1)} |v_n|^{p^*} \geq \int_{B(x_{i_0},1)} |v_n|^{p^*} dx \rightarrow \|\nu\| = 1$, which is impossible, we conclude then that $\|v\|_{p^*} = 1$. $\diamond$

Consequently, the function v is a (non trivial) extremal function that can be chosen non-negative (replacing v by $|v|$).

End of the proof of Theorem 1 :

From usual Lagrange multiplier rule, there is $\lambda_0 > 0$, such that :

$$-\sum_{i=1}^N \frac{\partial}{\partial x_i} \left(\left| \frac{\partial v}{\partial x_i} \right|^{p_i-2} \frac{\partial v}{\partial x_i} \right) = \lambda_0 v^{p^*-1} \text{ in } \mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)'.$$

A similar rescaling argument used above (say $v(\lambda_0^{-\frac{1}{p_1}} x_1, \dots, \lambda_0^{-\frac{1}{p_N}} x_N)$) gives the result. $\diamond$

The multiplicity of solutions comes directly from Lemma 2, that is :

Lemma 16 :

Let $\alpha \in \mathbb{R}$, $\alpha_i = \alpha \frac{p^*}{p_i} - \alpha$, $i = 1, \dots, N$ and $u \in \mathcal{S}$. Then, for all $\lambda \in \mathbb{R}_+^*$ for all $z = (z_1, \dots, z_N) \in \mathbb{R}^N$, the function defined by

$$u^{\lambda, z}(x) = \lambda^\alpha u(\lambda^{\alpha_1} x_1 + z_1, \dots, \lambda^{\alpha_N} x_N + z_N),$$

with $x = (x_1, \dots, x_N)$ belongs to $\mathcal{S}$.

Proof.

It is the same as for Lemma 2 using a direct computation.

3 Some properties of the solutions of (1)

We want to show first the :

Proposition 1

Any nonnegative solution u being in $\mathcal{D}^{1, \vec{p}}(\mathbb{R}^N)$ of (1) belongs to $L^q(\mathbb{R}^N)$ for all $p^* \leq q < +\infty$.

Proof.

We follow the proof of [4]. Let $a > 0$. Let j be fixed in $\{1, \dots, N\}$, for $L > 0$ (large) we define $\varphi_{j,L} = u \min[u^{ap_j}, L^{p_j}] \in \mathcal{D}^{1, \vec{p}}(\mathbb{R}^N)$ and for all i

$$|\partial_i u|^{p_j-2} \partial_i u \partial_i \varphi_{j,L} \geq \min[u^{ap_j}, L^{p_j}] |\partial_i u|^{p_j} \quad a.e., \quad (27)$$

and

$$|\partial_i(u \cdot \min[u^a, L])|^{p_j} \leq (a+1)^{p_j} \min[u^{ap_j}, L^{p_j}] |\partial_i u|^{p_j} \quad a.e. \quad (28)$$

Choosing $\varphi_{j,L}$ as a test function, one has :

$$\begin{aligned} \int_{\mathbb{R}^N} \min[u^{ap_j}, L^{p_j}] |\partial_j u|^{p_j} dx &\leq \sum_{i=1}^N \int_{\mathbb{R}^N} |\partial_i u|^{p_i-2} \partial_i u \partial_i \varphi_{j,L} dx \\ &= \int_{\mathbb{R}^N} u^{p^*} \min[u^{ap_j}, L^{p_j}] dx. \end{aligned} \quad (29)$$

Introducing $k > 0$, one has :

$$\int_{\mathbb{R}^N} u^{p^*} \min[u^{ap_j}, L^{p_j}] dx \leq k^{ap_j} \int_{\mathbb{R}^N} u^{p^*} dx + \int_{u \geq k} u^{p^*} \min[u^{ap_j}, L^{p_j}] dx. \quad (30)$$

Writing that :

$$\int_{u \geq k} u^{p^*} \min[u^{ap_j}, L^{p_j}] dx = \int_{u \geq k} u^{p^*-p_j} u^{p_j} (\min[u^a, L])^{p_j} dx. \quad (31)$$

The Hölder inequality applied to the right hand side of relation (31) shows that :

$$\int_{u \geq k} u^{p^*} \min[u^{ap_j}, L^{p_j}] dx \leq \left(\int_{u \geq k} u^{p^*} dx \right)^{1-\frac{p_j}{p^*}} \left(\int_{\mathbb{R}^N} (u \min[u^a, L])^{p^*} dx \right)^{\frac{p_j}{p^*}}. \quad (32)$$

By the Troisi's inequality (see Lemma 1)

$$\left(\int_{\mathbb{R}^N} (u \min[u^a, L])^{p^*} \right)^{\frac{1}{p^*}} \leq c \sum_{i=1}^N \left(\int_{\mathbb{R}^N} |\partial_i (u \min[u^a, L])|^{p_i} \right)^{\frac{1}{p_i}} \quad (33)$$

Setting $I_i = \left(\int |\partial_i (u \min[u^a, L])|^{p_i} \right)^{\frac{1}{p_i}}$, $\varepsilon_k = \int_{u \geq k} u^{p^*} dx$, relations (28) to (33), lead to :

$$\begin{aligned} \int |\partial_j (u \cdot \min[u^a, L])|^{p_j} dx &\leq (a+1)^{p_j} \int \min[u^{ap_j}, L^{p_j}] |\partial_j u|^{p_j} dx \\ &\leq (a+1)^{p_j} k^{ap_j} \left(\int u^{p^*} dx \right) \\ &\quad + c(a+1)^{p_j} \varepsilon_k^{1-\frac{p_j}{p^*}} \left[\sum_{i=1}^N \left(\int |\partial_i (u \min[u^a, L])|^{p_i} \right)^{\frac{1}{p_i}} \right]^{p_j}. \end{aligned}$$

Thus, for all j :

$$I_j \leq (a+1)k^a \left(\int u^{p^*} dx \right)^{\frac{1}{p_j}} + c(a+1) \varepsilon_k^{\frac{1}{p_j} - \frac{1}{p^*}} \left(\sum_{i=1}^N I_i \right) \quad (34)$$

The relation(34) infers :

$$\sum_{j=1}^N I_j \leq (a+1)k^a \left(\sum_{j=1}^N \|u\|_{p^*}^{\frac{p_j}{p^*}} \right) + c(a+1) \left(\sum_{j=1}^N \varepsilon_k^{\frac{1}{p_j} - \frac{1}{p^*}} \right) \left(\sum_{i=1}^N I_i \right). \quad (35)$$

Since $\lim_{k \rightarrow +\infty} \sum_{j=1}^N \varepsilon_k^{\frac{1}{p_j} - \frac{1}{p^*}} = 0$, there exists $k_a > 0$ such that for all $k \geq k_a$, such that

$c(a+1) \sum_{j=1}^N \varepsilon_k^{\frac{1}{p_j} - \frac{1}{p^*}} \leq \frac{1}{2}$. Thus relation (35) infers then

$$\sum_{i=1}^N I_i \leq 2(a+1)k^a \sum_{j=1}^N \|u\|_{p^*}^{\frac{p_j}{p^*}}, \text{ for } k \geq k_a.$$

By the Troisi's inequality, one has :

$$\|u \cdot \min[u^a, L]\|_{L^{p^*}} \leq c \sum_{j=1}^N I_j \leq 2c(a+1)k^a \sum_{j=1}^N \|u\|_{p^*}^{\frac{p_j}{p^*}}.$$

Letting $L \rightarrow +\infty$, one has :

$$\|u^{a+1}\|_{L^{p^*}} \leq 2c(a+1)k^a \sum_{j=1}^N \|u\|_{p^*}^{\frac{p_j}{p^*}}.$$

Let $q = (a+1)p^*$, then we obtain the result. $\diamond$

Proposition 2 Any nonnegative solution u being in $\mathcal{D}^{1,\vec{p}}(\mathbb{R}^N)$ of (1) belongs to $L^\infty(\mathbb{R}^N)$. Moreover, there exists a number τ_0 depending only on p_j, N such that

$$\|u\|_{p^*} \geq \tau_0 > 0, \text{ for } u \text{ non trivial.}$$

Proof.

For $u \geq 0$ solution of (1), we set $A_\tau = \{x \in \mathbb{R}^N, u(x) \geq \tau\}$ and $|A_\tau|$ its Lebesgue measure. Since $p^* > p_+$, one can choose $q > p^*$ so that

$$\varepsilon \doteq -\frac{1}{p^*} + \left(1 - \frac{p^*}{q}\right) \left(1 - \frac{1}{p^*}\right) \frac{1}{p_+ - 1} > 0.$$

Let $\varphi_k = (u - k)_+$, for $k > 0$ fixed. Chosing this function as a test function and using proposition 1, one has :

$$\sum_{i=1}^N \left\| \frac{\partial \varphi_k}{\partial x_i} \right\|_{p_i}^{p_i} = \int u^{p^*-1} (u - k)_+ \leq c_1 |A_k|^{(1-\frac{p^*}{q})(1-\frac{1}{p^*})} \|\varphi_k\|_{p^*}, \quad (36)$$

with $c_1 = \|u\|_q^{p^*-1}$.

Since $\|\varphi_k\|_{p^*} \leq \|u\|_{p^*}$, thus the corollary 1 of Lemma 3 and relation (36) imply :

$$\|\varphi_k\|_{p^*}^{p_+} \leq c_2 \sum_{i=1}^N \left\| \frac{\partial \varphi_k}{\partial x_i} \right\|_{p_i}^{p_i} \leq c_3 |A_k|^{(1-\frac{p^*}{q})(1-\frac{1}{p^*})} \|\varphi_k\|_{p^*}, \quad (37)$$

with $c_2 = \frac{1}{S \cdot p_-} \max_{1 \leq j \leq N} (\|u\|_{p^*}^{p_+ - p_j})$, $c_3 = c_1 c_2$.

Thus,

$$\|\varphi_k\|_{p^*} \leq c_4 |A_k|^{\frac{1}{p_+ - 1} (1 - \frac{p^*}{q})(1 - \frac{1}{p^*})}. \quad (38)$$

with $c_4 = c_3^{\frac{1}{p_+ - 1}}$. By Cavalieri's principle, Hölder inequality and relation(38), one has, for all $k > 0$:

$$\int_k^{+\infty} |A_\tau| d\tau = \int_{\mathbb{R}^N} (u - k)_+(x) dx \leq |A_k|^{1 - \frac{1}{p^*}} \|\varphi_k\|_{p^*} \leq c_4 |A_k|^{1 + \varepsilon}. \quad (39)$$

This last relation is a Gronwall inequality, which shows that $\forall k > 0$

$$\|u\|_\infty \leq k + \frac{1 + \varepsilon}{\varepsilon} \|(u - k)_+\|_1^{\frac{\varepsilon}{1 + \varepsilon}} c_4^{\frac{1}{1 + \varepsilon}}. \quad (40)$$

Setting

$$\gamma = (p^* - 1) \frac{\varepsilon}{1 + \varepsilon}, \quad b_0 = \frac{1 + \varepsilon}{\varepsilon} \|u\|_{p^*}^{\frac{\varepsilon p^*}{1 + \varepsilon}} c_4^{\frac{1}{1 + \varepsilon}},$$

and noticing that

$$\|(u - k)_+\|_1 \leq \frac{\|u\|_{p^*}^{p^*}}{k^{p^* - 1}},$$

thus relation(40) becomes :

$$\|u\|_\infty \leq \inf_{k > 0} \left[k + \frac{b_0}{k^\gamma} \right] = (\gamma + 1) \gamma^{-\frac{\gamma}{\gamma + 1}} b_0^{\frac{1}{1 + \gamma}}. \quad (41)$$

Separating the contribution of $\|u\|_q$ and $\|u\|_{p^*}$, we have a continuous map $\Lambda : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and constants $c_5 > 0$ and β depending only on p_+ , p_* so that

$$\|u\|_\infty \leq c_5 \|u\|_q^\beta \Lambda(\|u\|_{p^*}), \quad (42)$$

with $\beta = \frac{p^* - 1}{(p_+ - 1)(1 + \varepsilon)(1 + \gamma)}$, $\Lambda(\sigma) = \left[\sigma^{\varepsilon p^*} \max_{1 \leq j \leq N} (\sigma^{p_+ - p_j}) \right]^{\frac{1}{(1 + \varepsilon)(1 + \gamma)}}$.

Thus, from relation (42), we deduce

$$\|u\|_\infty^{1 - \beta(1 - \frac{p^*}{q})} \leq c_5 \|u\|_{p^*}^{\beta \frac{p^*}{q}} \Lambda(\|u\|_{p^*}) \text{ for } u \neq 0. \quad (43)$$

But the number $\kappa = 1 - \beta \left(1 - \frac{p^*}{q} \right) = 0$, so relation (43) implies that there is a number $\tau_0 > 0$ depending only p_j , p^* such that $\|u\|_{p^*} \geq \tau_0 > 0$. $\diamond$

Acknowledgment : The authors would like to thank the referee for his/her valuable comments.

References

- [1] Aubin T, *Problèmes isopérimétriques et espaces de Sobolev*. J. Differential Geometry (1976) 11, no. 4, 573–598.
- [2] Brezis H, Lieb E, *A relation between pointwise convergence of functions and convergence of functionals*. Proc. American Math. Soc. (1983) 88 486–490.
- [3] El Hamidi A, Rakotoson J. M, *Compactness and quasilinear problems with critical exponents*. Diff. Int. Equ. (2005) Vol 18 1201–1220.
- [4] Fragala I, Gazzola F, Kawohl B, *Existence and nonexistence results for anisotropic quasilinear elliptic equation*. Ann. I. H. Poincaré (2004) AN 21 715–734.
- [5] Lions P.L, *The concentration-compactness principle in the calculus of variations. The limit case, part1*. Rev. Matemática Iberoamericana (1985) Vol 1 No 1, 145–201. *part 2* Rev. Matemática Iberoamericana (1985) Vol 1 No 2, 45–121.
- [6] Nikol'skii S.M, *On imbedding, continuation and approximation theorems for differentiable functions of several variables*. Russian Math. Surv. (1961) 16 55–104.
- [7] Rakosnik J, *Some remarks to anisotropic Sobolev spaces I*. Beiträge zur Analysis (1979) 13 55–68.
- [8] Rakosnik J, *Some remarks to anisotropic Sobolev spaces II*. Beiträge zur Analysis (1981) 15 127–140.
- [9] Talenti G, *Best constant in Sobolev inequality*. Ann. Mat. Pura Appl. (1976) (4) 110 353–372.
- [10] Troisi M, *Teoremi di inclusione per spazi di Sobolev non isotropi*. Ricerche Mat. (1969) 18 3–24.

- [11] Ven'-tuan L, *On embedding theorems for spaces of functions with partial derivatives of various degrees of summability*. Vestnik Leningrad Univ. (1961) 16, pp. 23-37 (in Russian) .
- [12] Willem M, *Minimax Theorems*. Birkhäuser, (1996).
- [13] Yamabe H, *On a deformation of Riemannian structures on compact manifolds*. Osaka Math. J. (1960) 12 21–37.

Liste des prépublications

- 99-1 Monique Jeanblanc et Nicolas Privault. A complete market model with Poisson and Brownian components. A paraître dans *Proceedings of the Seminar on Stochastic Analysis, Random Fields and Applications*, Ascona, 1999.
- 99-2 Laurence Cherfils et Alain Miranville. Generalized Cahn-Hilliard equations with a logarithmic free energy. A paraître dans *Revista de la Real Academia de Ciencias*.
- 99-3 Jean-Jacques Prat et Nicolas Privault. Explicit stochastic analysis of Brownian motion and point measures on Riemannian manifolds. *Journal of Functional Analysis* **167** (1999) 201-242.
- 99-4 Changgui Zhang. Sur la fonction q -Gamma de Jackson. A paraître dans *Aequationes Math.*
- 99-5 Nicolas Privault. A characterization of grand canonical Gibbs measures by duality. A paraître dans *Potential Analysis*.
- 99-6 Guy Wallet. La variété des équations surstables. A paraître dans *Bulletin de la Société Mathématique de France*.
- 99-7 Nicolas Privault et Jiang-Lun Wu. Poisson stochastic integration in Hilbert spaces. *Annales Mathématiques Blaise Pascal*, **6** (1999) 41-61.
- 99-8 Augustin Fruchard et Reinhard Schäfke. Sursabilité et résonance.
- 99-9 Nicolas Privault. Connections and curvature in the Riemannian geometry of configuration spaces. *C. R. Acad. Sci. Paris, Série I* **330** (2000) 899-904.
- 99-10 Fabienne Marotte et Changgui Zhang. Multisommabilité des séries entières solutions formelles d'une équation aux q -différences linéaire analytique. A paraître dans *Annales de l'Institut Fourier*, 2000.
- 99-11 Knut Aase, Bernt Øksendal, Nicolas Privault et Jan Ubøe. White noise generalizations of the Clark-Hausmann-Ocone theorem with application to mathematical finance. *Finance and Stochastics*, **4** (2000) 465-496.
- 00-01 Eric Benoît. Canards en un point pseudo-singulier nœud. A paraître dans *Bulletin de la Société Mathématique de France*.
- 00-02 Nicolas Privault. Hypothesis testing and Skorokhod stochastic integration. *Journal of Applied Probability*, **37** (2000) 560-574.
- 00-03 Changgui Zhang. La fonction θ de Jacobi et la sommabilité des séries entières q -Gevrey, I. *C. R. Acad. Sci. Paris, Série I* **331** (2000) 31-34.
- 00-04 Guy Wallet. Déformation topologique par changement d'échelle.
- 00-05 Nicolas Privault. Quantum stochastic calculus for the uniform measure and Boolean convolution. A paraître dans *Séminaire de Probabilités XXXV*.
- 00-06 Changgui Zhang. Sur les fonctions q -Bessel de Jackson.
- 00-07 Laure Coutin, David Nualart et Ciprian A. Tudor. Tanaka formula for the fractional Brownian motion. A paraître dans *Stochastic Processes and their Applications*.
- 00-08 Nicolas Privault. On logarithmic Sobolev inequalities for normal martingales. *Annales de la Faculté des Sciences de Toulouse* **9** (2000) 509-518.
- 01-01 Emanuelle Augeraud-Veron et Laurent Augier. Stabilizing endogenous fluctuations by fiscal policies; Global analysis on piecewise continuous dynamical systems. A paraître dans *Studies in Nonlinear Dynamics and Econometrics*
- 01-02 Delphine Boucher. About the polynomial solutions of homogeneous linear differential equations depending on parameters. A paraître dans *Proceedings of the 1999 International Symposium on Symbolic and Algebraic Computation: ISSAC 99, Sam Dooley Ed., ACM, New York 1999*.
- 01-03 Nicolas Privault. Quasi-invariance for Lévy processes under anticipating shifts.
- 01-04 Nicolas Privault. Distribution-valued iterated gradient and chaotic decompositions of Poisson jump times functionals.
- 01-05 Christian Houdré et Nicolas Privault. Deviation inequalities: an approach via covariance representations.
- 01-06 Abdallah El Hamidi. Remarques sur les sentinelles pour les systèmes distribués
- 02-01 Eric Benoît, Abdallah El Hamidi et Augustin Fruchard. On combined asymptotic expansions in singular perturbation.

- 02-02 Rachid Bebbouchi et Eric Benoît. Equations différentielles et familles bien nées de courbes planes.
- 02-03 Abdallah El Hamidi et Gennady G. Laptev. Nonexistence of solutions to systems of higher-order semilinear inequalities in cone-like domains.
- 02-04 Hassan Lakhel, Youssef Ouknine, et Ciprian A. Tudor. Besov regularity for the indefinite Skorohod integral with respect to the fractional Brownian motion: the singular case.
- 02-05 Nicolas Privault et Jean-Claude Zambrini. Markovian bridges and reversible diffusions with jumps.
- 02-06 Abdallah El Hamidi et Gennady G. Laptev. Existence and Nonexistence Results for Reaction-Diffusion Equations in Product of Cones.
- 02-07 Guy Wallet. Nonstandard generic points.
- 02-08 Gilles Bailly-Maitre. On the monodromy representation of polynomials.
- 02-09 Abdallah El Hamidi. Necessary conditions for local and global solvability of nondiagonal degenerate systems.
- 02-10 Abdallah El Hamidi et Amira Obeid. Systems of Semilinear higher order evolution inequalities on the Heisenberg group.
- 03-01 Abdallah El Hamidi et Gennady G. Laptev. Non existence de solutions d'inéquations semilinéaires dans des domaines coniques.
- 03-02 Eric Benoît et Marie-Joëlle Rochet. A continuous model of biomass size spectra governed by predation and the effects of fishing on them.
- 03-03 Catherine Stenger: On a conjecture of Wolfgang Wasow concerning the nature of turning points.
- 03-04 Christian Houdré et Nicolas Privault. Surface measures and related functional inequalities on configuration spaces.
- 03-05 Abdallah El Hamidi et Mokhtar Kirane. Nonexistence results of solutions to systems of semilinear differential inequalities on the Heisenberg group.
- 03-06 Uwe Franz, Nicolas Privault et René Schott. Non-Gaussian Malliavin calculus on real Lie algebras.
- 04-01 Abdallah El Hamidi. Multiple solutions to a nonlinear elliptic equation involving Paneitz type operators.
- 04-02 Mohamed Amara, Amira Obeid et Guy Vallet. Relaxed formulation and existence result of the degenerated elliptic small disturbance model.
- 04-03 Hippolyte d'Albis et Emmanuelle Augeraud-Veron. Competitive Growth in a Life-cycle Model: Existence and Dynamics
- 04-04 Sadjia Aït-Mokhtar: Third order differential equations with fixed critical points.
- 04-05 Mokhtar Kirane et Nasser-eddine Tatar. Asymptotic Behavior for a Reaction Diffusion System with Unbounded Coefficients.
- 04-06 Mokhtar Kirane, Eric Nabana et Stanislav I. Pohozaev. Nonexistence of Global Solutions to an Elliptic Equation with a Dynamical Boundary Condition.
- 04-07 Khaled M. Furati, Nasser-eddine Tatar and Mokhtar Kirane. Existence and asymptotic behavior for a convection Problem.
- 04-08 José Alfredo López-Mimbela et Nicolas Privault. Blow-up and stability of semilinear PDE's with gamma generator.
- 04-09 Abdallah El Hamidi. Multiple solutions with changing sign energy to a nonlinear elliptic equation.
- 04-10 Sadjia Aït-Mokhtar: A singularly perturbed Riccati equation.
- 04-11 Mohamed Amara, Amira Obeid et Guy Vallet. Weighted Sobolev spaces for a degenerated nonlinear elliptic equation.
- 04-12 Abdallah El Hamidi. Existence results to elliptic systems with nonstandard growth conditions.
- 04-13 Eric Edo et Jean-Philippe Furter: Some families of polynomial automorphisms.
- 04-14 Laurence Cherfils et Yavdat Il'yasov. On the stationary solutions of generalized reaction diffusion equations with p & q - Laplacian.
- 04-15 Jean-Christophe Breton et Youri Davydov. Local limit theorem for supremum of an empirical processes for i.i.d. random variables.
- 04-16 Jean-Christophe Breton, Christian Houdré et Nicolas Privault. Dimension free and infinite variance tail estimates on Poisson space.

- 04-17 Abdallah El Hamidi et Gennady G. Laptev. Existence and nonexistence results for higher-order semilinear evolution inequalities with critical potential.
- 05-01 Mokhtar Kirane et Nasser-eddine Tatar. Nonexistence of Solutions to a Hyperbolic Equation with a Time Fractional Damping.
- 05-02 Mokhtar Kirane et Yamina Laskri. Nonexistence of Global Solutions to a Hyperbolic Equation with a Time Fractional Damping.
- 05-03 Mokhtar Kirane, Yamina Laskri et Nasser-eddine Tatar. Critical Exponents of Fujita Type for Certain Evolution Equations and Systems with Spatio-Temporal Fractional Derivatives.
- 05-04 Abdallah El Hamidi et Jean-Michel Rakotoson. Compactness and quasilinear problems with critical exponents
- 05-05 Claudianor O. Alves et Abdallah El Hamidi. Nehari manifold and existence of positive solutions to a class of quasilinear problems.
- 05-06 Khalid Adriouch et Abdallah El Hamidi. The Nehari manifold for systems of nonlinear elliptic equations.
- 05-07 Eric Benoît. Equation fonctionnelle: Transport et convolution.
- 05-08 Jean-Philippe Furter et Stefan Maubach. Locally Finite Polynomial Endomorphisms.
- 05-09 Thomas Forget. Solutions canards en des points tournants dégénérés.
- 05-10 José Alfredo López-Mimbela et Nicolas Privault. Critical Exponents for Semilinear PDEs with Bounded Potentials.
- 06-01 Aldéric Joulin. On maximal inequalities for stable stochastic integrals.
- 06-02 Aldéric Joulin. On local Poisson-type deviation inequalities for curved continuous time Markov chains, with applications to birth-death processes.
- 06-03 Abdallah El Hamidi et Jean-Michel Rakotoson. On a perturbed anisotropic equation. *Ricerche Di Matematica*, volume 55 No 1 (2006) 55–69.
- 06-04 Khalid Adriouch et Abdallah El Hamidi. On local compactness in quasilinear elliptic problems. A paraître dans *Diff. Integ. Equ.*
- 06-05 Jean-Christophe Breton. Convergence in variation of the joint laws of multiple stable stochastic integrals.
- 06-06 Laurence Cherfils et Alain Miranville. Some remarks on the asymptotic behavior of the Caginalp system with singular potentials.
- 06-07 Jean-Philippe Furter. Quasi-locally Finite Polynomial Endomorphisms.
- 06-08 Jean-Philippe Furter. Jet Groups.
- 06-09 Jean-Philippe Furter. Fat Points Embeddings.
- 06-10 Abdallah El Hamidi et Jean-Michel Rakotoson. Fonctions minimales pour des inégalités de Sobolev anisotropiques. À paraître dans *Ann. Int. H. Poincaré*, "Analyse non linéaire".